

Lectures on rough paths

F. CARAVENNA

M. GUBINELLI

L. ZAMBOTTI

(version du 14/05/25)

TABLE OF CONTENTS

INTRODUCTION	9
Notation	10
I. Rough Equations	11
1. THE SEWING BOUND	15
1.1. Controlled differential equation	15
1.2. Controlled difference equation	15
1.3. Some useful function spaces	17
1.4. Local uniqueness of solutions	19
1.5. The Sewing bound	20
1.6. End of proof of uniqueness	22
1.7. Weighted norms	23
1.8. A discrete Sewing Bound	25
1.9. Extra (to be completed)	27
2. DIFFERENCE EQUATIONS: THE YOUNG CASE	29
2.1. Summary	29
2.2. Set-up	30
2.3. A priori estimates	31
2.4. Uniqueness	33
2.5. Continuity of the solution map	36
2.6. Euler scheme and local/global existence	39
First part: globally Hölder case.	40
Second part: locally Lipschitz case.	41
2.7. Error estimate in the Euler scheme	42
2.8. Extra: a value for $\hat{\tau}$	43
3. DIFFERENCE EQUATIONS: THE ROUGH CASE	45
3.1. Enhanced Taylor expansion	45
3.2. Rough paths	47
3.3. Rough difference equations	49
3.4. Set-up	50
3.5. A priori estimates	51
3.6. Uniqueness	53
3.7. Continuity of the solution map	56
3.8. Global existence and uniqueness	60
3.9. Milstein scheme and local existence	60

4. STOCHASTIC DIFFERENTIAL EQUATIONS	63
4.1. Local expansion of stochastic integrals	64
4.2. Brownian rough path and SDEs	65
4.3. SDE with a drift	66
4.4. A refined Kolmogorov criterion	68
4.5. Proof of Theorem 4.3	71
5. WONG-ZAKAI	77
5.1. Itô versus Stratonovich	77
5.2. Wong-Zakai	78
II. Rough Integration	85
6. THE SEWING LEMMA	89
6.1. Local approximation	89
6.2. A general problem	90
6.3. An algebraic look	91
6.4. Enters analysis: the Sewing Lemma	91
6.5. The Sewing Map	93
6.6. Proof of the Sewing Lemma	94
7. THE YOUNG INTEGRAL	97
7.1. Construction of the Young integral	97
7.2. Itô versus Young	99
7.3. Integral formulation of Young equations	99
7.4. Properties of the Young integral	101
7.5. More on Hölder spaces	103
7.6. Uniqueness of the Young integral	104
7.7. Two technical proofs	106
8. ROUGH PATHS	109
8.1. Integral beyond Young	110
8.2. A negative result	111
8.3. A choice	112
8.4. One-dimensional rough paths	114
8.5. The vector case	115
8.6. Distance on rough paths	116
8.7. Canonical rough paths for $\alpha > \frac{1}{2}$	117
8.8. Lack of continuity	119
8.9. The paraintegral	120
9. ROUGH INTEGRATION	125
9.1. Controlled paths	125
9.2. The rough integral	126

9.3. Continuity properties of the rough integral	127
9.4. Stochastic and rough integrals	130
10. ROUGH INTEGRAL EQUATIONS	133
10.1. Localization argument	134
10.2. Invariance	134
10.3. Local Lipschitz continuity	136
10.4. Contraction	138
11. GEOMETRIC ROUGH PATHS	141
11.1. Geometric rough paths	141
11.2. The Stratonovich rough path	143
11.3. Non-geometric rough paths	144
11.4. Pure area rough paths	145
11.5. Doss-Sussmann	146
11.6. Lack of continuity (again)	148
11.7. The rough integral in the geometric case	149
12. ALGEBRA	151
12.1. A non-commutative group	151
12.2. Shuffle group	152
12.3. Unordered times	152
12.4. An example: the Brownian case	154
12.5. Sewing for unordered times	155
13. NEW TOPICS	155
BIBLIOGRAPHY	159

INTRODUCTION

The problem of interest in this book is the study of differential equations driven by *irregular functions* (more specifically: continuous but not differentiable). This will be achieved through the powerful and elegant theory of *rough paths*. A key motivation comes from stochastic differential equations driven by Brownian motion, but the goal is to develop a general theory which does not rely on probability.

We will focus on *controlled differential equations*: given a *driving path* $X = (X_t)_{t \geq 0}$ and a function $\sigma(\cdot)$, we look for a *solution* $Z = (Z_t)_{t \geq 0}$ of

$$\dot{Z}_t = \sigma(Z_t) \dot{X}_t. \quad (1)$$

The challenge is to make sense of this equation *when X is not differentiable*. We consider in particular the case when X is a generic α -Hölder continuous path, i.e. it satisfies $|X_t - X_s| = O(|t - s|^\alpha)$ for some $\alpha \in]0, 1[$.

The basic idea is to reformulate the ill-posed differential equation (1) in ways that do not contain the derivative $\dot{X}$. In Part I of this book (Chapters 1 to 4) we rewrite the differential equation (1) as a *finite difference equation* on some interval $[0, T]$:

$$Z_t - Z_s = \sigma(Z_s) (X_t - X_s) + \{\cdots\} + o(t - s) \quad \text{for } 0 \leq s < t \leq T, \quad (2)$$

where $\{\cdots\}$ denote suitable additional terms (to be described below). For such a difference equation we prove *well-posedness*, i.e. existence, uniqueness, regularity of solutions and continuous dependence on initial data, when X is a generic α -Hölder path with $\alpha > \frac{1}{3}$ and $\sigma(\cdot)$ is a sufficiently regular function. More precisely:

- in Chapter 1 we introduce our key tool, that we call the *Sewing Bound*¹;
- in Chapter 2 we consider the so-called *Young case* $\alpha > \frac{1}{2}$: we prove well-posedness for the difference equation (2) where we simply take $\{\cdots\} = 0$;
- in Chapter 3 we consider the so-called *Rough case* $\frac{1}{3} < \alpha \leq \frac{1}{2}$: in this regime we need to *enrich the path X* with a notion of *iterated integral* $\int X_u dX_u$, leading to a so-called *rough path*, then we prove well-posedness for the difference equation (2) for a natural additional term $\{\cdots\} \neq 0$;
- in Chapter 4 we apply the theory of Chapter 3 when $X = B$ is a typical path of *Brownian motion*: the solution of the difference equation (2) coincides with the solution of the *stochastic differential equation* $dZ_t = \sigma(Z_t) dB_t$ in the Itô or Stratonovich sense, depending on the choice of iterated integral $\int B_u dB_u$.

1. This may be viewed as “half” of the celebrated *Sewing Lemma*, to be discussed in Chapter 6.

Altogether, we show that the difference equation (2) driven by a rough path (i.e. a Hölder continuous path X enriched by an iterated integral) provides an elementary yet powerful reformulation of the ill-posed differential equation (1), which sheds a “pathwise light” on stochastic differential equations driven by Brownian motion.²

Part II of this book is devoted to the alternative (and possibly more customary) reformulation of the differential equation (1) as an integral equation:

$$Z_t = Z_0 + \mathcal{I}_t(\sigma(Z), X), \quad \text{where} \quad \mathcal{I}_t(Y, X) = \int_0^t Y_s \dot{X}_s ds. \quad (3)$$

The natural strategy to solve it (for given initial datum Z_0) is via the following steps:

- define a notion of *integral* $\mathcal{I}_t(Y, X)$ with respect to a non-differentiable X , for a suitable class $\mathcal{H}$ of integrands $Y = (Y_s)_{s \geq 0}$;
- show that the maps $Y \mapsto (\mathcal{I}_t(Y, X))_{t \geq 0}$ and $Z \mapsto \sigma(Z)$ are *continuous* with respect to a suitable metric on the space $\mathcal{H}$ of integrands;
- solve (3) as a *fixed point equation*, by showing that for small time horizon $T > 0$ the map $Z \mapsto \Phi(Z) := (\mathcal{I}_t(\sigma(Z), X))_{t \in [0, T]}$ is a contraction.

This is indeed the standard strategy to solve stochastic differential equations driven by Brownian motion $X = B$, in which case $\mathcal{I}_t(Y, B) = \int_0^t Y_s dB_s$ makes sense as a stochastic integral (e.g. Ito or Stratonovich).

We show that the same strategy can be applied in the theory of *rough paths*: one can define a notion of *rough integral* $\mathcal{I}_t(Y, X)$ for a wide class of integrands Y , known as *controlled paths*, and then solve the equation (3) by fixed point arguments. This is the approach in the book by Hairer and Friz **[Insert reference]**.

To be completed...

NOTATION

We fix a time horizon $T > 0$ and two dimensions $k, d \in \mathbb{N}$. We use “path” as a synonymous of “function defined on $[0, T]$ ” with values in $\mathbb{R}^d$. We denote by $|\cdot|$ the Euclidean norm.

The space of linear maps from $\mathbb{R}^d$ to $\mathbb{R}^k$ is denoted by $\mathcal{L}(\mathbb{R}^d, \mathbb{R}^k) = \mathbb{R}^k \otimes (\mathbb{R}^d)^*$. We can identify it with the space $\mathbb{R}^{k \times d}$ of $k \times d$ real matrices, that we equip with the Hilbert-Schmidt norm $|\cdot|$ (i.e. the Euclidean norm on $\mathbb{R}^{k \times d}$). Note that for a linear map $A \in \mathcal{L}(\mathbb{R}^d, \mathbb{R}^k)$ and a vector $v \in \mathbb{R}^d$ we have $|A v| \leq |A| |v|$.

2. The restriction to α -Hölder paths X with $\alpha > \frac{1}{3}$ is made for simplicity: the theory can be extended to any $\alpha > 0$ (for $\frac{1}{n+1} < \alpha \leq \frac{1}{n}$ with $n \in \mathbb{N}$ one needs to enrich the path X with n -order iterated integrals).

Part I

Rough Equations

CHAPTER 1

THE SEWING BOUND

This first chapter is dedicated to an elementary but fundamental tool, the *Sewing Bound*, that will be applied extensively throughout the book. It is a general Hölder-type bound for functions of two real variables that can be understood by itself, see Theorem 1.9 below. To provide motivation, we present it as a natural a priori estimate for solutions of differential equations.

1.1. CONTROLLED DIFFERENTIAL EQUATION

Consider the following *controlled ordinary differential equation (ODE)*: given a continuously differentiable path $X: [0, T] \rightarrow \mathbb{R}^d$ and a continuous function $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$, we look for a differentiable path $Z: [0, T] \rightarrow \mathbb{R}^k$ such that

$$\dot{Z}_t = \sigma(Z_t) \dot{X}_t, \quad t \in [0, T]. \quad (1.1)$$

By the fundamental theorem of calculus, this is equivalent to

$$Z_t = Z_0 + \int_0^t \sigma(Z_s) \dot{X}_s ds, \quad t \in [0, T]. \quad (1.2)$$

In the special case $k = d = 1$ and when $\sigma(x) = \lambda x$ is linear (with $\lambda \in \mathbb{R}$), we have the explicit solution $Z_t = z_0 \exp(\lambda (X_t - X_0))$, which has the interesting property of being well-defined also when X is non differentiable.

For any dimensions $k, d \in \mathbb{N}$, if we assume that $\sigma(\cdot)$ is Lipschitz, classical results in the theory of ODEs guarantee that *equation (1.1)-(1.2) is well-posed for any continuously differentiable path X* , namely for any $Z_0 \in \mathbb{R}^k$ there is one and only one solution Z (with no explicit formula, in general).

Our aim is to extend such a well-posedness result to a setting where X is *continuous but not differentiable* (also in cases where $\sigma(\cdot)$ may be non-linear). Of course, to this purpose it is first necessary to provide a generalized formulation of (1.1)-(1.2) where the derivative of X does not appear.

1.2. CONTROLLED DIFFERENCE EQUATION

Let us still suppose that X is continuously differentiable. We deduce by (1.1)-(1.2) that for $0 \leq s \leq t \leq T$

$$Z_t - Z_s = \sigma(Z_s) (X_t - X_s) + \int_s^t (\sigma(Z_u) - \sigma(Z_s)) \dot{X}_u du, \quad (1.3)$$

which implies that Z satisfies the following *controlled difference equation*:

$$Z_t - Z_s = \sigma(Z_s)(X_t - X_s) + o(t - s), \quad 0 \leq s \leq t \leq T, \quad (1.4)$$

because $u \mapsto \sigma(Z_u)$ is continuous and $u \mapsto \dot{X}_u$ is (continuous, hence) bounded on $[0, T]$.

Remark 1.1. (UNIFORMITY) Whenever we write $o(t - s)$, as in (1.4), we always mean *uniformly for* $0 \leq s \leq t \leq T$, i.e.

$$\forall \varepsilon > 0 \exists \delta > 0: \quad 0 \leq s \leq t \leq T, \quad t - s \leq \delta \quad \text{implies} \quad |o(t - s)| \leq \varepsilon(t - s). \quad (1.5)$$

This will be implicitly assumed in the sequel.

Let us make two simple observations.

- If X is continuously differentiable we deduced (1.4) from (1.1), but we can easily deduce (1.1) from (1.4): in other terms, the two equations (1.1) and (1.4) are *equivalent*.
- If X is *not* continuously differentiable, equation (1.4) is still *meaningful*, unlike equation (1.1) which contains explicitly $\dot{X}$.

For these reasons, henceforth we focus on the difference equation (1.4), which provides a generalized formulation of the differential equation (1.1) when X is continuous but not necessarily differentiable.

The problem is now to prove *well-posedness* for the difference equation (1.4). We are going to show that this is possible assuming a suitable *Hölder regularity* on X , but non trivial ideas are required. In this chapter we illustrate some key ideas, showing how to prove uniqueness of solutions via *a priori estimates* (existence of solutions will be studied in the next chapters). We start from a basic result, which ensures the continuity of solutions; more precise result will be obtained later.

LEMMA 1.2. (CONTINUITY OF SOLUTIONS) *Let X and σ be continuous. Then any solution Z of (1.4) is a continuous path, more precisely it satisfies*

$$|Z_t - Z_s| \leq C |X_t - X_s| + o(t - s), \quad 0 \leq s \leq t \leq T, \quad (1.6)$$

for a suitable constant $C < \infty$ which depends on Z .

Proof. Relation (1.6) follows by (1.4) with $C := \|\sigma(Z)\|_\infty = \sup_{0 \leq t \leq T} |\sigma(Z_t)|$, renaming $|o(t - s)|$ as $o(t - s)$. We only have to prove that $C < \infty$. Since σ is continuous by assumption, it is enough to show that Z is *bounded*.

Since $o(t - s)$ is uniform, see (1.5), we can fix $\bar{\delta} > 0$ such that $|o(t - s)| \leq 1$ for all $0 \leq s \leq t \leq T$ with $|t - s| \leq \bar{\delta}$. It follows that Z is bounded in any interval $[\bar{s}, \bar{t}]$ with $|\bar{t} - \bar{s}| \leq \bar{\delta}$, because by (1.4) we can bound

$$\sup_{t \in [\bar{s}, \bar{t}]} |Z_t| \leq |Z_{\bar{s}}| + |\sigma(Z_{\bar{s}})| \sup_{t \in [\bar{s}, \bar{t}]} |X_t - X_{\bar{s}}| + 1 < \infty.$$

We conclude that Z is bounded in the whole interval $[0, T]$, because we can write $[0, T]$ as a finite union of intervals $[\bar{s}, \bar{t}]$ with $|\bar{t} - \bar{s}| \leq \bar{\delta}$. $\square$

Remark 1.3. (COUNTEREXAMPLES) The weaker requirement that (1.4) holds for any fixed $s \in [0, T]$ as $t \downarrow s$ is not enough for our purposes, since in this case Z needs not be continuous. An easy counterexample is the following: given any continuous path $X: [0, 2] \rightarrow \mathbb{R}$, we define $Z: [0, 2] \rightarrow \mathbb{R}$ by

$$Z_t := \begin{cases} X_t & \text{if } 0 \leq t < 1, \\ X_t + 1 & \text{if } 1 \leq t \leq 2. \end{cases}$$

Note that $Z_t - Z_s = X_t - X_s$ when either $0 \leq s \leq t < 1$ or $1 \leq s \leq t \leq 2$, hence Z satisfies the difference equation (1.4) with $\sigma(\cdot) \equiv 1$ for any fixed $s \in [0, 2)$ as $t \downarrow s$, but not uniformly for $0 \leq s \leq t \leq 2$, since Z is discontinuous at $t = 1$.

For another counterexample, which is even unbounded, consider

$$Z_t := \begin{cases} \frac{1}{1-t} & \text{if } 0 \leq t < 1, \\ 0 & \text{if } 1 \leq t \leq 2, \end{cases}$$

which satisfies (1.4) as $t \downarrow s$ for any fixed $s \in [0, 2]$, for $X_t \equiv t$ and $\sigma(z) = z^2$.

1.3. SOME USEFUL FUNCTION SPACES

For $n \geq 1$ we define the simplex

$$[0, T]_{\leq}^n := \{(t_1, \dots, t_n): 0 \leq t_1 \leq \dots \leq t_n \leq T\} \quad (1.7)$$

(note that $[0, T]_{\leq}^1 = [0, T]$). We then write $C_n = C([0, T]_{\leq}^n, \mathbb{R}^k)$ as a shorthand for the space of continuous functions from $[0, T]_{\leq}^n$ to $\mathbb{R}^k$:

$$C_n := C([0, T]_{\leq}^n, \mathbb{R}^k) := \{F: [0, T]_{\leq}^n \rightarrow \mathbb{R}^k: F \text{ is continuous}\}. \quad (1.8)$$

We are going to work with functions of one (f_s), two (F_{st}) or three (G_{sut}) ordered variables in $[0, T]$, hence we focus on the spaces C_1, C_2, C_3 .

- On the spaces C_2 and C_3 we introduce a Hölder-like structure: given any $\eta \in (0, \infty)$, we define for $F \in C_2$ and $G \in C_3$

$$\|F\|_{\eta} := \sup_{0 \leq s < t \leq T} \frac{|F_{st}|}{(t-s)^{\eta}}, \quad \|G\|_{\eta} := \sup_{\substack{0 \leq s \leq u \leq t \leq T \\ s < t}} \frac{|G_{sut}|}{(t-s)^{\eta}}, \quad (1.9)$$

and we denote by C_2^{η} and C_3^{η} the corresponding function spaces:

$$C_2^{\eta} := \{F \in C_2: \|F\|_{\eta} < \infty\}, \quad C_3^{\eta} := \{G \in C_3: \|G\|_{\eta} < \infty\}, \quad (1.10)$$

which are Banach spaces endowed with the norm $\|\cdot\|_{\eta}$ (exercise).

- On the space C_1 of continuous functions $f: [0, T] \rightarrow \mathbb{R}^k$ we consider the usual Hölder structure. We first introduce the increment δf by

$$(\delta f)_{st} := f_t - f_s, \quad 0 \leq s \leq t \leq T, \quad (1.11)$$

and note that $\delta f \in C_2$ for any $f \in C_1$. Then, for $\alpha \in (0, 1]$, we define the classical space $\mathcal{C}^\alpha = \mathcal{C}^\alpha([0, T], \mathbb{R}^k)$ of α -Hölder functions

$$\mathcal{C}^\alpha := \left\{ f: [0, T] \rightarrow \mathbb{R}^k: \quad \|\delta f\|_\alpha = \sup_{0 \leq s < t \leq T} \frac{|f_t - f_s|}{(t - s)^\alpha} < \infty \right\} \quad (1.12)$$

(for $\alpha = 1$ it is the space of Lipschitz functions). Note that $\|\delta f\|_\alpha$ in (1.12) is consistent with (1.11) and (1.9).

Remark 1.4. (HÖLDER SEMI-NORM) We stress that $f \mapsto \|\delta f\|_\alpha$ is a semi-norm on $\mathcal{C}^\alpha$ (it vanishes on constant functions). The standard norm on $\mathcal{C}^\alpha$ is

$$\|f\|_{\mathcal{C}^\alpha} := \|f\|_\infty + \|\delta f\|_\alpha, \quad (1.13)$$

where we define the standard sup norm

$$\|f\|_\infty := \sup_{t \in [0, T]} |f_t|. \quad (1.14)$$

For $f: [0, T] \rightarrow \mathbb{R}^k$ we can bound $\|f\|_\infty \leq |f(0)| + T^\alpha \|\delta f\|_\alpha$ (see (1.40) below), hence

$$\|f\|_{\mathcal{C}^\alpha} \leq |f(0)| + (1 + T^\alpha) \|\delta f\|_\alpha. \quad (1.15)$$

This explains why it is often enough to focus on the semi-norm $\|\delta f\|_\alpha$.

Remark 1.5. (HÖLDER EXPONENTS) We only consider the Hölder space $\mathcal{C}^\alpha$ for $\alpha \in (0, 1]$ because for $\alpha > 1$ the only functions in $\mathcal{C}^\alpha$ are constant functions (note that $\|\delta f\|_\alpha < \infty$ for $\alpha > 1$ implies $\dot{f}_t = 0$ for every $t \in [0, T]$).

On the other hand, the spaces C_2^η and C_3^η in (1.10) are interesting for any exponent $\eta \in (0, \infty)$. For instance, the condition $\|F\|_\eta < \infty$ for a function $F \in C_2$ means that $|F_{st}| \leq C(t - s)^\eta$, which does not imply $F \equiv 0$ when $\eta > 1$ (unless $F = \delta f$ is the increment of some function $f \in C_1$).

In our results below we will have to assume that the non-linearity $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ belongs to classes of Hölder functions, in the following sense.

DEFINITION 1.6. Let $\gamma > 0$. A function $F: \mathbb{R}^k \rightarrow \mathbb{R}^N$ is said to be globally γ -Hölder (or globally of class $\mathcal{C}^\gamma$) if

1. for $\gamma \in (0, 1]$ we have

$$[F]_{\mathcal{C}^\gamma} := \sup_{x, y \in \mathbb{R}^k, x \neq y} \frac{|F(x) - F(y)|}{|x - y|^\gamma} < +\infty \quad (1.16)$$

2. for $\gamma \in (n, n + 1]$ and $n = \{1, 2, \dots\}$, F is n times continuously differentiable and

$$[D^{(n)}F]_{\mathcal{C}^\gamma} := \sup_{x, y \in \mathbb{R}^k, x \neq y} \frac{|D^{(n)}F(x) - D^{(n)}F(y)|}{|x - y|^{\gamma - n}} < +\infty$$

where $D^{(n)}$ is the n -fold differential of F .

Moreover $F: \mathbb{R}^k \rightarrow \mathbb{R}^N$ is said to be locally γ -Hölder (or locally of class $\mathcal{C}^\gamma$) if

- for $\gamma \in (0, 1]$ we have for all $R > 0$

$$\sup_{\substack{x, y \in \mathbb{R}^k, x \neq y \\ |x|, |y| \leq R}} \frac{|F(x) - F(y)|}{|x - y|^\gamma} < +\infty$$

- for $\gamma \in (n, n+1]$ and $n = \{1, 2, \dots\}$, F is n times continuously differentiable and

$$\sup_{\substack{x, y \in \mathbb{R}^k, x \neq y \\ |x|, |y| \leq R}} \frac{|D^{(n)}F(x) - D^{(n)}F(y)|}{|x - y|^{\gamma-n}} < +\infty.$$

We stress that in the previous definition we do not assume F or $D^{(n)}F$ to be bounded. The case $\gamma = 1$ corresponds to the classical *Lipschitz* condition.

1.4. LOCAL UNIQUENESS OF SOLUTIONS

We prove *uniqueness of solutions* for the controlled difference equation (1.4) when $X \in \mathcal{C}^\alpha$ is an Hölder path of exponent $\alpha > \frac{1}{2}$. For simplicity, we focus on the case when $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is a linear application: $\sigma \in (\mathbb{R}^k \otimes (\mathbb{R}^d)^*) \otimes (\mathbb{R}^k)^*$, and we write σZ instead of $\sigma(Z)$ (we discuss non linear $\sigma(\cdot)$ in Chapter 2).

THEOREM 1.7. (LOCAL UNIQUENESS OF SOLUTIONS, LINEAR CASE) *Fix a path $X: [0, T] \rightarrow \mathbb{R}^d$ in $\mathcal{C}^\alpha$, with $\alpha \in]\frac{1}{2}, 1]$, and a linear map $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$. If $T > 0$ is small enough (depending on X, α, σ), then for any $z_0 \in \mathbb{R}^k$ there is at most one path $Z: [0, T] \rightarrow \mathbb{R}^k$ with $Z_0 = z_0$ which solves the linear controlled difference equation (1.4), that is (recalling (1.11))*

$$\delta Z_{st} - (\sigma Z_s) \delta X_{st} = o(t - s), \quad 0 \leq s \leq t \leq T. \quad (1.17)$$

Proof. Suppose that we have two paths $Z, \bar{Z}: [0, T] \rightarrow \mathbb{R}^k$ satisfying (1.17) with $Z_0 = \bar{Z}_0$ and define $Y := Z - \bar{Z}$. Our goal is to show that $Y = 0$.

Let us introduce the function $R \in C_2 = C([0, T]_{\leq}^2, \mathbb{R}^k)$ defined by

$$R_{st} := \delta Y_{st} - (\sigma Y_s) \delta X_{st}, \quad 0 \leq s \leq t \leq T, \quad (1.18)$$

and note that by (1.17) and linearity we have

$$R_{st} = o(t - s). \quad (1.19)$$

Recalling (1.9), we can estimate

$$\|\delta Y\|_\alpha \leq |\sigma| \|Y\|_\infty \|\delta X\|_\alpha + \|R\|_\alpha,$$

and since $R_{st} = o(t - s) = o((t - s)^\alpha)$, we have $\|R\|_\alpha < +\infty$ and therefore $\|\delta Y\|_\alpha < +\infty$. Since $Y_0 = 0$, we can bound

$$\|Y\|_\infty \leq |Y_0| + \sup_{0 \leq t \leq T} |Y_t - Y_0| \leq T^\alpha \|\delta Y\|_\alpha.$$

Since $1 \leq T^\alpha (t-s)^{-\alpha}$ for $0 \leq s < t \leq T$, we can also bound

$$\|R\|_\alpha \leq T^\alpha \|R\|_{2\alpha},$$

so that

$$\|\delta Y\|_\alpha \leq T^\alpha (|\sigma| \|\delta Y\|_\alpha \|\delta X\|_\alpha + \|R\|_{2\alpha}).$$

Suppose we can prove that, for some constant $C = C(X, \alpha, \sigma) < \infty$,

$$\|R\|_{2\alpha} \leq C \|\delta Y\|_\alpha. \quad (1.20)$$

Then we obtain

$$\|\delta Y\|_\alpha \leq T^\alpha (|\sigma| \|\delta X\|_\alpha + C) \|\delta Y\|_\alpha.$$

If we fix T small enough, so that $T^\alpha (|\sigma| \|\delta X\|_\alpha + C) < 1$, we get $\|\delta Y\|_\alpha = 0$, hence $\delta Y \equiv 0$. This means that $Y_t = Y_s$ for all $s, t \in [0, T]$, and since $Y_0 = 0$ we obtain $Y \equiv 0$, namely our goal $Z \equiv \bar{Z}$. This completes the proof *assuming the estimate (1.20)* (where the hypothesis $\alpha > \frac{1}{2}$ will play a key role). $\square$

To actually complete the proof of Theorem 1.7, it remains to show that the inequality (1.20) holds. This is performed in the next two sections:

- in Section 1.5 we present a fundamental estimate, the *Sewing Bound*, which applies to *any function* $R_{st} = o(t-s)$ (recall (1.19));
- in Section 1.6 we apply the Sewing Bound to R_{st} in (1.18) and we prove the desired estimate (1.20) for $\alpha > \frac{1}{2}$ (see the assumptions of Theorem 1.7).

1.5. THE SEWING BOUND

Let us fix an arbitrary function $R \in C_2 = C([0, T]^2, \mathbb{R}^k)$ with $R_{st} = o(t-s)$. Our goal is to bound $|R_{ab}|$ for any given $0 \leq a < b \leq T$.

We first show that we can express R_{ab} via “Riemann sums” along partitions $\mathcal{P} = \{a = t_0 < t_1 < \dots < t_m = b\}$ of $[a, b]$. These are defined by

$$I_{\mathcal{P}}(R) := \sum_{i=1}^{\#\mathcal{P}} R_{t_{i-1}t_i}, \quad (1.21)$$

where we denote by $\#\mathcal{P} := m$ the number of intervals of the partition $\mathcal{P}$. Let us denote by $|\mathcal{P}| := \max_{1 \leq i \leq m} (t_i - t_{i-1})$ the *mesh* of $\mathcal{P}$.

LEMMA 1.8. (RIEMANN SUMS) *Given any $R \in C_2$ with $R_{st} = o(t-s)$, for any $0 \leq a < b \leq T$ and for any sequence $(\mathcal{P}_n)_{n \geq 0}$ of partitions of $[a, b]$ with vanishing mesh $\lim_{n \rightarrow \infty} |\mathcal{P}_n| = 0$ we have*

$$\lim_{n \rightarrow \infty} I_{\mathcal{P}_n}(R) = 0.$$

If furthermore $\mathcal{P}_0 = \{a, b\}$ is the trivial partition, then we can write

$$R_{ab} = \sum_{n=0}^{\infty} (I_{\mathcal{P}_n}(R) - I_{\mathcal{P}_{n+1}}(R)), \quad 0 \leq a < b \leq T. \quad (1.22)$$

Proof. Writing $\mathcal{P}_n = \{a = t_0^n < t_1^n < \dots < t_{\#\mathcal{P}_n}^n = b\}$, we can estimate

$$|I_{\mathcal{P}_n}(R)| \leq \sum_{i=1}^{\#\mathcal{P}_n} |R_{t_{i-1}^n t_i^n}| \leq \left\{ \max_{j=1, \dots, \#\mathcal{P}_n} \frac{|R_{t_{j-1}^n t_j^n}|}{(t_j^n - t_{j-1}^n)} \right\} \sum_{j=1}^{\#\mathcal{P}_n} (t_j^n - t_{j-1}^n),$$

hence $|I_{\mathcal{P}_n}(R)| \rightarrow 0$ as $n \rightarrow \infty$, because the final sum equals $b - a$ and the bracket vanishes (since $R_{st} = o(t - s)$ and $|\mathcal{P}_n| = \max_{1 \leq j \leq \#\mathcal{P}_n} (t_j^n - t_{j-1}^n) \rightarrow 0$).

We deduce relation (1.22) by the telescopic sum

$$I_{\mathcal{P}_0}(R) - I_{\mathcal{P}_N}(R) = \sum_{n=0}^{N-1} (I_{\mathcal{P}_n}(R) - I_{\mathcal{P}_{n+1}}(R)),$$

because $\lim_{N \rightarrow \infty} I_{\mathcal{P}_N}(R) = 0$ while $I_{\mathcal{P}_0}(R) = R_{ab}$ for $\mathcal{P}_0 = \{a, b\}$. $\square$

If we remove a single point t_i from a partition $\mathcal{P} = \{t_0 < t_1 < \dots < t_m\}$, we obtain a new partition $\mathcal{P}'$ for which, recalling (1.21), we can write

$$I_{\mathcal{P}'}(R) - I_{\mathcal{P}}(R) = R_{t_{i-1} t_{i+1}} - R_{t_{i-1} t_i} - R_{t_i t_{i+1}}. \quad (1.23)$$

The expression in the RHS deserves a name: given any two-variables function $F \in C_2$, we define its increment $\delta F \in C_3$ as the three-variables function

$$\delta F_{sut} := F_{st} - F_{su} - F_{ut}, \quad 0 \leq s \leq u \leq t \leq T. \quad (1.24)$$

We can then rewrite (1.23) as

$$I_{\mathcal{P}'}(R) - I_{\mathcal{P}}(R) = \delta R_{t_{i-1} t_i t_{i+1}}, \quad (1.25)$$

and recalling (1.9) we obtain the following estimate, for any $\eta > 0$:

$$|I_{\mathcal{P}'}(R) - I_{\mathcal{P}}(R)| \leq \|\delta R\|_{\eta} |t_{i+1} - t_{i-1}|^{\eta}. \quad (1.26)$$

We are now ready to state and prove the Sewing Bound.

THEOREM 1.9. (SEWING BOUND) *Given any $R \in C_2$ with $R_{st} = o(t - s)$, the following estimate holds for any $\eta \in (1, \infty)$ (recall (1.9)):*

$$\|R\|_{\eta} \leq K_{\eta} \|\delta R\|_{\eta} \quad \text{where} \quad K_{\eta} := (1 - 2^{1-\eta})^{-1}. \quad (1.27)$$

Proof. Fix $R \in C_2$ such that $\|\delta R\|_{\eta} < \infty$ for some $\eta > 1$ (otherwise there is nothing to prove). Also fix $0 \leq a < b \leq T$ and consider for $n \geq 0$ the dyadic partitions $\mathcal{P}_n := \{t_i^n := a + \frac{i}{2^n}(b-a) : 0 \leq i \leq 2^n\}$ of $[a, b]$. Since $\mathcal{P}_0 = \{a, b\}$ is the trivial partition, we can apply (1.22) to bound

$$|R_{ab}| \leq \sum_{n=0}^{\infty} |I_{\mathcal{P}_n}(R) - I_{\mathcal{P}_{n+1}}(R)|. \quad (1.28)$$

If we remove from $\mathcal{P}_{n+1}$ all the “odd points” t_{2j+1}^{n+1} , with $0 \leq j \leq 2^n - 1$, we obtain $\mathcal{P}_n$. Then, iterating relations (1.25)-(1.26), we have

$$\begin{aligned} |I_{\mathcal{P}_n}(R) - I_{\mathcal{P}_{n+1}}(R)| &\leq \sum_{j=0}^{2^n-1} |\delta R_{t_{2j}^{n+1} t_{2j+1}^{n+1} t_{2j+2}^{n+1}}| \\ &\leq 2^n \|\delta R\|_\eta \left(\frac{2(b-a)}{2^{n+1}} \right)^\eta \\ &= 2^{-(\eta-1)n} \|\delta R\|_\eta (b-a)^\eta. \end{aligned} \quad (1.29)$$

Plugging this into (1.28), since $\sum_{n=0}^{\infty} 2^{-(\eta-1)n} = (1 - 2^{1-\eta})^{-1}$, we obtain

$$|R_{ab}| \leq (1 - 2^{1-\eta})^{-1} \|\delta R\|_\eta (b-a)^\eta, \quad 0 \leq a < b \leq T, \quad (1.30)$$

which proves (1.27). $\square$

Remark 1.10. Recalling (1.11) and (1.24), we have defined linear maps

$$C_1 \xrightarrow{\delta} C_2 \xrightarrow{\delta} C_3 \quad (1.31)$$

which satisfy $\delta \circ \delta = 0$. Indeed, for any $f \in C_1$ we have

$$\delta(\delta f)_{sut} = (f_t - f_s) - (f_u - f_s) - (f_t - f_u) = 0.$$

Intuitively, $\delta F \in C_3$ measures how much a function $F \in C_2$ differs from being the increment δf of some $f \in C_1$, because $\delta F \equiv 0$ if and only if $F = \delta f$ for some $f \in C_1$ (it suffices to define $f_t := F_{0t}$ and to check that $\delta f_{st} = \delta F_{0st} + F_{st} = F_{st}$).

Remark 1.11. The assumption $R_{st} = o(t-s)$ in Theorem 1.9 cannot be avoided: if $R := \delta f$ for a non constant $f \in C_1$, then $\delta R = 0$ while $\|R\|_\eta > 0$.

1.6. END OF PROOF OF UNIQUENESS

In this section, we apply the Sewing Bound (1.27) to the function R_{st} defined in (1.18), in order to prove the estimate (1.20) for $\alpha > \frac{1}{2}$.

We first determine the increment δR through a simple and instructive computation: by (1.18), since $\delta(\delta Z) = 0$ (see Remark 1.10), we have

$$\begin{aligned} \delta R_{sut} &:= R_{st} - R_{su} - R_{ut} \\ &= (Y_t - Y_s) - (Y_u - Y_s) - (Y_t - Y_u) \\ &\quad - (\sigma Y_s)(X_t - X_s) + (\sigma Y_s)(X_u - X_s) + (\sigma Y_u)(X_t - X_u) \\ &= [\sigma(Y_u - Y_s)](X_t - X_u). \end{aligned} \quad (1.32)$$

Recalling (1.9), this implies

$$\|\delta R\|_{2\alpha} \leq |\sigma| \|\delta Y\|_\alpha \|\delta X\|_\alpha.$$

We next note that if $\alpha > \frac{1}{2}$ (as it is assumed in Theorem 1.7) we can apply the Sewing Bound (1.27) for $\eta = 2\alpha > 1$ to obtain

$$\|R\|_{2\alpha} \leq K_{2\alpha} \|\delta R\|_{2\alpha} \leq K_{2\alpha} |\sigma| \|\delta Y\|_\alpha \|\delta X\|_\alpha.$$

This is precisely our goal (1.20) with $C = C(X, \alpha, \sigma) := K_{2\alpha} |\sigma| \|\delta X\|_\alpha$.

Summarizing: thanks to the Sewing bound (1.27), we have obtained the estimate (1.20) and completed the proof of Theorem 1.7, showing uniqueness of solutions to the difference equation (1.4) for any $X \in \mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$. In the next chapters we extend this approach to non-linear $\sigma(\cdot)$ and to situations where $X \in \mathcal{C}^\alpha$ with $\alpha \leq \frac{1}{2}$.

Remark 1.12. For later purpose, let us record the computation (1.32) without σ : given any (say, real) paths X and Y , if

$$A_{st} = Y_s \delta X_{st}, \quad \forall 0 \leq s \leq t \leq T,$$

then

$$\delta A_{sut} = -\delta Y_{su} \delta X_{ut}, \quad \forall 0 \leq s \leq u \leq t \leq T. \quad (1.33)$$

1.7. WEIGHTED NORMS

We conclude this chapter defining *weighted versions* $\|\cdot\|_{\eta, \tau}$ of the norms $\|\cdot\|_\eta$ introduced in (1.9): given $F \in C_2$ and $G \in C_3$, we set for $\eta, \tau \in (0, \infty)$

$$\|F\|_{\eta, \tau} := \sup_{0 \leq s \leq t \leq T} \mathbb{1}_{\{0 < t-s \leq \tau\}} e^{-\frac{t}{\tau}} \frac{|F_{st}|}{(t-s)^\eta}, \quad (1.34)$$

$$\|G\|_{\eta, \tau} := \sup_{0 \leq s \leq u \leq t \leq T} \mathbb{1}_{\{0 < t-s \leq \tau\}} e^{-\frac{t}{\tau}} \frac{|G_{sut}|}{(t-s)^\eta}, \quad (1.35)$$

where C_2 and C_3 are the spaces of continuous functions from $[0, T]_{\leq}^2$ and $[0, T]_{\leq}^3$ to $\mathbb{R}^k$, see (1.8). Note that as $\tau \rightarrow \infty$ we recover the usual norms:

$$\|\cdot\|_\eta = \lim_{\tau \rightarrow \infty} \|\cdot\|_{\eta, \tau}. \quad (1.36)$$

Remark 1.13. (NORMS VS. SEMI-NORMS) While $\|\cdot\|_\eta$ is a norm, $\|\cdot\|_{\eta, \tau}$ is a norm for $\tau \geq T$ but *it is only a semi-norm for $\tau < T$* (for instance, $\|F\|_{\eta, \tau} = 0$ for $F \in C_2$ implies $F_{st} = 0$ only for $t-s \leq \tau$: no constraint is imposed on F_{st} for $t-s > \tau$).

However, if $F = \delta f$, that is $F_{st} = f_t - f_s$ for some $f \in C_1$, we have the equivalence

$$\|\delta f\|_{\eta, \tau} \leq \|\delta f\|_\eta \leq \left(1 + \frac{T}{\tau}\right) e^{\frac{T}{\tau}} \|\delta f\|_{\eta, \tau}. \quad (1.37)$$

The first inequality is clear. For the second one, given $0 \leq s < t \leq T$, we can write $s = t_0 < t_1 < \dots < t_N = t$ with $t_i - t_{i-1} \leq \tau$ and $N \leq 1 + \frac{T}{\tau}$ (for instance, we can consider $t_i = s + i \frac{t-s}{N}$ where $N := \lceil \frac{t-s}{\tau} \rceil$); we then obtain $\delta f_{st} = \sum_{i=1}^N \delta f_{t_{i-1}t_i}$ and $|\delta f_{t_{i-1}t_i}| \leq \|\delta f\|_{\eta, \tau} e^{t_i/\tau} (t_i - t_{i-1})^\eta \leq \|\delta f\|_{\eta, \tau} e^{T/\tau} (t-s)^\eta$, which yields (1.37).

Remark 1.14. (FROM LOCAL TO GLOBAL) The weighted semi-norms $\|\cdot\|_{\eta, \tau}$ will be useful to transform *local* results in *global* results. Indeed, using the standard norms $\|\cdot\|_\eta$ often requires the size $T > 0$ of the time interval $[0, T]$ to be *small*, as in Theorem 1.7, which can be annoying. Using $\|\cdot\|_{\eta, \tau}$ will allow us to *keep* $T > 0$ *arbitrary*, by choosing a sufficiently small $\tau > 0$.

Recalling the supremum norm $\|f\|_\infty$ of a function $f \in C_1$, see (1.14), we define the corresponding weighted version

$$\|f\|_{\infty,\tau} := \sup_{0 \leq t \leq T} e^{-\frac{t}{\tau}} |f_t|. \quad (1.38)$$

We stress that $\|\cdot\|_{\infty,\tau}$ is a norm equivalent to $\|\cdot\|_\infty$ for any $\tau > 0$, since

$$\|\cdot\|_{\infty,\tau} \leq \|\cdot\|_\infty \leq e^{\frac{T}{\tau}} \|\cdot\|_{\infty,\tau}. \quad (1.39)$$

Remark 1.15. (EQUIVALENT HÖLDER NORM) It follows by (1.37) and (1.39) that $\|\cdot\|_{\infty,\tau} + \|\cdot\|_{\alpha,\tau}$ is a norm equivalent to $\|\cdot\|_{C^\alpha} := \|\cdot\|_\infty + \|\cdot\|_\alpha$ on the space $\mathcal{C}^\alpha$ of Hölder functions, see Remark 1.4, for any $\tau > 0$.

We will often use the Hölder semi-norms $\|\delta f\|_\alpha$ and $\|\delta f\|_{\alpha,\tau}$ to bound the supremum norms $\|f\|_\infty$ and $\|f\|_{\infty,\tau}$, thanks to the following result.

LEMMA 1.16. (SUPREMUM-HÖLDER BOUND) For any $f \in C_1$ and $\eta \in (0, \infty)$

$$\|f\|_\infty \leq |f_0| + T^\eta \|\delta f\|_\eta, \quad (1.40)$$

$$\|f\|_{\infty,\tau} \leq |f_0| + 3(\tau \wedge T)^\eta \|\delta f\|_{\eta,\tau}, \quad \forall \tau > 0. \quad (1.41)$$

Proof. Let us prove (1.40): for any $f \in C_1$ and for $t \in]0, T]$ we have

$$|f_t| \leq |f_0| + |f_t - f_0| = |f_0| + t^\eta \frac{|f_t - f_0|}{t^\eta} \leq |f_0| + T^\eta \|\delta f\|_\eta.$$

The proof of (1.41) is slightly more involved. If $t \in]0, \tau \wedge T]$, then

$$e^{-\frac{t}{\tau}} |f_t| \leq |f_0| + t^\eta e^{-\frac{t}{\tau}} \frac{|f_t - f_0|}{t^\eta} \leq |f_0| + (\tau \wedge T)^\eta \|\delta f\|_{\eta,\tau},$$

which, in particular, implies (1.41) when $\tau \geq T$. When $\tau < T$, it remains to consider $\tau < t \leq T$: in this case, we define $N := \min \{n \in \mathbb{N} : n\tau \geq t\} \geq 2$ so that $\frac{t}{N} \leq \tau$. We set $t_k = k \frac{t}{N}$ for $k \geq 0$, so that $t_N = t$. Then

$$\begin{aligned} e^{-\frac{t}{\tau}} |f_t| &\leq |f_0| + \sum_{k=1}^N (t_k - t_{k-1})^\eta e^{-\frac{t-t_k}{\tau}} \left[e^{-\frac{t_k}{\tau}} \frac{|f_{t_k} - f_{t_{k-1}}|}{(t_k - t_{k-1})^\eta} \right] \\ &\leq |f_0| + (\tau \wedge T)^\eta \|\delta f\|_{\eta,\tau} \sum_{k=1}^N e^{-\frac{t-t_k}{\tau}}. \end{aligned}$$

By definition of N we have $(N-1)\tau < t$; since $\tau < t$ we obtain $N\tau < 2t$ and therefore $\frac{t}{N\tau} \geq \frac{1}{2}$. Since $t - t_k = (N-k) \frac{t}{N}$, renaming $\ell := N - k$ we obtain

$$\sum_{k=1}^N e^{-\frac{t-t_k}{\tau}} = \sum_{\ell=0}^{N-1} e^{-\ell \frac{t}{N\tau}} = \frac{1 - e^{-\frac{t}{N\tau}}}{1 - e^{-\frac{t}{N\tau}}} \leq \frac{1}{1 - e^{-\frac{1}{2}}} \leq 3.$$

The proof is complete. □

We finally show that the Sewing Bound (1.27) still holds if we replace $\|\cdot\|_\eta$ by $\|\cdot\|_{\eta,\tau}$, for any $\tau > 0$.

THEOREM 1.17. (WEIGHTED SEWING BOUND) *Given any $R \in C_2$ with $R_{st} = o(t-s)$, the following estimate holds for any $\eta \in (1, \infty)$ and $\tau > 0$:*

$$\|R\|_{\eta,\tau} \leq K_\eta \|\delta R\|_{\eta,\tau} \quad \text{where} \quad K_\eta := (1 - 2^{1-\eta})^{-1}. \quad (1.42)$$

Proof. Given $0 \leq a \leq b \leq T$, let us define

$$\|\delta R\|_{\eta,[a,b]} := \sup_{\substack{s,u,t \in [a,b]: \\ s \leq u \leq t, s < t}} \frac{|\delta R_{sut}|}{(t-s)^\eta}. \quad (1.43)$$

Following the proof of Theorem 1.9, we can replace $\|\delta R\|_\eta$ by $\|\delta R\|_{\eta,[a,b]}$ in (1.29) and in (1.30), hence we obtain $|R_{ab}| \leq K_\eta \|\delta R\|_{\eta,[a,b]} (b-a)^\eta$. Then for $b-a \leq \tau$ we can estimate

$$e^{-\frac{b}{\tau}} \frac{|R_{ab}|}{(b-a)^\eta} \leq e^{-\frac{b}{\tau}} K_\eta \|\delta R\|_{\eta,[a,b]} \leq K_\eta \|\delta R\|_{\eta,\tau},$$

and (1.42) follows taking the supremum over $0 \leq a \leq b \leq T$ with $b-a \leq \tau$. $\square$

1.8. A DISCRETE SEWING BOUND

We can prove a version of the Sewing Bound for functions $R = (R_{st})_{s < t \in \mathbb{T}}$ defined on a *finite set of points* $\mathbb{T} := \{0 = t_1 < \dots < t_{\#\mathbb{T}}\} \subseteq \mathbb{R}_+$ (this will be useful to construct solutions to difference equations via Euler schemes, see Sections 2.6 and 3.9). The condition $R_{st} = o(t-s)$ from Theorem 1.9 is now replaced by the requirement that R *vanishes on consecutive points of $\mathbb{T}$* , i.e. $R_{t_i t_{i+1}} = 0$ for all $1 \leq i < \#\mathbb{T}$.

We define versions $\|\cdot\|_{\eta,\tau}^\mathbb{T}$ of the norms $\|\cdot\|_{\eta,\tau}$ restricted on $\mathbb{T}$ for $\tau > 0$, recall (1.34)-(1.35):

$$\|A\|_{\eta,\tau}^\mathbb{T} := \sup_{\substack{0 \leq s < t \\ s,t \in \mathbb{T}}} \mathbb{1}_{\{0 < t-s \leq \tau\}} e^{-\frac{t}{\tau}} \frac{|A_{st}|}{|t-s|^\eta}, \quad (1.44)$$

$$\|B\|_{\eta,\tau}^\mathbb{T} := \sup_{\substack{0 \leq s \leq u \leq t \\ s,u,t \in \mathbb{T}, s < t}} \mathbb{1}_{\{0 < t-s \leq \tau\}} e^{-\frac{t}{\tau}} \frac{|B_{sut}|}{|t-s|^\eta} \quad (1.45)$$

for $A: \{(s,t) \in \mathbb{T}^2: 0 \leq s < t\} \rightarrow \mathbb{R}$ and $B: \{(s,u,t) \in \mathbb{T}^3: 0 \leq s \leq u \leq t, s < t\} \rightarrow \mathbb{R}$.

THEOREM 1.18. (DISCRETE SEWING BOUND) *If a function $R = (R_{st})_{s < t \in \mathbb{T}}$ vanishes on consecutive points of $\mathbb{T}$ (i.e. $R_{t_i t_{i+1}} = 0$), then for any $\eta > 1$ and $\tau > 0$ we have*

$$\|R\|_{\eta,\tau}^\mathbb{T} \leq C_\eta \|\delta R\|_{\eta,\tau}^\mathbb{T} \quad \text{with} \quad C_\eta := 2^\eta \sum_{n \geq 1} \frac{1}{n^\eta} = 2^\eta \zeta(\eta) < \infty. \quad (1.46)$$

Proof. We fix $s, t \in \mathbb{T}$ with $s < t$ and we start by proving that

$$|R_{st}| \leq C_\eta \|\delta R\|_{\eta,\tau}^\mathbb{T} (t-s)^\eta. \quad (1.47)$$

We have $s = t_k$ and $t = t_{k+m}$ and we may assume that $m \geq 2$ (otherwise there is nothing to prove, since for $m = 1$ we have $R_{t_i t_{i+1}} = 0$).

Consider the partition $\mathcal{P} = \{s = t_k < t_{k+1} < \dots < t_{k+m} = t\}$ with m intervals. Note that for some index $i \in \{k+1, \dots, k+m-1\}$ we must have $t_{i+1} - t_{i-1} \leq \frac{2(t-s)}{m-1}$, otherwise we would get the contradiction

$$2(t-s) \geq \sum_{i=k+1}^{k+m-1} (t_{i+1} - t_{i-1}) > \sum_{i=k+1}^{k+m-1} \frac{2(t-s)}{m-1} = 2(t-s).$$

Removing the point t_i from $\mathcal{P}$ we obtain a partition $\mathcal{P}'$ with $m-1$ intervals.

Let $A = (A_{st})_{s < t \in \mathbb{T}}$ be a generic real function and set $I_{\mathcal{P}}(A) := \sum_{i=k}^{k+m-1} A_{t_i t_{i+1}}$. As in (1.25) we have

$$|I_{\mathcal{P}}(A) - I_{\mathcal{P}'}(A)| = |\delta A_{t_{i-1} t_i t_{i+1}}| \leq \frac{2^\eta (t-s)^\eta}{(m-1)^\eta} \sup_{\substack{s \leq u < v < w \leq t \\ u, v, w \in \mathbb{T}}} \frac{|\delta A_{uvw}|}{|w-u|^\eta}.$$

Iterating this argument, until we arrive at the trivial partition $\{s, t\}$, we get

$$|I_{\mathcal{P}}(A) - A_{st}| \leq C_\eta (t-s)^\eta \sup_{\substack{s \leq u < v < w \leq t \\ u, v, w \in \mathbb{T}}} \frac{|\delta A_{uvw}|}{|w-u|^\eta}, \quad (1.48)$$

with $C_\eta := \sum_{n \geq 1} \frac{2^\eta}{n^\eta} < \infty$ because $\eta > 1$.

We apply now (1.48) to $A = R$; since $R_{t_i t_{i+1}} = 0$ we obtain $I_{\mathcal{P}}(R) = 0$ and therefore (1.47) follows. Finally if $t-s \leq \tau$ then $w-u \leq \tau$ in the supremum in (1.48) and since $e^{-\frac{t}{\tau}} \leq e^{-\frac{w}{\tau}}$ we obtain

$$e^{-\frac{t}{\tau}} |R_{st}| \leq C_\eta (t-s)^\eta \|\delta R\|_{\eta, \tau}^{\mathbb{T}},$$

and the proof is complete. $\square$

We also have an analog of Lemma 1.16. We set for $f: \mathbb{T} \rightarrow \mathbb{R}$ and $\tau > 0$

$$\|f\|_{\infty, \tau}^{\mathbb{T}} := \sup_{t \in \mathbb{T}} e^{-\frac{t}{\tau}} |f_t|.$$

LEMMA 1.19. (DISCRETE SUPREMUM-HÖLDER BOUND) *For $\mathbb{T} := \{0 = t_1 < \dots < t_{\#\mathbb{T}}\} \subseteq \mathbb{R}_+$ set*

$$M := \max_{i=2, \dots, \#\mathbb{T}} |t_i - t_{i-1}|.$$

Then for all $f: \mathbb{T} \rightarrow \mathbb{R}$, $\tau \geq 2M$ and $\eta > 0$

$$\|f\|_{\infty, \tau}^{\mathbb{T}} \leq |f_0| + 5\tau^\eta \|\delta f\|_{\eta, \tau}^{\mathbb{T}}. \quad (1.49)$$

Proof. We define $T_0 := 0$ and for $i \geq 1$, as long as $\mathbb{T} \cap (T_{i-1}, T_{i-1} + \tau]$ is not empty, we set

$$T_i := \max \mathbb{T} \cap (T_{i-1}, T_{i-1} + \tau], \quad i = 1, \dots, N,$$

so that $T_N = \max \mathbb{T}$. We have by construction $T_i + M > T_{i-1} + \tau$ for all $i = 1, \dots, N-1$, and since $M \leq \frac{\tau}{2}$

$$T_i - T_{i-1} \geq \tau - M \geq \frac{\tau}{2}.$$

For $i = N$ we have only $T_N > T_{N-1}$. Therefore for $i = 1, \dots, N$

$$\begin{aligned} e^{-\frac{T_i}{\tau}} |f_{T_i}| &\leq |f_0| + \sum_{k=1}^i (T_k - T_{k-1})^\eta e^{-\frac{T_i - T_k}{\tau}} \left[e^{-\frac{T_k}{\tau}} \frac{|f_{T_k} - f_{T_{k-1}}|}{(T_k - T_{k-1})^\eta} \right] \\ &\leq |f_0| + \tau^\eta \|\delta f\|_{\eta, \tau}^\mathbb{T} \sum_{k=1}^i e^{-\frac{T_i - T_k}{\tau}} \\ &\leq |f_0| + \tau^\eta \|\delta f\|_{\eta, \tau}^\mathbb{T} \left(1 + \sum_{k=0}^{\infty} e^{-\frac{k}{2}} \right) \\ &\leq |f_{t_0}| + 4\tau^\eta \|\delta f\|_{\eta, \tau}^\mathbb{T}. \end{aligned}$$

Now for $t \in \mathbb{T} \setminus \{T_i\}_i$ we have $T_i < t < T_{i+1}$ for some i and then

$$\begin{aligned} e^{-\frac{t}{\tau}} |f_t| &\leq e^{-\frac{t}{\tau}} |f_{T_i}| + (t - T_i)^\eta e^{-\frac{t}{\tau}} \frac{|f_t - f_{T_i}|}{(t - T_i)^\eta} \leq e^{-\frac{T_i}{\tau}} |f_{T_i}| + \tau^\eta \|\delta f\|_{\eta, \tau}^\mathbb{T} \\ &\leq |f_0| + 5\tau^\eta \|\delta f\|_{\eta, \tau}^\mathbb{T}. \end{aligned}$$

The proof is complete. $\square$

1.9. EXTRA (TO BE COMPLETED)

We also introduce the usual supremum norm, for $F \in C_2$ and $G \in C_3$:

$$\|F\|_\infty := \sup_{0 \leq s \leq t \leq T} |F_{st}|, \quad \|G\|_\infty := \sup_{0 \leq s \leq u \leq t \leq T} |G_{sut}|,$$

and a corresponding weighted version, for $\tau \in (0, \infty)$:

$$\|F\|_{\infty, \tau} := \sup_{0 \leq s \leq t \leq T} e^{-\frac{t}{\tau}} |F_{st}|, \quad \|G\|_{\infty, \tau} := \sup_{0 \leq s \leq u \leq t \leq T} e^{-\frac{t}{\tau}} |G_{sut}|. \quad (1.50)$$

Note that

$$\lim_{\tau \rightarrow +\infty} \|F\|_{\infty, \tau} = \|F\|_\infty, \quad \lim_{\tau \rightarrow +\infty} \|G\|_{\eta, \tau} = \|G\|_\eta, \quad \lim_{\tau \rightarrow +\infty} \|H\|_{\eta, \tau} = \|H\|_\eta.$$

We have

$$\|F\|_{\eta, \tau} \leq \|G\|_{\infty, \tau} \|H\|_\eta, \quad (F_{sut} = G_{su} H_{ut}), \quad (1.51)$$

Note that $\|\cdot\|_{\eta, \tau}$ is only a semi-norm on C_n^η if $\tau < T$; we have at least

$$\|\cdot\|_{\eta, \tau} \leq \|\cdot\|_\eta \leq e^{\frac{T}{\tau}} \left(\|\cdot\|_{\eta, \tau} + \frac{1}{\tau^\eta} \|\cdot\|_{\infty, \tau} \right). \quad (1.52)$$

However, if $\tau \geq T$ we have again equivalence of norms

$$\|\cdot\|_{\eta, \tau} \leq \|\cdot\|_\eta \leq e^{\frac{T}{\tau}} \|\cdot\|_{\eta, \tau}, \quad \tau \geq T. \quad (1.53)$$

CHAPTER 2

DIFFERENCE EQUATIONS: THE YOUNG CASE

Fix a time horizon $T > 0$ and two dimensions $k, d \in \mathbb{N}$. We study the following *controlled difference equation* for an unknown path $Z: [0, T] \rightarrow \mathbb{R}^k$:

$$Z_t - Z_s = \sigma(Z_s)(X_t - X_s) + o(t - s), \quad 0 \leq s \leq t \leq T, \quad (2.1)$$

where the “driving path” $X: [0, T] \rightarrow \mathbb{R}^d$ and the function $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ are given, and $o(t - s)$ is *uniform* for $0 \leq s \leq t \leq T$ (see Remark 1.1).

The difference equation (2.1) is a natural generalized formulation of the *controlled differential equation*

$$\dot{Z}_t = \sigma(Z_t) \dot{X}_t, \quad 0 \leq t \leq T. \quad (2.2)$$

Indeed, as we showed in Chapter 1 (see Section 1.2), equations (2.1) and (2.2) are *equivalent* when X is continuously differentiable and σ is continuous, but (2.1) is meaningful also when X is non differentiable.

In this chapter we prove *well-posedness for the difference equation (2.1)* when the driving path $X \in \mathcal{C}^\alpha$ is Hölder continuous in the regime $\alpha \in]\frac{1}{2}, 1]$, called the *Young case*. The more challenging regime $\alpha \leq \frac{1}{2}$, called the *rough case*, is the object of the next Chapter 3, where new ideas will be introduced.

2.1. SUMMARY

Using the increment notation $\delta f_{st} := f_t - f_s$ from (1.11), we rewrite (2.1) as

$$\delta Z_{st} = \sigma(Z_s) \delta X_{st} + o(t - s), \quad 0 \leq s \leq t \leq T, \quad (2.3)$$

so that a solution of (2.3) is any path $Z: [0, T] \rightarrow \mathbb{R}^k$ such that the “remainder”

$$Z_{st}^{[2]} := \delta Z_{st} - \sigma(Z_s) \delta X_{st} \quad \text{satisfies} \quad Z_{st}^{[2]} = o(t - s). \quad (2.4)$$

We summarize the main results of this chapter stating *local and global existence, uniqueness of solutions and continuity of the solution map* for the difference equation (2.3) under natural assumptions on σ . We will actually prove more precise results, which yield quantitative estimates.

THEOREM 2.1. (WELL-POSEDNESS) *Let $X: [0, T] \rightarrow \mathbb{R}^d$ be of class $\mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$ and let $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$. Then we have:*

- **local existence:** *if σ is locally γ -Hölder with $\gamma \in (\frac{1}{\alpha} - 1, 1]$ (e.g. of class C^1), then for every $z_0 \in \mathbb{R}^k$ there is a possibly shorter time horizon $T' = T'_{\alpha, X, \sigma}(z_0) \in]0, T]$ and a path $Z: [0, T'] \rightarrow \mathbb{R}^k$ starting from $Z_0 = z_0$ which solves (2.3) for $0 \leq s \leq t \leq T'$;*

- **global existence:** if σ is globally γ -Hölder with $\gamma \in (\frac{1}{\alpha} - 1, 1]$ (e.g. of class C^1 with $\|\nabla\sigma\|_\infty < \infty$), then we can take $T'_{\alpha,X,\sigma}(z_0) = T$ for any $z_0 \in \mathbb{R}^d$;
- **uniqueness:** if σ is of class C^γ with $\gamma \in (\frac{1}{\alpha}, 2]$ (e.g. if σ is of class C^2), then there is exactly one solution Z of (2.3) with $Z_0 = z_0$;
- **continuity of the solution map:** if σ is differentiable with bounded and globally $(\gamma - 1)$ -Hölder gradient with $\gamma \in (\frac{1}{\alpha}, 2]$ (i.e. $\|\nabla\sigma\|_\infty < \infty$, $[\nabla\sigma]_{C^{\gamma-1}} < \infty$), then the solution Z of (2.3) is a continuous function of the starting point z_0 and driving path X : the map $(z_0, X) \mapsto Z$ is continuous from $\mathbb{R}^k \times \mathcal{C}^\alpha \rightarrow \mathcal{C}^\alpha$.

In the first part of this chapter, we give for granted the existence of solutions and we focus on their properties: we prove *a priori estimates* in Section 2.3, *uniqueness of solutions* in Section 2.4 and *continuity of the solution map* in Section 2.5. A key role is played by the Sewing Bound from Chapter 1, see Theorems 1.9 and 1.17, and its discrete version, see Theorem 1.18.

The proof of local and global *existence of solutions* of (2.3) is given at the end of this chapter, see Section 2.6, exploiting a suitable Euler scheme.

2.2. SET-UP

We collect here some notions and tools that will be used extensively.

We recall that C_1 denotes the space of continuous functions $f: [0, T] \rightarrow \mathbb{R}^k$. Similarly, C_2 and C_3 are the spaces of continuous functions of two and three ordered variables, i.e. defined on $[0, T]_{\leq}^2$ and $[0, T]_{\leq}^3$, see (1.7)-(1.8).

We are going to exploit the *weighted semi-norms* $\|\cdot\|_{\eta,\tau}$, see (1.34)-(1.35) (see also (1.9) for the original norm $\|\cdot\|_\eta$). These are useful to bound the *weighted supremum norm* $\|f\|_{\infty,\tau}$ of a function $f \in C_1$, see (1.38) and (1.41):

$$\|f\|_{\infty,\tau} \leq |f_0| + 3(\tau \wedge T)^\eta \|\delta f\|_{\eta,\tau}, \quad \forall \eta, \tau > 0. \quad (2.5)$$

It follows directly from the definitions (1.34)-(1.35) that

$$\|\cdot\|_{\eta,\tau} \leq (\tau \wedge T)^{\eta'} \|\cdot\|_{\eta+\eta',\tau}, \quad \forall \eta, \eta' > 0, \quad (2.6)$$

because $(t-s)^\eta \geq (t-s)^{\eta+\eta'} (\tau \wedge T)^{-\eta'}$ for $0 \leq s \leq t \leq T$ with $t-s \leq \tau$.

Remark 2.2. The factor $(\tau \wedge T)^{\eta'}$ in the RHS of (2.6) can be made small by choosing τ small while keeping T fixed. This is why we included the indicator function $\mathbb{1}_{\{0 < t-s \leq \tau\}}$ in the definition (1.34)-(1.35) of the norms $\|\cdot\|_{\eta,\tau}$: without this indicator function, instead of $(\tau \wedge T)^{\eta'}$ we would have $T^{\eta'}$, which is small only when T is small.

We will often work with functions $F \in C_2$ or $F \in C_3$ that are *product of two factors*, like $F_{st} = g_s H_{st}$ or $F_{sut} = G_{su} H_{ut}$. We show in the next result that the semi-norm $\|F\|_{\eta,\tau}$ can be controlled by a product of suitable norms for each factor.

LEMMA 2.3. (WEIGHTED BOUNDS) *For any $\eta, \eta' \in (0, \infty)$ and $\tau > 0$, we have*

$$\text{if } F_{st} = g_s H_{st} \text{ or } F_{st} = g_t H_{st} \quad \text{then} \quad \|F\|_{\eta,\tau} \leq \|g\|_{\infty,\tau} \|H\|_{\eta}, \quad (2.7)$$

$$\text{if } F_{sut} = G_{su} H_{ut} \quad \text{then} \quad \|F\|_{\eta+\eta',\tau} \leq \|G\|_{\eta,\tau} \|H\|_{\eta'}. \quad (2.8)$$

Proof. If $F_{st} = g_t H_{st}$, by (1.38) we can estimate $e^{-t/\tau} |g_t| \leq \|g\|_{\infty, \tau}$ to get (2.7). If $F_{st} = g_s H_{st}$, for $s \leq t$ we can bound $e^{-t/\tau} \leq e^{-s/\tau}$ in the definition (1.34)-(1.35) of $\|\cdot\|_{\eta, \tau}$, hence again by (1.38) we can estimate $e^{-s/\tau} |g_s| \leq \|g\|_{\infty, \tau}$ to get (2.7).

If $F_{sut} = G_{su} H_{ut}$, we can further bound $(t-s)^{\eta+\eta'} \geq (t-u)^\eta (u-s)^{\eta'}$ in (1.35) and then estimate $e^{-s/\tau} G_{su} / (u-s)^\eta \leq \|G\|_{\eta, \tau}$, which yields (2.8). $\square$

We stress that in the RHS of (2.7) and (2.8) *only one factor gets the weighted norm or semi-norm*, while the other factor gets the non-weighted norm $\|\cdot\|_\eta$. We will sometimes need an extra weight, which can be introduced as follows.

LEMMA 2.4. (EXTRA WEIGHT) *For any $\eta, \bar{\tau} \in (0, \infty)$ and $0 < \tau \leq \bar{\tau}$, we have*

$$\text{if } F_{st} = g_s H_{st} \quad \text{or} \quad F_{st} = g_t H_{st} \quad \text{then} \quad \|F\|_{\eta, \tau} \leq \|g\|_{\infty, \tau} e^{\frac{\tau}{\bar{\tau}}} \|H\|_{\eta, \bar{\tau}}. \quad (2.9)$$

Proof. Recall the definition (1.34)-(1.35) of $\|\cdot\|_{\eta, \tau}$ and note that for $0 \leq s \leq t \leq T$ we have $e^{-t/\tau} |g_t| \leq \|g\|_{\infty, \tau}$ and $e^{-s/\tau} |g_s| \leq \|g\|_{\infty, \tau}$ (see the proof of Lemma 2.3). Finally, for $t-s \leq \tau \leq \bar{\tau}$ we can estimate $|H_{st}| \leq e^{T/\bar{\tau}} e^{-t/\bar{\tau}} |H_{st}| \leq e^{T/\bar{\tau}} \|H\|_{\eta, \bar{\tau}} (t-s)^\eta$. $\square$

We recall that $\mathbb{R}^k \otimes (\mathbb{R}^d)^* \simeq \mathbb{R}^{k \times d}$ is the space of linear applications from $\mathbb{R}^d$ to $\mathbb{R}^k$ equipped with the Hilbert-Schmidt (Euclidean) norm $|\cdot|$. We say that a function is of class C^m if it is continuously differentiable m times. Given $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ of class C^2 , that we represent by $\sigma_j^i(z)$ with $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, d\}$, we denote by $\nabla \sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^k)^*$ its gradient and by $\nabla^2 \sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^k)^* \otimes (\mathbb{R}^k)^*$ its Hessian, represented for $i, a, b \in \{1, \dots, k\}$ and $j \in \{1, \dots, d\}$ by

$$(\nabla \sigma(z))_{ja}^i = \frac{\partial \sigma_j^i}{\partial z_a}(z), \quad (\nabla^2 \sigma(z))_{jab}^i = \frac{\partial^2 \sigma_j^i}{\partial z_a \partial z_b}(z).$$

Remark 2.5. (NORM OF THE GRADIENT OF LIPSCHITZ FUNCTIONS) For a *locally Lipschitz function* $\psi: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$ we can define the “norm of the gradient” at any point (even where ψ may not be differentiable):

$$|\nabla \psi(z)| := \limsup_{y \rightarrow z} \frac{|\psi(y) - \psi(z)|}{|y - z|} \in [0, \infty).$$

Similarly, $|\nabla^2 \psi(z)|$ is well defined as soon as ψ is *differentiable with locally Lipschitz gradient* $\nabla \psi$ (which is slightly less than requiring $\psi \in C^2$).

2.3. A PRIORI ESTIMATES

In this section we prove *a priori estimates* for solutions of (2.3) assuming that σ is *globally Hölder*, that is $[\sigma]_{C^\gamma} < \infty$ with $\gamma > \frac{1}{\alpha}$, recall the definition (1.16) of the Hölder seminorm $[\sigma]_{C^\gamma}$. Note that σ is *not* assumed to be bounded.

We first observe that if the driving path X is of class $\mathcal{C}^\alpha$, then any solution Z of (2.3) is also of class $\mathcal{C}^\alpha$, as soon as σ is continuous.

LEMMA 2.6. (HÖLDER REGULARITY) *Let X be of class $\mathcal{C}^\alpha$ with $\alpha \in]0, 1]$ and let σ be continuous. Then any solution Z of (2.3) is of class $\mathcal{C}^\alpha$.*

Proof. We know by Lemma 1.2 that Z is continuous, more precisely by (1.6) we have $|\delta Z_{st}| \leq C |\delta X_{st}| + o(t-s)$ with $C < \infty$. Since $|\delta X_{st}| \leq \|\delta X\|_\alpha (t-s)^\alpha$ and $o(t-s) = o((t-s)^\alpha)$ for any $\alpha \leq 1$, it follows that $Z \in \mathcal{C}^\alpha$. $\square$

We next formulate the announced a priori estimates. It is convenient to use the weighted semi-norms $\|\cdot\|_{\eta, \tau}$ in (1.34)-(1.35) (note that the usual norms $\|\cdot\|_\eta$ in (1.9) can be recovered by letting $\tau \rightarrow \infty$).

THEOREM 2.7. (A PRIORI ESTIMATES) *Let X be of class $\mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$ and let σ be globally γ -Hölder with $\gamma \in (\frac{1}{\alpha} - 1, 1]$. Then, for any solution $Z: [0, T] \rightarrow \mathbb{R}^k$ of (2.3), the remainder $Z_{st}^{[2]} := \delta Z_{st} - \sigma(Z_s) \delta X_{st}$ satisfies $Z^{[2]} \in C_2^{(\gamma+1)\alpha}$, more precisely for any $\tau > 0$*

$$\|Z^{[2]}\|_{(\gamma+1)\alpha, \tau} \leq C_{\alpha, \gamma, X, \sigma} \|\delta Z\|_{\alpha, \tau}^\gamma \quad \text{with } C_{\alpha, \gamma, X, \sigma} := K_{(\gamma+1)\alpha} \|\delta X\|_\alpha [\sigma]_{\mathcal{C}^\gamma}, \quad (2.10)$$

where $K_\eta = (1 - 2^{1-\eta})^{-1}$. Moreover, if either T or τ is small enough, we have

$$\|\delta Z\|_{\alpha, \tau} \leq 1 \vee (2 \|\delta X\|_\alpha |\sigma(Z_0)|) \quad \text{for } (\tau \wedge T)^\alpha \leq \varepsilon_{\alpha, \gamma, X, \sigma}, \quad (2.11)$$

where we define

$$\varepsilon_{\alpha, \gamma, X, \sigma} := \frac{1}{2(K_{(\gamma+1)\alpha} + 3) \|\delta X\|_\alpha [\sigma]_{\mathcal{C}^\gamma}}. \quad (2.12)$$

If σ is globally Lipschitz, namely if we can take $\gamma = 1$, we can improve (2.11) to

$$\|\delta Z\|_{\alpha, \tau} \leq 2 \|\delta X\|_\alpha |\sigma(Z_0)| \quad \text{for } (\tau \wedge T)^\alpha \leq \varepsilon_{\alpha, 1, X, \sigma}. \quad (2.13)$$

Proof. We first prove (2.10). Since $Z_{st}^{[2]} = o(t-s)$ by definition of solution, see (2.4), we can estimate $Z^{[2]}$ in terms of $\delta Z^{[2]}$, by the weighted Sewing Bound (1.42). Let us compute $\delta Z_{sut}^{[2]} = Z_{st}^{[2]} - Z_{su}^{[2]} - Z_{ut}^{[2]}$: recalling (2.4) and (1.33), since $\delta \circ \delta = 0$, we have

$$\delta Z_{sut}^{[2]} = \delta \sigma(Z)_{su} \delta X_{ut} = (\sigma(Z_u) - \sigma(Z_s)) (X_t - X_u). \quad (2.14)$$

Since $|\sigma(z) - \sigma(\bar{z})| \leq [\sigma]_{\mathcal{C}^\gamma} |z - \bar{z}|^\gamma$ for all $z, \bar{z} \in \mathbb{R}^d$, we can bound

$$\|\delta \sigma(Z)\|_{\gamma\alpha, \tau} \leq [\sigma]_{\mathcal{C}^\gamma} \|\delta Z\|_{\alpha, \tau}^\gamma, \quad (2.15)$$

hence by (2.8) we obtain

$$\|\delta Z^{[2]}\|_{(\gamma+1)\alpha, \tau} \leq \|\delta X\|_\alpha [\sigma]_{\mathcal{C}^\gamma} \|\delta Z\|_{\alpha, \tau}^\gamma.$$

Applying the weighted Sewing Bound (1.42), for $(\gamma+1)\alpha > 1$ we then obtain

$$\|Z^{[2]}\|_{(\gamma+1)\alpha, \tau} \leq K_{(\gamma+1)\alpha} \|\delta X\|_\alpha [\sigma]_{\mathcal{C}^\gamma} \|\delta Z\|_{\alpha, \tau}^\gamma, \quad (2.16)$$

which proves (2.10).

We next prove (2.11). To simplify notation, let us set $\varepsilon := (\tau \wedge T)^\alpha$. Recalling (2.7) and (2.6), we obtain by (2.4)

$$\begin{aligned} \|\delta Z\|_{\alpha,\tau} &\leq \|\sigma(Z) \delta X\|_{\alpha,\tau} + \|Z^{[2]}\|_{\alpha,\tau} \\ &\leq \|\sigma(Z)\|_{\infty,\tau} \|\delta X\|_\alpha + \varepsilon^\gamma \|Z^{[2]}\|_{(\gamma+1)\alpha,\tau}. \end{aligned} \quad (2.17)$$

We can estimate $\|\sigma(Z)\|_{\infty,\tau}$ by (2.5) and (2.15):

$$\|\sigma(Z)\|_{\infty,\tau} \leq |\sigma(Z_0)| + 3\varepsilon^\gamma [\sigma]_{C^\gamma} \|\delta Z\|_{\alpha,\tau}^\gamma.$$

Plugging this and (2.16) into (2.17), we get

$$\begin{aligned} \|\delta Z\|_{\alpha,\tau} &\leq (|\sigma(Z_0)| + 3\varepsilon^\gamma [\sigma]_{C^\gamma} \|\delta Z\|_{\alpha,\tau}^\gamma) \|\delta X\|_\alpha + \\ &\quad + \varepsilon^\gamma K_{(\gamma+1)\alpha} \|\delta X\|_\alpha [\sigma]_{C^\gamma} \|\delta Z\|_{\alpha,\tau}^\gamma \\ &= \|\delta X\|_\alpha |\sigma(Z_0)| + \frac{1}{2} \frac{\varepsilon^\gamma}{\varepsilon_{\alpha,\gamma,X,\sigma}} \|\delta Z\|_{\alpha,\tau}^\gamma, \end{aligned}$$

where $\varepsilon_{\alpha,\gamma,X,\sigma}$ is defined in (2.12). For $\varepsilon^\gamma \leq \varepsilon_{\alpha,\gamma,X,\sigma}$ the last term is bounded by $\frac{1}{2} \|\delta Z\|_{\alpha,\tau}^\gamma$ which is finite by Lemma 2.6. If $\|\delta Z\|_{\alpha,\tau} \leq 1$ then (2.11) holds trivially; if not, $\frac{1}{2} \|\delta Z\|_{\alpha,\tau}^\gamma \leq \frac{1}{2} \|\delta Z\|_{\alpha,\tau}$. Bringing this term in the LHS we obtain (2.11).

To prove (2.13), we argue as for (2.11) and since $\gamma = 1$ we obtain

$$\|\delta Z\|_{\alpha,\tau} \leq \|\delta X\|_\alpha |\sigma(Z_0)| + \frac{1}{2} \frac{\varepsilon}{\varepsilon_{\alpha,1,X,\sigma}} \|\delta Z\|_{\alpha,\tau}.$$

For $\varepsilon \leq \varepsilon_{\alpha,1,X,\sigma}$ the last term is bounded by $\frac{1}{2} \|\delta Z\|_{\alpha,\tau}$ which is finite by Lemma 2.6. Bringing this term in the LHS we obtain (2.13), and this completes the proof. $\square$

2.4. UNIQUENESS

In this section we prove uniqueness of solutions to (2.3) assuming that σ is of class C^1 with locally Hölder gradient (without boundedness assumption on σ). This improves on Theorem 1.7, both because we allow for non-linear σ and because we do not require that the time horizon $T > 0$ is small.

We first need an elementary but fundamental estimate on the difference of increments of a function. Given $\Psi: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$, we use the notation

$$C_{\Psi,R} := \sup \{ |\Psi(x)| : x \in \mathbb{R}^k, |x| \leq R \}. \quad (2.18)$$

LEMMA 2.8. (DIFFERENCE OF INCREMENTS) *Let $\psi: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$ be of class $\mathcal{C}_{\text{loc}}^{1+\rho}$ for some $0 < \rho \leq 1$ (i.e. ψ is differentiable with $\nabla \psi$ of class $\mathcal{C}_{\text{loc}}^\rho$). Then for any $R > 0$ and for all $x, \bar{x}, y, \bar{y} \in \mathbb{R}^k$ with $\max \{|x|, |y|, |\bar{x}|, |\bar{y}|\} \leq R$ we can estimate*

$$\begin{aligned} |[\psi(x) - \psi(y)] - [\psi(\bar{x}) - \psi(\bar{y})]| \\ \leq C'_R |(x - y) - (\bar{x} - \bar{y})| + C''_R \{|x - y|^\rho + |\bar{x} - \bar{y}|^\rho\} |y - \bar{y}|, \end{aligned} \quad (2.19)$$

where $C'_R := \sup \{ |\nabla \psi(x)| : |x| \leq R \}$ and $C''_R := \sup \left\{ \frac{|\nabla \psi(x) - \nabla \psi(y)|}{|x - y|^\rho} : |x|, |y| \leq R \right\}.$

Proof. For $z, w \in \mathbb{R}^k$ we can write

$$\psi(z) - \psi(w) = \hat{\psi}(z, w) (z - w),$$

where $\hat{\psi}(z, w) := \int_0^1 \nabla \psi(u z + (1 - u) w) du \in \mathbb{R}^\ell \otimes (\mathbb{R}^k)^*$, therefore

$$\begin{aligned} [\psi(x) - \psi(y)] - [\psi(\bar{x}) - \psi(\bar{y})] &= [\psi(x) - \psi(\bar{x})] - [\psi(y) - \psi(\bar{y})] \\ &= \hat{\psi}(x, \bar{x}) (x - \bar{x}) - \hat{\psi}(y, \bar{y}) (y - \bar{y}) \\ &= \hat{\psi}(x, \bar{x}) [(x - \bar{x}) - (y - \bar{y})] \\ &\quad + [\hat{\psi}(x, \bar{x}) - \hat{\psi}(y, \bar{y})] (y - \bar{y}). \end{aligned}$$

By definition of C'_R and C''_R we have $|\hat{\psi}(x, \bar{x})| \leq C'_R$ and

$$\begin{aligned} |\hat{\psi}(x, \bar{x}) - \hat{\psi}(y, \bar{y})| &\leq |\hat{\psi}(x, \bar{x}) - \hat{\psi}(y, \bar{x})| + |\hat{\psi}(y, \bar{x}) - \hat{\psi}(y, \bar{y})| \\ &\leq C''_R \{|x - y|^\rho + |\bar{x} - \bar{y}|^\rho\}, \end{aligned}$$

hence (2.19) follows. $\square$

We are now ready to state and prove the announced uniqueness result.

THEOREM 2.9. (UNIQUENESS) *Let X be of class $\mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$ and let σ be of class $\mathcal{C}^\gamma$ for some $\gamma > \frac{1}{\alpha}$ (for instance, we can take $\sigma \in \mathcal{C}^2$). Then for every $z_0 \in \mathbb{R}^k$ there exists at most one solution Z to (2.3) with $Z_0 = z_0$.*

Proof. Let Z and $\bar{Z}$ be two solutions of (2.3), i.e. they satisfy (2.4), and set

$$Y := Z - \bar{Z}.$$

We want to show that, for $\tau > 0$ small enough, we have

$$\|Y\|_{\infty, \tau} \leq 2 \|Y_0\|,$$

where the weighted norm $\|\cdot\|_{\infty, \tau}$ was defined in (1.38). In particular, if we assume that $Z_0 = \bar{Z}_0$, we obtain $\|Y\|_{\infty, \tau} = 0$ and hence $Z = \bar{Z}$.

We know by (2.5) that for any $\tau > 0$

$$\|Y\|_{\infty, \tau} \leq \|Y_0\| + 3\tau^\alpha \|\delta Y\|_{\alpha, \tau}, \quad (2.20)$$

where we recall that the weighted semi-norm $\|\cdot\|_{\alpha, \tau}$ was defined in (1.34). We now define $Y^{[2]}$ as the difference between the remainders $Z^{[2]}$ and $\bar{Z}^{[2]}$ of the solutions Z and $\bar{Z}$ as defined in (2.4), that is

$$Y_{st}^{[2]} := Z_{st}^{[2]} - \bar{Z}_{st}^{[2]} = \delta Y_{st} - (\sigma(Z_s) - \sigma(\bar{Z}_s)) \delta X_{st}. \quad (2.21)$$

(We are slightly abusing notation, since $Y^{[2]}$ is not the remainder of Y when σ is not linear.) By assumption $\sigma \in \mathcal{C}^\gamma$ for some $\gamma > \frac{1}{\alpha}$: renaming γ as $\gamma \wedge 2$, we may assume that $\gamma \in]\frac{1}{\alpha}, 2]$. We are going to prove the following inequalities: for any $\tau > 0$

$$\|\delta Y\|_{\alpha, \tau} \leq c_1 \|Y\|_{\infty, \tau} + \tau^{(\gamma-1)\alpha} \|Y^{[2]}\|_{\gamma\alpha, \tau}, \quad (2.22)$$

$$\|Y^{[2]}\|_{\gamma\alpha, \tau} \leq c_2 \|Y\|_{\infty, \tau} + c'_2 \tau^{(\gamma-1)\alpha} \|Y^{[2]}\|_{\gamma\alpha, \tau}, \quad (2.23)$$

for finite constants c_i, c'_i that may depend on $X, \sigma, Z, \bar{Z}$ but not on τ .

Let us complete the proof assuming (2.22) and (2.23). Note that $(\gamma - 1)\alpha > 0$ by assumption. If we fix $\tau > 0$ small, so that $c'_2 \tau^{(\gamma-1)\alpha} < \frac{1}{2}$, from (2.23) we get $\|Y^{[2]}\|_{\gamma\alpha, \tau} \leq 2c_2 \|Y\|_{\infty, \tau}$ which plugged into (2.22) yields $\|\delta Y\|_{\alpha, \tau} \leq 2c_1 \|Y\|_{\infty, \tau}$ for $\tau > 0$ small (it suffices that $2c_2 \tau^{(\gamma-1)\alpha} < c_1$). Finally, plugging this into (2.20) and possibly choosing $\tau > 0$ even smaller, we obtain our goal $\|Y\|_{\infty, \tau} \leq 2|Y_0|$ which completes the proof.

It remains to prove (2.22) and (2.23). Using the notation from Lemma 2.8 we set

$$\begin{aligned} C'_1 &:= \sup \{ |\nabla \sigma(x)| : |x| \leq \|Z\|_\infty \vee \|\bar{Z}\|_\infty \}, \\ C''_1 &:= \sup \left\{ \frac{|\nabla \sigma(x) - \nabla \sigma(y)|}{|x - y|^\rho} : |x|, |y| \leq \|Z\|_\infty \vee \|\bar{Z}\|_\infty \right\}. \end{aligned}$$

so that $|\sigma(Z_t) - \sigma(\bar{Z}_t)| \leq C'_1 |Z_t - \bar{Z}_t|$ and, therefore,

$$\|\sigma(Z) - \sigma(\bar{Z})\|_{\infty, \tau} \leq C'_1 \|Y\|_{\infty, \tau}. \quad (2.24)$$

We now exploit (2.21) to estimate $\|\delta Y\|_{\alpha, \tau}$: applying (2.7) we obtain

$$\begin{aligned} \|\delta Y\|_{\alpha, \tau} &\leq \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty, \tau} \|\delta X\|_\alpha + \|Y^{[2]}\|_{\alpha, \tau} \\ &\leq C'_1 \|Y\|_{\infty, \tau} \|\delta X\|_\alpha + \tau^{(\gamma-1)\alpha} \|Y^{[2]}\|_{\gamma\alpha, \tau}, \end{aligned} \quad (2.25)$$

where we note that $\|Y^{[2]}\|_{\alpha, \tau} \leq \tau^{(\gamma-1)\alpha} \|Y^{[2]}\|_{\gamma\alpha, \tau}$ by (2.6). We have shown that (2.22) holds with $c_1 = C'_1 \|\delta X\|_\alpha$.

We finally prove (2.23). Since $Y_{st}^{[2]} = o(t - s)$, see (2.21) and (2.4), we bound $Z^{[2]}$ by its increment $\delta Z^{[2]}$ through the weighted Sewing Bound (1.42):

$$\|Y^{[2]}\|_{\gamma\alpha, \tau} \leq K_{\gamma\alpha} \|\delta Y^{[2]}\|_{\gamma\alpha, \tau}, \quad (2.26)$$

hence we focus on $\|\delta Y^{[2]}\|_{\gamma\alpha, \tau}$. By (2.21) and (1.33), since $\delta \circ \delta = 0$, we have

$$\delta Y_{sut}^{[2]} = (\delta\sigma(Z)_{su} - \delta\sigma(\bar{Z})_{su}) \delta X_{ut}. \quad (2.27)$$

Applying the estimate (2.19) for $x = Z_u, y = Z_s, \bar{x} = \bar{Z}_u, \bar{y} = \bar{Z}_s$, we can write

$$\begin{aligned} |\delta\sigma(Z)_{su} - \delta\sigma(\bar{Z})_{su}| &\leq C'_1 |\delta Z_{su} - \delta \bar{Z}_{su}| + C''_1 \{ |\delta Z_{su}|^{\gamma-1} + |\delta \bar{Z}_{su}|^{\gamma-1} \} |Z_s - \bar{Z}_s| \\ &= C'_1 |\delta Y_{su}| + C''_1 \{ |\delta Z_{su}|^{\gamma-1} + |\delta \bar{Z}_{su}|^{\gamma-1} \} |Y_s|. \end{aligned} \quad (2.28)$$

hence by (2.7) we get

$$\begin{aligned} \|\delta\sigma(Z) - \delta\sigma(\bar{Z})\|_{(\gamma-1)\alpha, \tau} &\leq C'_1 \|\delta Y\|_{(\gamma-1)\alpha, \tau} + \\ &\quad + C''_1 \{ \|\delta Z\|_\alpha^{\gamma-1} + \|\delta \bar{Z}\|_\alpha^{\gamma-1} \} \|Y\|_{\infty, \tau}. \end{aligned} \quad (2.29)$$

If we take $\tau \leq 1$ we can bound $\|\delta Y\|_{(\gamma-1)\alpha, \tau} \leq \|\delta Y\|_{\alpha, \tau}$ by (2.6) (recall that we are assuming $\gamma \leq 2$). Then by (2.27) we obtain, recalling (2.8),

$$\|\delta Y^{[2]}\|_{\gamma\alpha, \tau} \leq \|\delta X\|_\alpha \|\delta\sigma(Z) - \delta\sigma(\bar{Z})\|_{(\gamma-1)\alpha, \tau} \leq \tilde{c}_1 (\|\delta Y\|_{\alpha, \tau} + \|Y\|_{\infty, \tau}),$$

for a suitable (explicit) constant $\tilde{c}_1 = \tilde{c}_1(\sigma, Z, \bar{Z}, X)$. Applying (2.22), we obtain

$$\|\delta Y^{[2]}\|_{\gamma\alpha, \tau} \leq (c_1 + 1) \tilde{c}_1 \|Y\|_{\infty, \tau} + \tilde{c}_1 \tau^{(\gamma-1)\alpha} \|Y^{[2]}\|_{\gamma\alpha, \tau},$$

which plugged into (2.26) shows that (2.23) holds. The proof is complete. $\square$

We conclude with an example of (2.19).

Example 2.10. If $\sigma: \mathbb{R} \rightarrow \mathbb{R}$ is $\sigma(x) = x^2$, then we have

$$\begin{aligned} & (\sigma(x) - \sigma(y)) - (\sigma(\bar{x}) - \sigma(\bar{y})) \\ &= (x^2 - y^2) - (\bar{x}^2 - \bar{y}^2) = (x^2 - \bar{x}^2) - (y^2 - \bar{y}^2) \\ &= (x - \bar{x})(x + \bar{x}) - (y - \bar{y})(y + \bar{y}) \\ &= [(x - \bar{x}) - (y - \bar{y})](y + \bar{y}) + (x - \bar{x})[(x + \bar{x}) - (y + \bar{y})] \\ &= [(x - \bar{x}) - (y - \bar{y})](y + \bar{y}) + (x - \bar{x})[(x - y) + (\bar{x} - \bar{y})], \end{aligned}$$

where in the second last equality we have summed and subtracted $(y - \bar{y})(x + \bar{x})$. If we use this formula for $x = Z_t$, $y = Z_s$ and $\bar{x} = \bar{Z}_t$, $\bar{y} = \bar{Z}_s$, then we obtain

$$\delta(Z^2 - \bar{Z}^2)_{st} = \delta(Z - \bar{Z})_{st}(Z_s + \bar{Z}_s) + (Z_t - \bar{Z}_t)[\delta Z_{st} + \delta \bar{Z}_{st}],$$

which is in the spirit of (2.19) with $\rho = 1$. It follows that

$$\|\delta(Z^2 - \bar{Z}^2)\|_\alpha \leq 2 \|\bar{Z}\|_\infty \|\delta(Z - \bar{Z})\|_\alpha + \|Z - \bar{Z}\|_\infty [\|\delta Z\|_\alpha + \|\delta \bar{Z}\|_\alpha],$$

which is the form that (2.29) takes in this particular case.

2.5. CONTINUITY OF THE SOLUTION MAP

In this section we assume that σ is *globally Lipschitz* and of class C^1 with a *globally γ -Hölder gradient*, i.e. $\|\nabla \sigma\|_\infty < \infty$ and $[\nabla \sigma]_{C^\gamma} < \infty$, with $\gamma > \frac{1}{\alpha}$ (recall the definition (1.16) of the Hölder seminorm $[\nabla \sigma]_{C^\gamma}$), while σ itself is not assumed to be bounded. Under these assumptions, we have *global existence and uniqueness* of solutions $Z: [0, T] \rightarrow \mathbb{R}^k$ to (2.3) for any time horizon $T > 0$, for any starting point $Z_0 \in \mathbb{R}^k$ and for any driving path X of class $\mathcal{C}^\alpha$ with $\frac{1}{2} < \alpha \leq 1$ (as we will prove in Section 2.6).

We can thus consider the *solution map*:

$$\begin{aligned} \Phi: \mathbb{R}^k \times \mathcal{C}^\alpha &\longrightarrow \mathcal{C}^\alpha \\ (Z_0, X) &\longmapsto Z := \begin{cases} \text{unique solution of (2.3) for } t \in [0, T] \\ \text{starting from } Z_0 \end{cases} \end{aligned} \quad (2.30)$$

We prove in this section that this map is *continuous*, in fact *locally Lipschitz*.

Remark 2.11. The continuity of the solution map is a highly non-trivial property. Indeed, when X is of class C^1 , note that Z solves the equation

$$Z_t = Z_0 + \int_0^t \sigma(Z_s) \dot{X}_s \, ds, \quad (2.31)$$

which is based on the derivative $\dot{X}$ of X . We instead consider driving paths $X \in \mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$ which are continuous but may be non-differentiable.

We shall see in the next chapters that the continuity of the solution map holds also in more complex situations such as $X \in \mathcal{C}^\alpha$ with $\alpha \leq \frac{1}{2}$, which cover the case when X is a Brownian motion and Z is the solution to a SDE.

Before stating the continuity of the solution map, we recall that the space $\mathcal{C}^\alpha$ is equipped with the norm $\|f\|_{\mathcal{C}^\alpha} := \|f\|_\infty + \|\delta f\|_\alpha$, see Remark 1.4, but *an equivalent norm is $\|f\|_{\infty, \tau} + \|\delta f\|_{\alpha, \tau}$ for any choice of the weight $\tau > 0$* , see Remark 1.15.

THEOREM 2.12. (CONTINUITY OF THE SOLUTION MAP) *Let σ be globally Lipschitz with a globally $(\gamma - 1)$ -Hölder gradient: $\|\nabla \sigma\|_\infty < \infty$ and $[\nabla \sigma]_{\mathcal{C}^{\gamma-1}} < \infty$, with $\gamma \in (\frac{1}{\alpha}, 2]$. Then, for any $T > 0$ and $\alpha \in]\frac{1}{2}, 1]$, the solution map $(Z_0, X) \mapsto Z$ in (2.30) is locally Lipschitz.*

More explicitly, given $M_0, M, D < \infty$, if we assume that

$$\max \{ \|\nabla \sigma\|_\infty, [\nabla \sigma]_{\mathcal{C}^{\gamma-1}} \} \leq D,$$

and we consider starting points $Z_0, \bar{Z}_0 \in \mathbb{R}^d$ and driving paths $X, \bar{X} \in \mathcal{C}^\alpha$ with

$$\max \{ |\sigma(Z_0)|, |\sigma(\bar{Z}_0)| \} \leq M_0, \quad \max \{ \|\delta X\|_\alpha, \|\delta \bar{X}\|_\alpha \} \leq M, \quad (2.32)$$

then the corresponding solutions $Z = (Z_s)_{s \in [0, T]}$, $\bar{Z} = (\bar{Z}_s)_{s \in [0, T]}$ of (2.3) satisfy

$$\|Z - \bar{Z}\|_{\infty, \tau} + \|\delta Z - \delta \bar{Z}\|_{\alpha, \tau} \leq \mathfrak{C}_M |Z_0 - \bar{Z}_0| + 6 M_0 \|\delta X - \delta \bar{X}\|_\alpha, \quad (2.33)$$

provided $0 < \tau \wedge T \leq \hat{\tau}$ for a suitable $\hat{\tau} = \hat{\tau}_{\alpha, \gamma, T, D, M_0, M} > 0$, where we set

$$\mathfrak{C}_M := 2 (\|\nabla \sigma\|_\infty M + 1) \leq 2 (D M + 1).$$

Proof. Let us define the constant

$$\mathfrak{c}_M := \|\nabla \sigma\|_\infty M \leq D M. \quad (2.34)$$

We fix two solutions Z and $\bar{Z}$ of (2.3) with respective driving paths X and $\bar{X}$. If we define $Y := Z - \bar{Z}$, we can rewrite our goal (2.33) as

$$\|Y\|_{\infty, \tau} + \|\delta Y\|_{\alpha, \tau} \leq 6 M_0 \|\delta X - \delta \bar{X}\|_\alpha + 2 (\mathfrak{c}_M + 1) |Y_0|. \quad (2.35)$$

Let us introduce the shorthand

$$\varepsilon := (\tau \wedge T)^\alpha$$

and let us agree that, whenever we write *for ε small enough* we mean for $0 < \varepsilon \leq \varepsilon_0$ for a suitable $\varepsilon_0 > 0$ which depends on α, T, M_0, M, D . By (2.5), for ε small enough,

$$\|Y\|_{\infty, \tau} \leq |Y_0| + \varepsilon \|\delta Y\|_{\alpha, \tau} \leq |Y_0| + \frac{1}{5} \|\delta Y\|_{\alpha, \tau}, \quad (2.36)$$

hence to prove (2.35) we can focus on $\|\delta Y\|_{\alpha, \tau}$.

Recalling (2.4), let us define $Y^{[2]} := Z^{[2]} - \bar{Z}^{[2]}$. We are going to establish the following two relations, for ε small enough:

$$\frac{4}{5} \|\delta Y\|_{\alpha, \tau} \leq 2 M_0 \|\delta X - \delta \bar{X}\|_\alpha + \mathfrak{c}_M |Y_0| + \|Y^{[2]}\|_{\alpha, \tau}, \quad (2.37)$$

$$\|Y^{[2]}\|_{\alpha, \tau} \leq M_0 \|\delta X - \delta \bar{X}\|_\alpha + \frac{1}{2} |Y_0| + \frac{1}{5} \|\delta Y\|_{\alpha, \tau}. \quad (2.38)$$

Plugging (2.38) into (2.37) and applying (2.36), we obtain (2.35).

It remains to prove (2.37) and (2.38). We record some useful bounds. Let us set

$$\bar{\varepsilon} = \bar{\varepsilon}_{\alpha, D, M} := \frac{1}{2(K_{2\alpha} + 3)DM}. \quad (2.39)$$

We exploit the a priori estimate (2.13) from Theorem 2.7: by (2.32), we have

$$\text{for } \varepsilon = (\tau \wedge T)^\alpha \leq \bar{\varepsilon}: \quad \max\{\|\delta Z\|_{\alpha, \tau}, \|\delta \bar{Z}\|_{\alpha, \tau}\} \leq 2M_0M, \quad (2.40)$$

therefore

$$\|\delta\sigma(Z)\|_{\alpha, \tau} \leq \|\nabla\sigma\|_\infty \|\delta Z\|_{\alpha, \tau} \leq 2\|\nabla\sigma\|_\infty M_0M = 2M_0\mathfrak{c}_M, \quad (2.41)$$

and applying (2.5) and (2.32) we get, *for ε small enough*,

$$\|\sigma(Z)\|_{\infty, \tau} \leq |\sigma(Z_0)| + 3\varepsilon \|\delta\sigma(Z)\|_{\alpha, \tau} \leq M_0(1 + 6\mathfrak{c}_M\varepsilon) \leq 2M_0. \quad (2.42)$$

We can now prove (2.37). Defining $Y^{[2]} := Z^{[2]} - \bar{Z}^{[2]}$, we obtain from (2.4)

$$\begin{aligned} \delta Y_{st} &= \delta Z_{st} - \delta \bar{Z}_{st} = \sigma(Z_s) \delta X_{st} - \sigma(\bar{Z}_s) \delta \bar{X}_{st} + Y_{st}^{[2]} \\ &= \sigma(Z_s) (\delta X - \delta \bar{X})_{st} + (\sigma(Z_s) - \sigma(\bar{Z}_s)) \delta \bar{X}_{st} + Y_{st}^{[2]}, \end{aligned}$$

hence by (2.7) we can bound

$$\begin{aligned} \|\delta Y\|_{\alpha, \tau} &\leq \|\sigma(Z)\|_{\infty, \tau} \|\delta X - \delta \bar{X}\|_\alpha \\ &\quad + \|\delta \bar{X}\|_\alpha \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty, \tau} + \|Y^{[2]}\|_{\alpha, \tau}. \end{aligned} \quad (2.43)$$

Let us look at the second term in the RHS of (2.43): by (2.5)

$$\begin{aligned} \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty, \tau} &\leq \|\nabla\sigma\|_\infty \|Z - \bar{Z}\|_{\infty, \tau} \\ &\leq \|\nabla\sigma\|_\infty (|Y_0| + 3\varepsilon \|\delta Y\|_{\alpha, \tau}). \end{aligned} \quad (2.44)$$

Hence by (2.32) and (2.34) we get, *for ε small enough*,

$$\|\delta \bar{X}\|_\alpha \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty, \tau} \leq \mathfrak{c}_M |Y_0| + \frac{1}{5} \|\delta Y\|_{\alpha, \tau}. \quad (2.45)$$

Plugging this into (2.43) we then obtain, by (2.42),

$$\frac{4}{5} \|\delta Y\|_{\alpha, \tau} \leq 2M_0 \|\delta X - \delta \bar{X}\|_\alpha + \mathfrak{c}_M |Y_0| + \|Y^{[2]}\|_{\alpha, \tau}, \quad (2.46)$$

which proves (2.37).

We finally prove (2.38). Since $Y_{st}^{[2]} = Z_{st}^{[2]} - \bar{Z}_{st}^{[2]} = o(t-s)$, see (2.4), the weighted Sewing Bound (1.42) and (2.6) give

$$\|Y^{[2]}\|_{\alpha, \tau} \leq \varepsilon^{\gamma-1} \|Y^{[2]}\|_{\gamma\alpha, \tau} \leq K_{\gamma\alpha} \varepsilon^{\gamma-1} \|\delta Y^{[2]}\|_{\gamma\alpha, \tau}. \quad (2.47)$$

To estimate $\delta Y^{[2]} = \delta Z^{[2]} - \delta \bar{Z}^{[2]}$, note that by (2.4) and (1.33) we can write

$$\delta Y_{sut}^{[2]} = \delta\sigma(Z)_{su} (\delta X - \delta \bar{X})_{ut} + (\delta\sigma(Z) - \delta\sigma(\bar{Z}))_{su} \delta \bar{X}_{ut}, \quad (2.48)$$

hence by (2.8)

$$\|\delta Y^{[2]}\|_{\gamma\alpha, \tau} \leq \|\delta\sigma(Z)\|_{(\gamma-1)\alpha, \tau} \|\delta X - \delta \bar{X}\|_\alpha + \|\delta \bar{X}\|_\alpha \|\delta\sigma(Z) - \delta\sigma(\bar{Z})\|_{(\gamma-1)\alpha, \tau}. \quad (2.49)$$

The first term is easy to control: by (2.41), for ε small enough,

$$K_{\gamma\alpha} \varepsilon^{\gamma-1} \|\delta\sigma(Z)\|_{(\gamma-1)\alpha,\tau} \|\delta X - \delta\bar{X}\|_{\alpha} \leq M_0 \|\delta X - \delta\bar{X}\|_{\alpha}. \quad (2.50)$$

Let us now focus on the second term. By (2.19) we have, see also (2.28),

$$|\delta\sigma(Z)_{su} - \delta\sigma(\bar{Z})_{su}| \leq \|\nabla\sigma\|_{\infty} |\delta Y_{su}| + [\nabla\sigma]_{\mathcal{C}^{\gamma-1}} \{|\delta Z_{su}|^{\gamma-1} + |\delta\bar{Z}_{su}|^{\gamma-1}\} |Y_s|.$$

We apply (2.9) for $H = \delta Z$, $g = Y$ and $\bar{\tau} = (\bar{\varepsilon})^{1/\alpha}$ from (2.39):

$$\begin{aligned} \|\delta\sigma(Z) - \delta\sigma(\bar{Z})\|_{(\gamma-1)\alpha,\tau} &\leq \|\nabla\sigma\|_{\infty} \|\delta Y\|_{(\gamma-1)\alpha,\tau} + \\ &\quad + [\nabla\sigma]_{\mathcal{C}^{\gamma-1}} e^{\frac{T}{\bar{\tau}}} (\|\delta Z\|_{\alpha,\bar{\tau}}^{\gamma-1} + \|\delta\bar{Z}\|_{\alpha,\bar{\tau}}^{\gamma-1}) \|Y\|_{\infty,\tau} \\ &\leq D \|\delta Y\|_{\alpha,\tau} + 2(2M_0 M)^{\gamma-1} e^{\frac{T}{\bar{\tau}}} D \|Y\|_{\infty,\tau}, \end{aligned} \quad (2.51)$$

where we applied (2.40). Hence by (2.51), recalling (2.32), for ε small enough we obtain

$$K_{\gamma\alpha} \varepsilon^{\gamma-1} \|\delta\bar{X}\|_{\alpha} \|\delta\sigma(Z) - \delta\sigma(\bar{Z})\|_{(\gamma-1)\alpha,\tau} \leq \frac{1}{10} \|\delta Y\|_{\alpha,\tau} + \frac{1}{2} \|Y\|_{\infty,\tau}, \quad (2.52)$$

and since $\|Y\|_{\infty,\tau} \leq |Y_0| + \frac{1}{5} \|\delta Y\|_{\alpha,\tau}$, see (2.36), we obtain

$$K_{\gamma\alpha} \varepsilon^{\gamma-1} \|\delta\bar{X}\|_{\alpha} \|\delta\sigma(Z) - \delta\sigma(\bar{Z})\|_{(\gamma-1)\alpha,\tau} \leq \frac{1}{2} |Y_0| + \frac{1}{5} \|\delta Y\|_{\alpha,\tau}.$$

Finally, plugging this bound and (2.50) into (2.49) and (2.47), we obtain

$$\|Y^{[2]}\|_{\alpha,\tau} \leq M_0 \|\delta X - \delta\bar{X}\|_{\alpha} + \frac{1}{2} |Y_0| + \frac{1}{5} \|\delta Y\|_{\alpha,\tau},$$

which proves (2.38) and completes the proof. $\square$

Remark 2.13. An explicit choice for $\hat{\tau}$ in Theorem 2.12 is

$$\hat{\tau}^{\alpha} := \frac{e^{-\frac{T}{\bar{\tau}}}}{10(K_{2\alpha} + 3)(1 + M_0)(1 + D(M + M^2))}, \quad (2.53)$$

with $\bar{\tau} = \bar{\tau}_{\alpha,D,M}$ defined in (2.39). This is obtained by tracking all the points in the proof of Theorem 2.12 where $\varepsilon = (\tau \wedge T)^{\alpha}$ was assumed to be *small enough*: see Section 2.8 for the details.

2.6. EULER SCHEME AND LOCAL/GLOBAL EXISTENCE

In this section we discuss *global existence of solutions*, under the assumption that σ is globally γ -Hölder with $\gamma \in (\frac{1}{\alpha} - 1, 1]$, i.e. $[\sigma]_{\mathcal{C}^{\gamma}} < \infty$ (again without any boundedness assumption on σ). We also state a result of *local existence of solutions* for equation (2.3), where we only assume that σ is *locally* γ -Hölder with $\gamma \in (\frac{1}{\alpha} - 1, 1]$ (with no boundedness assumption on σ).

We fix $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^{\alpha}$ with $\alpha \in]\frac{1}{2}, 1]$ and a starting point $z_0 \in \mathbb{R}^k$. We split the proof in two parts: we first assume that $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is globally γ -Hölder, then we consider the case when σ is locally γ -Hölder.

First part: globally Hölder case.

We consider a finite set $\mathbb{T} = \{0 = t_1 < \dots < t_{\#\mathbb{T}}\} \subset \mathbb{R}_+$ and we define an approximate solution $Z = Z^\mathbb{T} = (Z_t)_{t \in \mathbb{T}}$ through the *Euler scheme*

$$Z_0 := z_0, \quad Z_{t_{i+1}} := Z_{t_i} + \sigma(Z_{t_i}) \delta X_{t_i, t_{i+1}} \quad \text{for } 1 \leq i \leq \#\mathbb{T} - 1. \quad (2.54)$$

Let us define the “remainder”

$$R_{st} := \delta Z_{st} - \sigma(Z_s) \delta X_{st} \quad \text{for } s < t \in \mathbb{T}. \quad (2.55)$$

We assume that σ is *globally γ -Hölder*, namely $[\sigma]_{C^\gamma} < \infty$, with $\gamma \in (\frac{1}{\alpha} - 1, 1]$. We set

$$\hat{\varepsilon}_{\alpha, \gamma, X, \sigma} := \frac{1}{2(C_{(\gamma+1)\alpha} + 5) \|\delta X\|_\alpha [\sigma]_{C^\gamma}}, \quad (2.56)$$

where the constant C_η is defined in (1.46). We prove the following *a priori estimates* on the Euler scheme (2.54), which are analogous to those in Theorem 2.7.

LEMMA 2.14. *If σ is globally γ -Hölder, namely $[\sigma]_{C^\gamma} < \infty$, with $\gamma \in (\frac{1}{\alpha} - 1, 1]$, then*

$$\|R\|_{(\gamma+1)\alpha}^\mathbb{T} \leq C_{(\gamma+1)\alpha} [\sigma]_{C^\gamma} (\|\delta Z\|_\alpha^\mathbb{T})^\gamma \|\delta X\|_\alpha, \quad (2.57)$$

$$\text{and for } \tau^{\gamma\alpha} \leq \hat{\varepsilon}_{\alpha, \gamma, X, \sigma}: \quad \|\delta Z\|_\alpha^\mathbb{T} \leq 1 \vee (2|\sigma(z_0)| \|\delta X\|_\alpha). \quad (2.58)$$

Proof. Since $\delta R_{sut} = (\sigma(Z_s) - \sigma(Z_u)) \delta X_{ut}$, recall (1.33), and since $R_{t_i t_{i+1}} = 0$ by (2.54), we can apply the discrete Sewing Bound (1.46) with $\eta = (\gamma + 1)\alpha > 1$ to get

$$\|R\|_{(\gamma+1)\alpha, \tau}^\mathbb{T} \leq C_{(\gamma+1)\alpha} \|\delta R\|_{(\gamma+1)\alpha, \tau}^\mathbb{T} \leq C_{(\gamma+1)\alpha} [\sigma]_{C^\gamma} (\|\delta Z\|_{\alpha, \tau}^\mathbb{T})^\gamma \|\delta X\|_\alpha. \quad (2.59)$$

We have proved (2.57).

We next prove (2.58). Recalling (2.55) we can bound, by (2.6) for $\|\cdot\|_{\gamma\alpha, \mathbb{T}_n}$,

$$\|\delta Z\|_{\alpha, \tau}^\mathbb{T} \leq \|\sigma(Z)\|_{\infty, \tau}^\mathbb{T} \|\delta X\|_\alpha + \tau^{\gamma\alpha} \|R\|_{(\gamma+1)\alpha, \tau}^\mathbb{T}.$$

By (1.49)

$$\|\sigma(Z)\|_{\infty, \tau}^\mathbb{T} \leq |\sigma(z_0)| + 5\tau^{\gamma\alpha} \|\delta\sigma(Z)\|_{\gamma\alpha, \tau}^\mathbb{T} \leq |\sigma(z_0)| + 5\tau^{\gamma\alpha} [\sigma]_{C^\gamma} (\|\delta Z\|_{\alpha, \tau}^\mathbb{T})^\gamma.$$

We thus obtain, combining the previous bounds,

$$\|\delta Z\|_{\alpha, \tau}^\mathbb{T} \leq |\sigma(z_0)| \|\delta X\|_\alpha + \{\tau^{\gamma\alpha} (C_{\gamma\alpha} + 5) [\sigma]_{C^\gamma} \|\delta X\|_\alpha\} (\|\delta Z\|_{\alpha, \tau}^\mathbb{T})^\gamma.$$

Now if $\|\delta Z\|_{\alpha, \tau}^\mathbb{T} \leq 1$ then (2.58) is proved, otherwise $(\|\delta Z\|_{\alpha, \tau}^\mathbb{T})^\gamma \leq \|\delta Z\|_{\alpha, \tau}^\mathbb{T}$ and then for τ as in (2.56) the term in brackets is less than $\frac{1}{2}$ and we obtain (2.58). $\square$

We can now prove the following

THEOREM 2.15. (GLOBAL EXISTENCE) *Let X be of class $\mathcal{C}^\alpha$, with $\alpha \in [\frac{1}{2}, 1]$, and let σ be globally γ -Hölder with $\gamma \in (\frac{1}{\alpha} - 1, 1]$, i.e. $[\sigma]_{C^\gamma} < \infty$. For every $z_0 \in \mathbb{R}^k$, with no restriction on $T > 0$, there exists a solution $(Z_t)_{t \in [0, T]}$ of (2.3) with $Z_0 = z_0$.*

Proof. Given $n \in \mathbb{N}$, we construct an approximate solution $Z^n = (Z_t^n)_{t \in \mathbb{T}_n}$ of (2.3) defined in the discrete set of times $\mathbb{T}_n := (\{i2^{-n} : i = 0, 1, \dots\} \cap [0, T]) \cup \{T\}$ through the *Euler scheme* (2.54).

$$Z_0^n := z_0, \quad Z_{t_{i+1}}^n := Z_{t_i}^n + \sigma(Z_{t_i}^n) \delta X_{t_i, t_{i+1}} \quad \text{for } t_i, t_{i+1} \in \mathbb{T}_n. \quad (2.60)$$

Let us define the “remainder”

$$R_{st}^n := \delta Z_{st}^n - \sigma(Z_s^n) \delta X_{st} \quad \text{for } s < t \in \mathbb{T}_n. \quad (2.61)$$

We fix $T > 0$ such that

We extend Z^n by linear interpolation to a continuous function defined on $[0, T]$, still denoted by Z^n . Given two points $t_i \leq s < t \leq t_{i+1}$ inside the same interval $[t_i, t_{i+1}]$ of the partition $\mathbb{T}_n$, since $\delta Z_{st}^n = \frac{t-s}{t_{i+1}-t_i} \delta Z_{t_i t_{i+1}}^n$, we can bound for $\alpha \in (0, 1]$

$$\frac{|\delta Z_{st}^n|}{(t-s)^\alpha} = \left(\frac{t-s}{t_{i+1}-t_i} \right)^{1-\alpha} \frac{|\delta Z_{t_i t_{i+1}}^n|}{(t_{i+1}-t_i)^\alpha} \leq \frac{|\delta Z_{t_i t_{i+1}}^n|}{(t_{i+1}-t_i)^\alpha}.$$

Given two points $s < t$ in different intervals, say $t_i \leq s \leq t_{i+1} \leq t_j \leq t \leq t_{j+1}$ for some $i < j$, by the triangle inequality we can bound $|\delta Z_{st}^n| \leq |\delta Z_{st_{i+1}}^n| + |\delta Z_{t_{i+1} t_j}^n| + |\delta Z_{t_j t}^n|$. Recalling (1.9) and (1.44), we then obtain $\|\cdot\|_\alpha \leq 3 \|\cdot\|_\alpha^{\mathbb{T}_n}$, hence by (2.58) we get

$$\|\delta Z^n\|_{\alpha, \tau} \leq 3 \vee (6 \|\sigma(z_0)\| \|\delta X\|_\alpha). \quad (2.62)$$

The family $(Z^n)_{n \in \mathbb{N}}$ is *equi-continuous* by (2.62) and *equi-bounded*, since $Z_0^n = z_0$ for all $n \in \mathbb{N}$, hence by the Arzelà-Ascoli Theorem it is *compact* in the space $C([0, T], \mathbb{R}^k)$. Let us denote by $Z: [0, T] \rightarrow \mathbb{R}^k$ any limit point. Plugging (2.58) into (2.57), by (2.61) we can write

$$\text{if } T^\alpha \leq \hat{\varepsilon}_{\alpha, X, \sigma}: \quad |\delta Z_{st}^n - \sigma(Z_s^n) \delta X_{st}| \leq c(z_0) (t-s)^{2\alpha} \quad \forall s < t \in \mathbb{T}_n, \quad (2.63)$$

where $c(z_0) := C_{(\gamma+1)\alpha}[\sigma] C^\gamma (3 \vee (6 \|\sigma(z_0)\| \|\delta X\|_\alpha))^\gamma \|\delta X\|_\alpha$. Letting $n \rightarrow \infty$ and observing that $\mathbb{T}_n \subseteq \mathbb{T}_{n+1}$, we see that (2.63) still holds with Z^n replaced by Z and $\mathbb{T}_n$ replaced by the set $\mathbb{T} := \bigcup_{\ell \in \mathbb{N}} \mathbb{T}_{2^\ell} = (\{\frac{i}{2^n} : i, n \in \mathbb{N}\} \cap [0, T]) \cup \{T\}$ of dyadic rationals:

$$\text{if } T^\alpha \leq \hat{\varepsilon}_{\alpha, X, \sigma}: \quad |\delta Z_{st} - \sigma(Z_s) \delta X_{st}| \leq c(z_0) (t-s)^{2\alpha} \quad \forall s < t \in \mathbb{T}.$$

Since $\mathbb{T}$ is dense in $[0, T]$ and Z is continuous, this bound extends to all $0 \leq s < t \leq T$, which shows that Z is a solution of (2.3). This completes the proof. $\square$

Second part: locally Lipschitz case.

We now assume that σ is *locally* γ -Hölder and we fix $z_0 \in \mathbb{R}^k$. We also fix $T > 0$ such that $T \leq \tilde{\varepsilon}_{\alpha, X, \sigma}(z_0)$, see (2.64), and we prove that there exists a solution $Z: [0, T] \rightarrow \mathbb{R}^k$ of (2.3) with $Z_0 = z_0$.

THEOREM 2.16. (LOCAL EXISTENCE) *Let X be of class C^α , with $\alpha \in [\frac{1}{2}, 1]$, and let σ be locally Lipschitz (e.g. of class C^1). For any $z_0 \in \mathbb{R}^k$ and for $T > 0$ small enough, i.e.*

$$T^\alpha \leq \tilde{\varepsilon}_{\alpha, X, \sigma}(z_0) := \frac{1}{2} \frac{1}{(C_{2\alpha} + 3) \|\delta X\|_\alpha \{1 + \sup_{|z-z_0| \leq |\sigma(z_0)|} |\nabla \sigma(z)|\}}, \quad (2.64)$$

there exists a solution $(Z_t)_{t \in [0, T]}$ of (2.3) with $Z_0 = z_0$.

Let $\tilde{\sigma}$ be a globally γ -Hölder function (depending on z_0) such that

$$\tilde{\sigma}(z) = \sigma(z) \quad \forall |z - z_0| \leq \sigma(z_0) \quad \text{and} \quad [\tilde{\sigma}]_{C^\gamma} = \sup_{|z - z_0| \leq \sigma(z_0)} |\nabla \sigma(z)|. \quad (2.65)$$

Since $T \leq \tilde{\varepsilon}_{\alpha, X, \sigma}(z_0) \leq \hat{\varepsilon}_{\alpha, X, \sigma}$, see (2.64) and (2.56), by the first part of the proof there exists a solution Z of (2.3) with $\tilde{\sigma}$ in place of σ and $Z_0 = z_0$. We will prove that

$$|Z_t - z_0| \leq \sigma(z_0) \quad \text{for all } t \in [0, T], \quad (2.66)$$

therefore $\tilde{\sigma}(Z_t) = \sigma(Z_t)$ for all $t \in [0, T]$, see (2.65). This means that Z is a solution of the original (2.3) with σ , which completes the proof of Theorem 2.16.

To prove (2.66), we apply the a priori estimate (2.13) with $\tau = \infty$: we note that $T \leq \tilde{\varepsilon}_{\alpha, X, \sigma}(z_0) \leq \varepsilon_{\alpha, X, \sigma}$ (see (2.64) and (2.12), and note that $C_{2\alpha} \geq K_{2\alpha}$), therefore

$$\|\delta Z\|_\alpha \leq 2 \|\delta X\|_\alpha |\sigma(z_0)|,$$

because $\tilde{\sigma}(z_0) = \sigma(z_0)$. Then for every $t \in [0, T]$ we can bound

$$|Z_t - z_0| \leq T^\alpha \|\delta Z\|_\alpha \leq 2 T^\alpha \|\delta X\|_\alpha |\sigma(z_0)| \leq |\sigma(z_0)|,$$

where the last inequality holds because $T^\alpha \leq \tilde{\varepsilon}_{\alpha, X, \sigma}(z_0) \leq (2 \|\delta X\|_\alpha)^{-1}$, see (2.64). This completes the proof of (2.66).

2.7. ERROR ESTIMATE IN THE EULER SCHEME

We suppose in this section that σ is of class C^2 with $\|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty < +\infty$.

THEOREM 2.17. *The Euler scheme converges at speed $n^{2\alpha-1}$.*

Proof. Let us set $z_i := \partial y_i / \partial y_0$, where $(y_i)_{i \geq 0}$ is defined by (2.60). Then

$$z_{i+1} = z_i + \nabla \sigma(y_i) z_i \delta X_{t_i t_{i+1}}, \quad i \geq 0.$$

This shows that the pair $(y_i, z_i)_{i \geq 0}$ satisfies a recurrence which is similar to (2.60) with a map Σ of class C^1 and therefore we can apply the above results to obtain that $|z_i| \leq \text{const}$. In particular the map $y_0 \rightarrow y_k$ is Lipschitz-continuous, uniformly over $k \geq 0$.

Let us call, for $k \geq 0$, $(z_\ell^{(k)})_{\ell \geq k}$ as the sequence which satisfies (2.60) but has initial value $z_k^{(k)} = y(t_k)$. Since $(y(t))_{t \geq 0}$ is a solution to (2.4), we have

$$|z_{k+1}^{(k)} - y(t_{k+1})| \lesssim n^{-2\alpha}.$$

Since the map $y_0 \rightarrow y_k$ is Lipschitz-continuous uniformly over $k \geq 0$, we have

$$|z_\ell^{(k)} - z_\ell^{(k+1)}| \lesssim |z_{k+1}^{(k)} - y(t_{k+1})| \lesssim n^{-2\alpha}, \quad \ell \geq k+1.$$

Therefore

$$|y_\ell - y(t_\ell)| = |z_\ell^{(0)} - z_\ell^{(\ell)}| \leq \sum_{k=0}^{\ell-1} |z_\ell^{(k)} - z_\ell^{(k+1)}| \lesssim \frac{\ell}{n^{2\alpha}} = \frac{t_\ell}{n^{2\alpha-1}} \rightarrow 0$$

as t_ℓ is bounded and $n \rightarrow \infty$. □

2.8. EXTRA: A VALUE FOR $\hat{\tau}$

We can give an explicit expression for $\hat{\tau} = \hat{\tau}_{M_0, M, T}$ in Theorem 2.12, by tracking all the points in the proof where τ is *small enough*, namely:

- for (2.36) we need $\tau^\alpha \leq \frac{1}{15}$;
- for (2.40) we need $\tau^\alpha \leq (\hat{\rho}_M)^\alpha := (2(K_{2\alpha} + 3)\mathfrak{c}_M)^{-1}$;
- for (2.42) we need $\tau^\alpha \leq (6\mathfrak{c}_M)^{-1}$, for (2.45) we need $\tau^\alpha \leq (15\mathfrak{c}_M)^{-1}$;
- for (2.50) we need $\tau^{(\gamma-1)\alpha} \leq (2K_{\gamma\alpha}\mathfrak{c}_M)^{-1}$;
- for (2.52) we need $\tau^{(\gamma-1)\alpha} \leq (10K_{\gamma\alpha}\mathfrak{c}_M)^{-1}$ (first term in the RHS) and also $\tau^{(\gamma-1)\alpha} \leq \left(K_{\gamma\alpha} e^{\frac{T}{\hat{\rho}_M}} M_0 M^2 \|\nabla^2 \sigma\|_\infty\right)^{-1}$ (second term in the RHS).

Since $\mathfrak{c}_M = M \|\nabla \sigma\|_\infty$, see (2.34), it is easy to check that all these constraints are satisfied for $0 < \tau \leq \hat{\tau}$ given by formula (2.53) in Remark 2.13.

CHAPTER 3

DIFFERENCE EQUATIONS: THE ROUGH CASE

We have so far considered the difference equation (2.3), that is

$$Z_t - Z_s = \sigma(Z_s) (X_t - X_s) + o(t - s), \quad 0 \leq s \leq t \leq T, \quad (3.1)$$

where X is given, Z is the unknown and $\sigma(\cdot)$ is sufficiently regular. This is a generalization of the differential equation $\dot{Z}_t = \sigma(Z_t) \dot{X}_t$ which is meaningful for non smooth X , as we showed in Chapter 2, where we proved *well-posedness* in the so-called *Young case*, i.e. assuming that $X \in \mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$.

However, the restriction $\alpha > \frac{1}{2}$ is a substantial limitation: in particular, we cannot take $X = B$ as a typical path of Brownian motion, which is in $\mathcal{C}^\alpha$ only for $\alpha < \frac{1}{2}$. For this reason, we show in this chapter how to *enrich* the difference equation (3.1) and prove *well-posedness when $X \in \mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{3}, \frac{1}{2}]$* , called the *rough case*. This will be applied to Brownian motion in the next Chapter 4, in order to obtain a robust formulation of classical *stochastic differential equations*.

NOTATION. Throughout this book we write $f_{st} \lesssim g_{st}$ to mean that $f_{st} \leq C g_{st}$ for all $0 \leq s \leq t \leq T$, where $C < \infty$ is a suitable random constant.

Remark 3.1. (YOUNG VS. ROUGH CASE) The restriction $\alpha > \frac{1}{2}$ for the study of the difference equation (3.1) has a substantial reason, namely *there is no solution to (3.1) for general $X \in \mathcal{C}^\alpha$ with $\alpha \leq \frac{1}{2}$* . Indeed, taking the “increment” δ of both sides of (3.1) and recalling (1.24) and (1.33), we obtain

$$(\sigma(Z_u) - \sigma(Z_s)) (X_t - X_u) = o(t - s) \quad \text{for } 0 \leq s \leq u \leq t \leq T. \quad (3.2)$$

If $X \in \mathcal{C}^\alpha$, for any $\alpha \in (0, 1]$, then we know from Lemma 2.6 that $Z \in \mathcal{C}^\alpha$, but not better in general (e.g. when $\sigma(\cdot) \equiv c$ is constant we have $Z = cX$), hence the LHS of (3.2) is $\lesssim (u - s)^\alpha (t - u)^\alpha \lesssim (t - s)^{2\alpha}$, but not better in general. This shows that the restriction $\alpha > \frac{1}{2}$ is generally necessary for (3.1) to have solutions.

3.1. ENHANCED TAYLOR EXPANSION

We fix $d, k \in \mathbb{N}$, a time horizon $T > 0$ and a sufficiently regular function $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$. Our goal is to give a meaning to the integral equation

$$Z_t = Z_0 + \int_0^t \sigma(Z_s) \dot{X}_s ds, \quad 0 \leq t \leq T, \quad (3.3)$$

where $Z: [0, T] \rightarrow \mathbb{R}^k$ is the unknown and $X: [0, T] \rightarrow \mathbb{R}^d$ is a non smooth path, more precisely $X \in \mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{3}, \frac{1}{2}]$.

The difference equation (3.1) is no longer enough, for the crucial reason that typically *it admits no solutions for $\alpha \leq \frac{1}{2}$* , see Remark 3.1. We are going to solve this problem by *enriching the RHS of (3.1)* in a suitable, but non canonical way: this leads to the key notion of *rough path* which is central in this book.

To provide motivation, suppose for the moment that X is continuously differentiable, so that (3.3) is meaningful. As we saw in (1.3), an integration yields for $s < t$

$$Z_t - Z_s - \sigma(Z_s)(X_t - X_s) = \int_s^t (\sigma(Z_u) - \sigma(Z_s)) \dot{X}_u du. \quad (3.4)$$

In Chapter 1 we observed that the integral is $o(t-s)$, which lead to the difference equation (3.1). More precisely, the integral is $O((t-s)^2)$ if $X \in C^1$ and σ is locally Lipschitz (note that $Z \in C^1$). The idea is now to go further, expanding the integral to get a more accurate local description, with a better remainder $O((t-s)^3)$.

To this purpose, we assume that σ is differentiable and we introduce the key function $\sigma_2: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^d)^*$ by

$$\sigma_2(z) := \nabla \sigma(z) \sigma(z), \quad \text{i.e.} \quad [\sigma_2(z)]_{j\ell}^i := \sum_{a=1}^k \frac{\partial \sigma_j^i}{\partial z_a}(z) \sigma_\ell^a(z). \quad (3.5)$$

Since $\frac{d}{dr} \sigma(Z_r) = \nabla \sigma(Z_r) \dot{Z}_r = \sigma_2(Z_r) \dot{X}_r$ by (3.3), we can write for $s < u$

$$\sigma(Z_u) - \sigma(Z_s) - \sigma_2(Z_s)(X_u - X_s) = \int_s^u (\sigma_2(Z_r) - \sigma_2(Z_s)) \dot{X}_r dr, \quad (3.6)$$

where for $z \in \mathbb{R}^d$ and $a \in \mathbb{R}^d$ we define $\sigma_2(z) a \in \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ by

$$[\sigma_2(z) a]_j^i = \sum_{\ell=1}^d [\sigma_2(z)]_{j\ell}^i a^\ell.$$

If we assume that σ_2 is continuous, then the right-hand side of (3.6) is $o(u-s)$ (recall that $X \in C^1$). Plugging this into (3.4), we then obtain

$$\begin{aligned} Z_t - Z_s &= \sigma(Z_s)(X_t - X_s) - \sigma_2(Z_s) \int_s^t (X_u - X_s) \otimes \dot{X}_u du \\ &= \int_s^t (\sigma(Z_r) - \sigma(Z_s) - \sigma_2(Z_s)(X_r - X_s)) \dot{X}_r dr, \end{aligned} \quad (3.7)$$

where now for $z \in \mathbb{R}^d$ and $B \in \mathbb{R}^d \otimes \mathbb{R}^d$ we define $\sigma_2(z) B \in \mathbb{R}^k$ by

$$[\sigma_2(z) B]^i = \sum_{\ell, m=1}^d [\sigma_2(z)]_{\ell m}^i B^{m\ell}. \quad (3.8)$$

Now, since the right-hand side of (3.6) is $o(u-s)$, the right-hand side of (3.7) is $o((t-s)^2)$.

In order to rewrite (3.7) more conveniently, we introduce the shorthands

$$\mathbb{X}_{st}^1 := X_t - X_s, \quad \mathbb{X}_{st}^2 := \int_s^t (X_r - X_s) \otimes \dot{X}_r dr, \quad 0 \leq s \leq t \leq T, \quad (3.9)$$

so that $\mathbb{X}^1: [0, T]_{\leq}^2 \rightarrow \mathbb{R}^d$ and $\mathbb{X}^2: [0, T]_{\leq}^2 \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$, see (1.7). More explicitly:

$$(\mathbb{X}_{st}^2)^{ij} := \int_s^t (X_r^i - X_s^i) \dot{X}_r^j dr, \quad i, j \in \{1, \dots, d\}.$$

We can thus rewrite (3.7), in the compact form

$$Z_t - Z_s = \sigma(Z_s) \mathbb{X}_{st}^1 + \sigma_2(Z_s) \mathbb{X}_{st}^2 + o(t - s), \quad 0 \leq s \leq t \leq T, \quad (3.10)$$

where for the product $\sigma_2(Z_s) \mathbb{X}_{st}^2$ we use the contraction rule (3.8), and we know that the $o(t - s)$ term is in fact $o((t - s)^2)$ but we are not going to use this further.

We have obtained an *enhanced Taylor expansion*: comparing with (3.1), we added a “second order term” containing $\mathbb{X}_{st}^2$. The idea is to take this new difference equation, that we call *rough difference equation*, as a generalized formulation of the differential equation (3.3), just as we did in Chapter 1 (see Section 1.2). However, there is a problem: the term $\mathbb{X}_{st}^2$ depends on the derivative $\dot{X}$, see (3.9), so it is not clearly defined for a non-differentiable X .

To overcome this problem, we will *choose* a suitable function $\mathbb{X}^2 = (\mathbb{X}_{st}^2)_{0 \leq s \leq t \leq T}$ playing the role of the integral (3.9) when X is not differentiable: this leads to the notion of *rough paths*, defined in the next section and studied in depth in Chapter 8. We will show in this chapter that rough paths are the key to a robust solution theory of rough difference equations when X of class $\mathcal{C}^\alpha$ with $\alpha \in (\frac{1}{3}, \frac{1}{2}]$.

3.2. ROUGH PATHS

Let us fix a path $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^\alpha$ with $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. Motivated by the previous section, we are going to reformulate the ill-posed integral equation (3.3) as the difference equation (3.10), which contains $\mathbb{X}^1$ and $\mathbb{X}^2$.

We can certainly define $\mathbb{X}_{st}^1 := X_t - X_s$ as in (3.9), but there is no canonical definition of $\mathbb{X}_{st}^2 = \int_s^t (X_r - X_s) \otimes \dot{X}_r dr$, since X may not be differentiable. We therefore *assign* a function $\mathbb{X}_{st}^2$ which satisfies *suitable properties*. Note that when X is continuously differentiable the function $\mathbb{X}^2$ in (3.9) satisfies:

- an algebraic identity known as *Chen’s relation*: for $0 \leq s \leq u \leq t \leq T$

$$\mathbb{X}_{st}^2 - \mathbb{X}_{su}^2 - \mathbb{X}_{ut}^2 = \mathbb{X}_{su}^1 \otimes \mathbb{X}_{ut}^1 = (X_u - X_s) \otimes (X_t - X_u), \quad (3.11)$$

which follows from (3.9) noting that

$$\mathbb{X}_{st}^2 - \mathbb{X}_{su}^2 - \mathbb{X}_{ut}^2 = \int_u^t (X_r - X_s) \otimes \dot{X}_r dr = (X_u - X_s) \otimes (X_t - X_u);$$

- the analytic bounds

$$|\mathbb{X}_{st}^1| \lesssim |t - s|, \quad |\mathbb{X}_{st}^2| \lesssim |t - s|^2, \quad (3.12)$$

which follow from the fact that $\dot{X}$ is bounded.

The algebraic relation (3.11) is still meaningful for non-differentiable X , while the analytic bounds (3.12) can naturally be adapted to the case of Hölder paths $X \in \mathcal{C}^\alpha$ by changing the exponents 1, 2 to $\alpha, 2\alpha$. This leads to the following key definition.

DEFINITION 3.2. (ROUGH PATHS) Fix $\alpha \in [\frac{1}{3}, \frac{1}{2}]$, $d \in \mathbb{N}$ and a path $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^\alpha$. An **α -rough path over X** is a pair $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ where the functions $\mathbb{X}^1: [0, T] \rightarrow \mathbb{R}^d$ and $\mathbb{X}^2: [0, T]^2 \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ satisfy, for $0 \leq s \leq u \leq t \leq T$:

- the algebraic relations

$$\mathbb{X}_{st}^1 = X_t - X_s, \quad \delta \mathbb{X}_{sut}^2 := \mathbb{X}_{st}^2 - \mathbb{X}_{su}^2 - \mathbb{X}_{ut}^2 = \mathbb{X}_{su}^1 \otimes \mathbb{X}_{ut}^1, \quad (3.13)$$

where the second identity is called *Chen's relation*;

- the analytic bounds

$$|\mathbb{X}_{st}^1| \lesssim |t - s|^\alpha, \quad |\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}. \quad (3.14)$$

We call $\mathcal{R}_{\alpha,d}(X)$ the set of d -dimensional α -rough paths $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ over X and $\mathcal{R}_{\alpha,d} = \bigcup_{X \in \mathcal{C}^\alpha} \mathcal{R}_{\alpha,d}(X)$ the set of all d -dimensional α -rough paths.

When X is of class C^1 , the choice (3.9) yields by (3.11)-(3.12) a α -rough path for any $\alpha \in (\frac{1}{3}, \frac{1}{2}]$ which we call the *canonical rough path*, see Section 8.7 below.

When $X = B$ is Brownian motion, the theory of stochastic integration provides a natural candidate for $\mathbb{X}^2$, in fact *multiple candidates* (think of Ito vs. Stratonovich integration), as we discuss in Chapter 4 below. Incidentally, this makes it clear that the construction of $\mathbb{X}^2$ is in general *non canonical*, i.e. there are multiple choices of $\mathbb{X}^2$ for a given path X . *This is a strength of the theory of rough paths*, since it allows to treat different non equivalent forms of integration.

Remark 3.3. The existence of rough paths over any given path X (i.e. the fact that $\mathcal{R}_{\alpha,d}(X) \neq \emptyset$) is a non trivial fact, which will be proved in Chapter 8.

Remark 3.4. ($\mathbb{X}^2$ AS A “PATH”) The two-parameters function $\mathbb{X}_{st}^2$ is determined by the one-parameter function

$$\mathbb{I}_t := \mathbb{X}_{0t}^2 + X_0 \otimes (X_t - X_0), \quad (3.15)$$

which intuitively describes the integral $\int_0^t X_r \otimes \dot{X}_r dr$. Indeed, we can write

$$\mathbb{X}_{st}^2 = \mathbb{I}_t - \mathbb{I}_s - X_s \otimes (X_t - X_s), \quad (3.16)$$

since $\mathbb{X}_{st}^2 = \mathbb{X}_{0t}^2 - \mathbb{X}_{0s}^2 - (X_s - X_0) \otimes (X_t - X_s)$ by Chen's relation (3.13).

Vice versa, given a function $\mathbb{I}: [0, T] \rightarrow \mathbb{R}^d$, if we *define* $\mathbb{X}^2$ by (3.16), then Chen's relation (3.13) is automatically satisfied (recall (1.33)). In order to satisfy the analytic bound in (3.14), we must require that

$$|\mathbb{I}_t - \mathbb{I}_s - X_s \otimes (X_t - X_s)| \lesssim (t - s)^{2\alpha}, \quad (3.17)$$

which is a natural estimate if $\mathbb{I}_t - \mathbb{I}_s$ should describe “ $= \int_s^t X_r \otimes \dot{X}_r dr$ ”.

Summarizing: given any path $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^\alpha$, it is equivalent to assign $\mathbb{X}^2: [0, T]^2 \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ satisfying (3.13)-(3.14) or to assign $\mathbb{I}: [0, T] \rightarrow \mathbb{R}^d$ satisfying (3.17), the correspondence being given by (3.15)-(3.16).

3.3. ROUGH DIFFERENCE EQUATIONS

Given a time horizon $T > 0$ and two dimensions $d, k \in \mathbb{N}$, let us fix:

- a path $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{3}, \frac{1}{2}]$;
- an α -rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ over X , see Definition 3.2;
- a *differentiable* function $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$, which lets us define the function

$$\sigma_2: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^d)^* \quad (\text{see (3.5)}).$$

Motivated by the previous discussions, see in particular (3.10), we study in this chapter the following *rough difference equation* for an unknown path $Z: [0, T] \rightarrow \mathbb{R}^k$:

$$\delta Z_{st} = \sigma(Z_s) \mathbb{X}_{st}^1 + \sigma_2(Z_s) \mathbb{X}_{st}^2 + o(t-s), \quad 0 \leq s \leq t \leq T, \quad (3.18)$$

where we recall the increment notation $\delta Z_{st} := Z_t - Z_s$ and the contraction rule (3.8), and we stress that $o(t-s)$ is *uniform* for $0 \leq s \leq t \leq T$, see Remark 1.1. In analogy with (2.3)-(2.4), a solution of (3.18) is a path $Z: [0, T] \rightarrow \mathbb{R}^k$ such that

$$Z_{st}^{[3]} := \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma_2(Z_s) \mathbb{X}_{st}^2 = o(t-s). \quad (3.19)$$

We stress that the rough difference equation (3.18) is a generalization of the integral equation (3.3), as we show in the next result.

PROPOSITION 3.5. *If X and σ are of class C^1 and σ_2 is continuous (e.g. if σ is of class C^1), then any solution Z to the integral equation (3.3) satisfies the difference equation (3.18) for the canonical rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ in (3.9).*

Proof. If $X \in C^1$, then $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ defined in (3.9) is an α -rough path over X for any $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, as we showed in (3.11)-(3.12). Given a solution Z of (3.3), if σ_2 is continuous we derived the Taylor expansion (3.10), hence (3.18) holds. $\square$

We now state *local and global existence, uniqueness of solutions and continuity of the solution map* for the rough difference equation (3.18) under natural assumptions on σ and σ_2 , summarizing the main results of this chapter. We refer to the next sections for more precise and quantitative results.

PROPOSITION 3.6. *Let $z_0 \in \mathbb{R}^d$. We suppose that σ and σ_2 are of class C^1 and globally Lipschitz, namely $\|\nabla \sigma\|_\infty + \|\nabla \sigma_2\|_\infty < +\infty$. Let $D := \max\{1, \|\nabla \sigma\|_\infty, \|\nabla \sigma_2\|_\infty\}$ and $M > 0$.*

There exists $T_{M,D,\alpha} > 0$ such that, for all $T \in (0, T_{M,D,\alpha})$ and $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha,d}$ such that $\|\mathbb{X}^1\|_\alpha + \|\mathbb{X}^2\|_{2\alpha} \leq M$, there exists a solution Z to (3.19) on the interval $[0, T]$ such that $Z_0 = z_0$ and

$$\|Z\|_\alpha \leq 15 M (|\sigma(z_0)| + |\sigma_2(z_0)|). \quad (3.20)$$

The proof of this Proposition, based on a discretization argument, is postponed to section 3.9 below.

We are going to use the Sewing Bound (1.27), its weighted version (1.42) and its discrete formulation (1.46).

3.4. SET-UP

We recall that the *weighted semi-norms* $\|\cdot\|_{\eta,\tau}$ are defined in (1.34)-(1.35). We are going to use the various properties that we recalled in Section 2.2, see in particular (2.5), (2.6) and (2.7)-(2.8), as well as the natural generalization

$$\text{if } F_{sut} = G_{su} H_{ut} \text{ then } \|F\|_{3\eta,\tau} \begin{cases} \leq \|G\|_{2\eta,\tau} \|H\|_{\eta} , \\ \leq \|G\|_{\eta,\tau} \|H\|_{2\eta} . \end{cases} \quad (3.21)$$

In all these bounds, whenever there is a product, *only one factor gets the weighted semi-norm, while the other factor gets the ordinary semi-norm*. We sometimes need to introduce an additional weight, which is possible applying (2.9).

In Chapter 2 a key tool to study the Young difference equation (2.4) was the estimate on the “difference of increments” in Lemma 2.8. This tool is still crucial in this chapter, but we will need an additional ingredient that we now present.

LEMMA 3.7. (TAYLOR IDENTITY) *Let $z_1, z_2 \in \mathbb{R}^k$ and $x \in \mathbb{R}^d$. If $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is of class C^1 , defining $\sigma_2: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^d)^*$ by (3.5) and setting $\delta z_{12} := z_2 - z_1$, we have the identities*

$$\begin{aligned} & \sigma(z_2) - \sigma(z_1) - \sigma_2(z_1) x \\ &= \nabla \sigma(z_1) (\delta z_{12} - \sigma(z_1) x) + \int_0^1 [(\nabla \sigma(z_1 + r \delta z_{12}) - \nabla \sigma(z_1)) \delta z_{12}] dr, \end{aligned} \quad (3.22)$$

and

$$\begin{aligned} \sigma(z_2) - \sigma(z_1) - \sigma_2(z_1) x &= \int_0^1 [(\sigma_2(z_1 + r \delta z_{12}) - \sigma_2(z_1)) x] dr \\ &+ \int_0^1 [\nabla \sigma(z_1 + r \delta z_{12}) (\delta z_{12} - \sigma(z_1) x)] dr \\ &- \int_0^1 \nabla \sigma(z_1 + r \delta z_{12}) \left(\int_0^r [\nabla \sigma(z_1 + v \delta z_{12}) \delta z_{12} x] dv \right) dr. \end{aligned} \quad (3.23)$$

Proof. The first formula is based on elementary manipulations and on the fact that

$$\sigma(z_2) - \sigma(z_1) = \int_0^1 [\nabla \sigma(z_1 + r \delta z_{12}) \delta z_{12}] dr.$$

For the second formula, setting $\delta z := \delta z_{12}$ for short, we similarly write

$$\begin{aligned} \sigma(z_2) - \sigma(z_1) &= \int_0^1 [\nabla \sigma(z_1 + r \delta z) \delta z] dr \\ &= \int_0^1 [\nabla \sigma(z_1 + r \delta z) (\delta z - \sigma(z_1) x)] dr + \underbrace{\int_0^1 [\nabla \sigma(z_1 + r \delta z) \sigma(z_1) x] dr}_A \end{aligned}$$

and then, recalling the definition (3.5) of σ_2 ,

$$A = \int_0^1 [\sigma_2(z_1 + r \delta z) x] dr - \underbrace{\int_0^1 [\nabla \sigma(z_1 + r \delta z) (\sigma(z_1 + r \delta z) - \sigma(z_1)) x] dr}_B.$$

Finally

$$B = \int_0^1 \nabla \sigma(z_1 + r \delta z) \left(\int_0^r [\nabla \sigma(z_1 + v \delta z) \delta z x] dv \right) dr$$

from which (3.23) follows easily. $\square$

We will see below that (3.22) is useful for the comparison between *two solutions*, as in the proofs of uniqueness (Theorem 3.10) and continuity of the solution map (Theorem 3.11), while (3.23) is well suited for a priori estimates on a *single solution* (Theorem 3.9) or on a discretization scheme (Lemma 3.13).

3.5. A PRIORI ESTIMATES

In this section we prove *a priori estimates* for solutions of the rough difference equation (3.18) for *globally Lipschitz* σ and σ_2 , i.e. $\|\nabla \sigma\|_\infty < \infty$ and $\|\nabla \sigma_2\|_\infty < \infty$. A sufficient condition is that $\sigma, \nabla \sigma, \nabla^2 \sigma$ are bounded, see (3.5), but it is interesting that *boundedness of σ is not necessary* (think of the case of linear σ).

Given a solution Z of (3.18), we define the “remainders” $Z^{[3]}$ and $Z^{[2]}$ by

$$Z_{st}^{[3]} = \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma_2(Z_s) \mathbb{X}_{st}^2, \quad Z_{st}^{[2]} = \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1. \quad (3.24)$$

Let us first show, by easy arguments, that any solution Z of (3.18) has the same Hölder regularity $\mathcal{C}^\alpha$ of the driving path X (in analogy with Lemmas 1.2 and 2.6), and that the “level 2 remainder” $Z_{st}^{[2]}$ is in $C_2^{2\alpha}$, that is $|Z_{st}^{[2]}| \lesssim (t-s)^{2\alpha}$.

LEMMA 3.8. (HÖLDER REGULARITY) *Let σ be of class C^1 and let Z be a solution of (3.18). There is a constant $C = C(Z) < \infty$ such that*

$$\begin{cases} |Z_{st}^{[2]}| \leq C |\mathbb{X}_{st}^2| + o(t-s), \\ |\delta Z_{st}| \leq C (|\mathbb{X}_{st}^1| + |\mathbb{X}_{st}^2|) + o(t-s), \end{cases} \quad 0 \leq s \leq t \leq T. \quad (3.25)$$

In particular, if $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ is an α -rough path, then $Z^{[2]} \in C_2^{2\alpha}$ and Z is of class $\mathcal{C}^\alpha$.

Proof. If $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ is an α -rough path, then by the first bound in (3.25) we have $|Z_{st}^{[2]}| \lesssim (t-s)^{2\alpha} + o(t-s) \lesssim (t-s)^{2\alpha}$, that is $Z^{[2]} \in C_2^{2\alpha}$. Similarly, the second bound in (3.25) gives $|\delta Z_{st}| \lesssim (t-s)^\alpha + (t-s)^{2\alpha} + o(t-s) \lesssim (t-s)^\alpha$, that is Z is of class $\mathcal{C}^\alpha$.

It remains to prove (3.25). This follows by (3.18) with $C := \sup_{0 \leq s \leq T} \{|\sigma(Z_s)| + |\sigma_2(Z_s)|\}$, so we need to show that $C < \infty$. Since σ and σ_2 are continuous (because σ is of class C^1), it is enough to prove that Z is bounded: $\sup_{0 \leq t \leq T} |Z_t| < \infty$.

Arguing as in the proof of Lemma 1.2, we fix $\bar{\delta} > 0$ such that $|o(t-s)| \leq 1$ for all $0 \leq s \leq t \leq T$ with $|t-s| \leq \bar{\delta}$. Since $[0, T]$ is a finite union of intervals $[\bar{s}, \bar{t}]$ with $\bar{t} - \bar{s} \leq \bar{\delta}$, we may focus on one such interval: by (3.18) we can bound

$$\sup_{t \in [\bar{s}, \bar{t}]} |Z_t| \leq |Z_{\bar{s}}| + |\sigma(Z_{\bar{s}})| \sup_{t \in [\bar{s}, \bar{t}]} |\mathbb{X}_{st}^1| + |\sigma_2(Z_{\bar{s}})| \sup_{t \in [\bar{s}, \bar{t}]} |\mathbb{X}_{st}^2| + 1 < \infty.$$

This completes the proof that $\sup_{0 \leq t \leq T} |Z_t| < \infty$. $\square$

We next get to our main a priori estimates, showing in particular that the “level 3 remainder” $Z_{st}^{[3]}$ is in $C_2^{3\alpha}$, that is $|Z_{st}^{[3]}| \lesssim |t-s|^{3\alpha}$. Let us first record a useful computation: recalling (1.24) and (1.33), by $\delta \circ \delta = 0$ and (3.13), we have

$$\begin{aligned} \delta Z_{sut}^{[3]} &= Z_{st}^{[3]} - Z_{su}^{[3]} - Z_{ut}^{[3]} \\ &= \underbrace{(\sigma(Z_u) - \sigma(Z_s) - \sigma_2(Z_s) \mathbb{X}_{su}^1)}_{B_{su}} \mathbb{X}_{ut}^1 + (\sigma_2(Z_u) - \sigma_2(Z_s)) \mathbb{X}_{ut}^2. \end{aligned} \quad (3.26)$$

THEOREM 3.9. (ROUGH A PRIORI ESTIMATES) *Let X be of class $\mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{3}, \frac{1}{2}]$ and let $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ be an α -rough path over X . Let σ and σ_2 be globally Lipschitz.*

For any solution Z of (3.18), recalling the “remainders” $Z^{[3]}$ and $Z^{[2]}$ from (3.24), we have $Z^{[3]} \in C_2^{3\alpha}$: more precisely, for any $\tau > 0$,

$$\|Z^{[3]}\|_{3\alpha, \tau} \leq K_{3\alpha} c'_{\alpha, \mathbb{X}, \sigma} (\|\delta Z\|_{\alpha, \tau} + \|Z^{[2]}\|_{2\alpha, \tau}), \quad (3.27)$$

where we recall that $K_{3\alpha} = (1 - 2^{1-3\alpha})^{-1}$ and we define the constant

$$c'_{\alpha, \mathbb{X}, \sigma} := \|\nabla \sigma\|_\infty \|\mathbb{X}^1\|_\alpha + \|\nabla \sigma_2\|_\infty \|\mathbb{X}^2\|_{2\alpha} + (\|\nabla \sigma\|_\infty^2 + \|\nabla \sigma_2\|_\infty) \|\mathbb{X}^1\|_\alpha^2. \quad (3.28)$$

Moreover, if either T or τ is small enough, we have

$$\begin{aligned} \|\delta Z\|_{\alpha, \tau} + \|Z^{[2]}\|_{2\alpha, \tau} &\leq 2 (\sigma(Z_0) \|\mathbb{X}^1\|_\alpha + \sigma_2(Z_0) \|\mathbb{X}^2\|_{2\alpha}) \\ &\text{for } (T \wedge \tau)^\alpha \leq \varepsilon'_{\alpha, \mathbb{X}, \sigma}, \end{aligned} \quad (3.29)$$

where we set

$$\varepsilon'_{\alpha, \mathbb{X}, \sigma} := \frac{1}{4(K_{3\alpha} + 3)(c'_{\alpha, \mathbb{X}, \sigma} + 1)}. \quad (3.30)$$

Proof. Let us prove (3.27). Since $3\alpha > 1$ and $Z_{st}^{[3]} = o(t-s)$, see (3.19), we can apply the weighted Sewing Bound (1.42) which gives $\|Z^{[3]}\|_{3\alpha, \tau} \leq K_{3\alpha} \|\delta Z^{[3]}\|_{3\alpha, \tau}$. It remains to estimate $\delta Z^{[3]}$ from (3.26): applying (3.21) we can write

$$\|\delta Z^{[3]}\|_{3\alpha, \tau} \leq \|B\|_{2\alpha, \tau} \|\mathbb{X}^1\|_\alpha + \|\delta \sigma_2(Z)\|_{\alpha, \tau} \|\mathbb{X}^2\|_{2\alpha}. \quad (3.31)$$

We now focus on B_{su} from (3.26): by (3.23) we have

$$\begin{aligned} B_{su} &= \int_0^1 [(\sigma_2(Z_s + u \delta Z_{su}) - \sigma_2(Z_s)) \mathbb{X}_{su}^1] du + \int_0^1 [\nabla \sigma(Z_s + u \delta Z_{su}) Z_{su}^{[2]}] du \\ &\quad - \int_0^1 \nabla \sigma(Z_s + u \delta Z_{su}) \left(\int_0^u [\nabla \sigma(Z_s + v \delta Z_{su}) \delta Z_{su} \mathbb{X}_{su}^1] dv \right) du, \end{aligned}$$

so that, by (2.8),

$$\|B\|_{2\alpha, \tau} \leq (\|\nabla \sigma_2\|_\infty + \|\nabla \sigma\|_\infty^2) \|\mathbb{X}^1\|_\alpha \|\delta Z\|_{\alpha, \tau} + \|\nabla \sigma\|_\infty \|Z^{[2]}\|_{2\alpha, \tau}. \quad (3.32)$$

We can plug this estimate into (3.31), together with the elementary bound

$$\|\delta \sigma_2(Z)\|_{\alpha, \tau} \leq \|\nabla \sigma_2\|_\infty \|\delta Z\|_{\alpha, \tau}. \quad (3.33)$$

Recalling that $\|Z^{[3]}\|_{3\alpha,\tau} \leq K_{3\alpha} \|\delta Z^{[3]}\|_{3\alpha,\tau}$, we have proved (3.27)-(3.28).

We next prove (3.29), for which we need to estimate $Z^{[2]}$ and δZ . Writing $Z_{st}^{[2]} = \sigma_2(Z_s) \mathbb{X}_{st}^2 + Z_{st}^{[3]}$ and setting $\varepsilon := (\tau \wedge T)^\alpha$ for short, we can bound by (2.6) and (2.7)

$$\|Z^{[2]}\|_{2\alpha,\tau} \leq \|\sigma_2(Z)\|_{\infty,\tau} \|\mathbb{X}^2\|_{2\alpha} + \varepsilon \|Z^{[3]}\|_{3\alpha,\tau}.$$

By (2.5) we have $\|\sigma_2(Z)\|_{\infty,\tau} \leq \sigma_2(Z_0) + 3\varepsilon \|\delta\sigma_2(Z)\|_{\alpha,\tau}$ and we can bound $\|\delta\sigma_2(Z)\|_{\alpha,\tau}$ by (3.33). Applying (3.27) and recalling (3.28), we then obtain

$$\begin{aligned} \|Z^{[2]}\|_{2\alpha,\tau} &\leq \sigma_2(Z_0) \|\mathbb{X}^2\|_{2\alpha} + \varepsilon (K_{3\alpha} + 3) c'_{\alpha,\mathbb{X},\sigma} (\|\delta Z\|_{\alpha,\tau} + \|Z^{[2]}\|_{2\alpha,\tau}) \\ &\leq \sigma_2(Z_0) \|\mathbb{X}^2\|_{2\alpha} + \frac{1}{4} \frac{\varepsilon}{\varepsilon'_{\alpha,\mathbb{X},\sigma}} (\|\delta Z\|_{\alpha,\tau} + \|Z^{[2]}\|_{2\alpha,\tau}), \end{aligned} \quad (3.34)$$

where we recall that $\varepsilon'_{\alpha,\mathbb{X},\sigma}$ is defined in (3.30).

Similarly, writing $\delta Z_{st} = \sigma(Z_s) \mathbb{X}_{st}^1 + Z_{st}^{[2]}$ we can bound, by (2.6) and (2.7),

$$\|\delta Z\|_{\alpha,\tau} \leq \|\sigma(Z)\|_{\infty,\tau} \|\mathbb{X}^1\|_{\alpha} + \varepsilon \|Z^{[2]}\|_{2\alpha,\tau},$$

and since $\|\sigma(Z)\|_{\infty,\tau} \leq \sigma(Z_0) + 3\varepsilon \|\delta\sigma(Z)\|_{\alpha,\tau} \leq \sigma(Z_0) + 3\varepsilon \|\nabla\sigma\|_{\infty} \|\delta Z\|_{\alpha,\tau}$ we get, recalling (3.28),

$$\begin{aligned} \|\delta Z\|_{\alpha,\tau} &\leq \sigma(Z_0) \|\mathbb{X}^1\|_{\alpha} + 3\varepsilon c'_{\alpha,\mathbb{X},\sigma} \|\delta Z\|_{\alpha,\tau} + \varepsilon \|Z^{[2]}\|_{2\alpha,\tau} \\ &\leq \sigma(Z_0) \|\mathbb{X}^1\|_{\alpha} + \frac{1}{4} \frac{\varepsilon}{\varepsilon'_{\alpha,\mathbb{X},\sigma}} \|\delta Z\|_{\alpha,\tau} + \varepsilon \|Z^{[2]}\|_{2\alpha,\tau}. \end{aligned} \quad (3.35)$$

Finally, for $\varepsilon \leq \varepsilon'_{\alpha,\mathbb{X},\sigma}$ (hence $\varepsilon \leq \frac{1}{4}$, see (3.28)), by (3.34) and (3.35) we obtain

$$\|\delta Z\|_{\alpha,\tau} + \|Z^{[2]}\|_{2\alpha,\tau} \leq \sigma(Z_0) \|\mathbb{X}^1\|_{\alpha} + \sigma_2(Z_0) \|\mathbb{X}^2\|_{2\alpha} + \frac{1}{2} (\|\delta Z\|_{\alpha,\tau} + \|Z^{[2]}\|_{2\alpha,\tau}).$$

Since $\|\delta Z\|_{\alpha,\tau} + \|Z^{[2]}\|_{2\alpha,\tau} < \infty$ by Lemma 3.8, we have proved (3.29). $\square$

3.6. UNIQUENESS

In this section we prove uniqueness of solutions of (3.18) under the assumption that $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is of class C^γ with $\gamma > \frac{1}{\alpha}$ (e.g. it suffices that σ is of class C^3). This implies that σ_2 from (3.5) is of class C^1 with locally $(\gamma - 2)$ -Hölder gradient $\nabla\sigma_2$. We stress that σ and σ_2 are *not* required to be bounded.

THEOREM 3.10. (UNIQUENESS) *Let X be of class $\mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, let $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ be an α -rough path over X , and let σ be of class C^γ with $\gamma > \frac{1}{\alpha}$ (e.g. if σ is of class C^3). Then for every $z_0 \in \mathbb{R}^k$ there exists at most one solution Z to (3.18) such that $Z_0 = z_0$.*

Proof. Let us fix two solutions $Z, \bar{Z}$ of (3.18) and define their difference

$$Y := Z - \bar{Z}.$$

Our goal is to show that, for $\tau > 0$ small, we have $\|Y\|_{\infty, \tau} \leq 2|Y_0|$. In particular, if $Z_0 = \bar{Z}_0$, then $Y_0 = 0$ and therefore $\|Y\|_{\infty, \tau} = 0$, i.e. $Z = \bar{Z}$, which completes the proof.

We know by (2.5) that

$$\|Y\|_{\infty, \tau} \leq |Y_0| + 3\tau^\alpha \|\delta Y\|_{\alpha, \tau}. \quad (3.36)$$

With some abuse of notation, we denote by $Y_{st}^{[2]} := Z_{st}^{[2]} - \bar{Z}_{st}^{[2]}$ and $Y_{st}^{[3]} := Z_{st}^{[3]} - \bar{Z}_{st}^{[3]}$ the “differences of remainders”, recall (3.24), so that we can write

$$\delta Y_{st} = (\sigma(Z_s) - \sigma(\bar{Z}_s)) \mathbb{X}_{st}^1 + Y_{st}^{[2]}, \quad (3.37)$$

$$Y_{st}^{[2]} = (\sigma_2(Z_s) - \sigma_2(\bar{Z}_s)) \mathbb{X}_{st}^2 + Y_{st}^{[3]}. \quad (3.38)$$

We are going to show that, for $\tau > 0$ small enough, the following bounds hold:

$$\|\delta Y\|_{\alpha, \tau} \leq c_1 \|Y\|_{\infty, \tau} + \tau^\alpha \|Y^{[2]}\|_{2\alpha, \tau}, \quad (3.39)$$

$$\|Y^{[2]}\|_{2\alpha, \tau} \leq c_2 \|Y\|_{\infty, \tau} + \tau^{(\gamma-2)\alpha} \|Y^{[3]}\|_{\gamma\alpha, \tau}, \quad (3.40)$$

$$\|Y^{[3]}\|_{\gamma\alpha, \tau} \leq c_3 \|Y\|_{\infty, \tau} + c'_3 \tau^{(\gamma-2)\alpha} \|Y^{[3]}\|_{\gamma\alpha, \tau}, \quad (3.41)$$

for suitable constants c_i, c'_i that may depend on $Z, \bar{Z}, \mathbb{X}^1, \mathbb{X}^2, \sigma$, but not on τ .

We can easily complete the proof, assuming (3.39)-(3.41): if we fix $\tau > 0$ small enough so that $c'_3 \tau^{(\gamma-2)\alpha} < \frac{1}{2}$, by (3.41) we have $\|Y^{[3]}\|_{\gamma\alpha, \tau} \leq 2c_3 \|Y\|_{\infty, \tau}$; plugging this into (3.40) and taking $\tau > 0$ small, we obtain $\|Y^{[2]}\|_{2\alpha, \tau} \leq 2c_2 \|Y\|_{\infty, \tau}$, which plugged into (3.39) yields $\|\delta Y\|_{\alpha, \tau} \leq 2c_1 \|Y\|_{\infty, \tau}$, for $\tau > 0$ is small enough. Finally, by (3.36) we obtain, for $\tau > 0$ small, our goal $\|Y\|_{\infty, \tau} \leq 2|Y_0|$.

It remains to prove (3.39)-(3.41). Recalling (2.18), let us define the constants

$$C'_1 := C_{\nabla\sigma, \|Z\|_\infty \vee \|\bar{Z}\|_\infty}, \quad C''_1 := C_{\nabla^2\sigma, \|Z\|_\infty \vee \|\bar{Z}\|_\infty}, \quad C'_2 := C_{\nabla\sigma_2, \|Z\|_\infty \vee \|\bar{Z}\|_\infty},$$

$$C'''_1 := \sup \left\{ \frac{|\nabla^2\sigma(x) - \nabla^2\sigma(y)|}{|x - y|^{\gamma-2}} : |x|, |y| \leq \|Z\|_\infty \vee \|\bar{Z}\|_\infty \right\},$$

$$C''_2 := \sup \left\{ \frac{|\nabla\sigma_2(x) - \nabla\sigma_2(y)|}{|x - y|^{\gamma-2}} : |x|, |y| \leq \|Z\|_\infty \vee \|\bar{Z}\|_\infty \right\}.$$

(Note that $\|Z\|_\infty, \|\bar{Z}\|_\infty < \infty$ because $Z, \bar{Z}$ are continuous, see Lemma 3.8.)

We can prove (3.39) and (3.40) arguing as in the proof of Theorem 2.9, see (2.24) and (2.25). Indeed, from (3.37) we can bound, by (2.6) and (2.7),

$$\begin{aligned} \|\delta Y\|_{\alpha, \tau} &\leq \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty, \tau} \|\mathbb{X}^1\|_\alpha + \tau^\alpha \|Y^{[2]}\|_{2\alpha, \tau} \\ &\leq C'_1 \|Y\|_{\infty, \tau} \|\mathbb{X}^1\|_\alpha + \tau^\alpha \|Y^{[2]}\|_{2\alpha, \tau}, \end{aligned} \quad (3.42)$$

because $|\sigma(Z_t) - \sigma(\bar{Z}_t)| \leq C'_1 |Z_t - \bar{Z}_t|$, hence (3.39) holds with $c_1 = C'_1 \|\mathbb{X}^1\|_\alpha$. Similarly, by (3.38) we can bound

$$\begin{aligned} \|Y^{[2]}\|_{2\alpha, \tau} &\leq \|\sigma_2(Z) - \sigma_2(\bar{Z})\|_{\infty, \tau} \|\mathbb{X}^2\|_{2\alpha} + \tau^{(\gamma-2)\alpha} \|Y^{[3]}\|_{\gamma\alpha, \tau} \\ &\leq C'_2 \|Y\|_{\infty, \tau} \|\mathbb{X}^2\|_{2\alpha} + \tau^{(\gamma-2)\alpha} \|Y^{[3]}\|_{\gamma\alpha, \tau}, \end{aligned} \quad (3.43)$$

because $|\sigma_2(Z_t) - \sigma_2(\bar{Z}_t)| \leq C'_2 |Z_t - \bar{Z}_t|$, hence also (3.40) holds with $c_2 = C'_2 \|\mathbb{X}^2\|_{2\alpha}$.

We finally prove (3.41). Since $Y_{st}^{[3]} = Z_{st}^{[3]} - \bar{Z}_{st}^{[3]} = o(t-s)$, see (3.19), we can bound $Z^{[3]}$ by its increment $\delta Z^{[3]}$ through the weighted Sewing Bound (1.42):

$$\|Y^{[3]}\|_{\gamma\alpha,\tau} \leq K_{\gamma\alpha} \|\delta Y^{[3]}\|_{\gamma\alpha,\tau}. \quad (3.44)$$

We are going to prove the following estimate:

$$\|\delta Y^{[3]}\|_{\gamma\alpha,\tau} \leq \tilde{c}_3 \|Y\|_{\infty,\tau} + \tilde{c}_3' \|\delta Y\|_{\alpha,\tau} + \tilde{c}_3'' \|Y^{[2]}\|_{2\alpha,\tau}, \quad (3.45)$$

for suitable constants $\tilde{c}_3, \tilde{c}_3', \tilde{c}_3''$ that depend on $Z, \bar{Z}, \mathbb{X}^1, \mathbb{X}^2, \sigma$, *but not on* τ . Plugging the estimates (3.39) and (3.40) (that we already proved) for $\|\delta Y\|_{\alpha,\tau}$ and $\|Y^{[2]}\|_{2\alpha,\tau}$, we obtain (3.41) for suitable (explicit) constants c_3, c_3' .

Let us then prove (3.45). Recalling (3.26), for $0 \leq s \leq u \leq t \leq T$ we can write

$$\delta Y_{sut}^{[3]} = (B_{su} - \bar{B}_{su}) \mathbb{X}_{ut}^1 + (\delta\sigma_2(Z) - \delta\sigma_2(\bar{Z}))_{su} \mathbb{X}_{ut}^2,$$

where $B_{su} := \sigma(Z_u) - \sigma(Z_s) - \sigma_2(Z_s) \mathbb{X}_{su}^1$ and similarly for $\bar{B}_{su}$, hence by (3.21)

$$\|\delta Y^{[3]}\|_{\gamma\alpha,\tau} \leq \|B - \bar{B}\|_{(\gamma-1)\alpha,\tau} \|\mathbb{X}\|_{\alpha} + \|\delta\sigma_2(Z) - \delta\sigma_2(\bar{Z})\|_{(\gamma-2)\alpha,\tau} \|\mathbb{X}^2\|_{2\alpha}. \quad (3.46)$$

To obtain (3.45) we need to show that $\|B - \bar{B}\|_{(\gamma-1)\alpha,\tau}$ and $\|\delta\sigma_2(Z) - \delta\sigma_2(\bar{Z})\|_{(\gamma-2)\alpha,\tau}$ can be bounded by *linear combinations of* $\|Y\|_{\infty,\tau}$, $\|\delta Y\|_{\alpha,\tau}$ and $\|Y^{[2]}\|_{2\alpha,\tau}$.

We start from $\|\delta\sigma_2(Z) - \delta\sigma_2(\bar{Z})\|_{(\gamma-2)\alpha,\tau}$, which can be bounded as in (2.29):

$$\|\delta\sigma_2(Z) - \delta\sigma_2(\bar{Z})\|_{(\gamma-2)\alpha,\tau} \leq C_2' \|\delta Y\|_{\alpha,\tau} + C_2'' \{\|\delta Z\|_{\alpha}^{\gamma-1} + \|\delta \bar{Z}\|_{\alpha}^{\gamma-1}\} \|Y\|_{\infty,\tau}.$$

We next focus on $\|B - \bar{B}\|_{(\gamma-1)\alpha,\tau}$, which we are going to estimate by the following explicit linear combination of $\|Y\|_{\infty,\tau}$, $\|\delta Y\|_{\alpha,\tau}$ and $\|Y^{[2]}\|_{2\alpha,\tau}$:

$$\begin{aligned} \|B - \bar{B}\|_{(\gamma-1)\alpha,\tau} &\leq C_1'' \|Y\|_{\infty,\tau} \|Z^{[2]}\|_{2\alpha} + C_1' \|Y^{[2]}\|_{2\alpha,\tau} \\ &\quad + C_1'' \|\delta Y\|_{\alpha,\tau} \|\delta Z\|_{\alpha} + 2 C_1''' \|Y\|_{\infty,\tau} \|\delta Z\|_{\alpha}^2 \\ &\quad + C_1'' \|\delta \bar{Z}\|_{\alpha} \|\delta Y\|_{\alpha,\tau}, \end{aligned} \quad (3.47)$$

which completes the proof of (3.45) when plugged into (3.46).

It only remains to prove (3.47). Recalling (3.24), it follows by (3.22) that

$$\begin{aligned} B_{su} &:= \sigma(Z_u) - \sigma(Z_s) - \sigma_2(Z_s) \mathbb{X}_{su}^1 \\ &= \nabla\sigma(Z_s) Z_{su}^{[2]} + \underbrace{\int_0^1 (\nabla\sigma(Z_u + r \delta Z_{su}) - \nabla\sigma(Z_u)) \delta Z_{su} dr}_{F_{su}}, \end{aligned}$$

and likewise for $\bar{B}_{su}$ (with $\bar{F}_{su}$ defined similarly), therefore

$$|B_{su} - \bar{B}_{su}| \leq |\nabla\sigma(Z_s) Z_{su}^{[2]} - \nabla\sigma(\bar{Z}_s) \bar{Z}_{su}^{[2]}| + \int_0^1 |F_{su} \delta Z_{su} - \bar{F}_{su} \delta \bar{Z}_{su}| dr. \quad (3.48)$$

By the elementary estimate $|ab - \bar{a}\bar{b}| = |ab - \bar{a}b + \bar{a}b - \bar{a}\bar{b}| \leq |a - \bar{a}| |b| + |\bar{a}| |b - \bar{b}|$, that we apply repeatedly, we can bound

$$\begin{aligned} |\nabla\sigma(Z_s) Z_{su}^{[2]} - \nabla\sigma(\bar{Z}_s) \bar{Z}_{su}^{[2]}| &\leq |\nabla\sigma(Z_s) - \nabla\sigma(\bar{Z}_s)| |Z_{su}^{[2]}| + |\nabla\sigma(\bar{Z}_s)| |Z_{su}^{[2]} - \bar{Z}_{su}^{[2]}| \\ &\leq C_1'' |Y_s| |Z_{su}^{[2]}| + C_1' |Y_{su}^{[2]}|, \end{aligned}$$

and note that by (2.7) we obtain the first line in the RHS of (3.47).

To complete the proof of (3.47), we look at the second term in the RHS of (3.48):

$$\begin{aligned} |F_{su} \delta Z_{su} - \bar{F}_{su} \delta \bar{Z}_{su}| &\leq |F_{su} - \bar{F}_{su}| |\delta Z_{su}| + |\bar{F}_{su}| |\delta Z_{su} - \delta \bar{Z}_{su}| \\ &\leq |F_{su} - \bar{F}_{su}| |\delta Z_{su}| + C_1'' r |\delta \bar{Z}_{su}| |\delta Y_{su}|, \end{aligned} \quad (3.49)$$

because $|\bar{F}_{su}| \leq C_1'' r |\delta \bar{Z}_{su}|$. We then see, applying (2.8), that the last term in (3.49) produces the third line in (3.47). Finally, by (2.19) we estimate

$$\begin{aligned} |F_{su} - \bar{F}_{su}| &= |(\nabla \sigma(Z_u + r \delta Z_{su}) - \nabla \sigma(Z_u)) - (\nabla \sigma(\bar{Z}_u + r \delta \bar{Z}_{su}) - \nabla \sigma(\bar{Z}_u))| \\ &\leq C_1'' r |\delta Y_{su}| + C_1''' \{ |r \delta Z_{su}|^{\gamma-2} + |r \delta \bar{Z}_{su}|^{\gamma-2} \} |Y_s|. \end{aligned}$$

We obtain by (2.7) for $0 \leq r \leq 1$

$$\|F - \bar{F}\|_{(\gamma-2)\alpha, \tau} \leq C_1'' \|\delta Y\|_{\alpha, \tau} + 2 C_1''' \|Y\|_{\infty, \tau} \|\delta Z\|_{\alpha}^{\gamma-2}.$$

Applying again (2.8), we finally see that the first term in (3.49) yields the second line in (3.47), which completes the proof. $\square$

3.7. CONTINUITY OF THE SOLUTION MAP

In this section we assume that σ has bounded first, second and third derivatives, while σ_2 has bounded first and second derivatives:

$$\|\nabla \sigma\|_{\infty}, \|\nabla^2 \sigma\|_{\infty}, \|\nabla^3 \sigma\|_{\infty} < \infty, \quad \|\nabla \sigma_2\|_{\infty}, \|\nabla^2 \sigma_2\|_{\infty} < \infty. \quad (3.50)$$

(We stress that no boundedness assumption is made on σ and σ_2 .) Under these assumptions, given any time horizon $T > 0$, any starting point $Z_0 \in \mathbb{R}^k$ and any α -rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ with $\frac{1}{3} < \alpha \leq \frac{1}{2}$, we have *global existence and uniqueness* of solutions $Z: [0, T] \rightarrow \mathbb{R}^k$ to (3.18) (as we will prove in Theorem 3.12).

Denoting by $\mathcal{R}_{\alpha, d}$ the space of d -dimensional α -rough paths $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$, that we endow with the norm $\|\mathbb{X}^1\|_{\alpha} + \|\mathbb{X}^2\|_{2\alpha}$ we can thus consider the *solution map*:

$$\begin{aligned} \Phi: \mathbb{R}^k \times \mathcal{R}_{\alpha, d} &\longrightarrow \mathcal{C}^{\alpha} \\ (Z_0, \mathbb{X}) &\longmapsto Z := \begin{cases} \text{unique solution of (3.18) for } t \in [0, T] \\ \text{starting from } Z_0 \end{cases}. \end{aligned} \quad (3.51)$$

We prove the highly non-trivial result that this map is *locally Lipschitz*. In the space $\mathcal{C}^{\alpha}$ of Hölder functions we work with the weighted norm $\|f\|_{\infty, \tau} + \|\delta f\|_{\alpha, \tau}$, which is equivalent to the usual norm $\|f\|_{\mathcal{C}^{\alpha}} := \|f\|_{\infty} + \|\delta f\|_{\alpha}$, see Remark 1.15.

THEOREM 3.11. (CONTINUITY OF THE SOLUTION MAP) *Let σ and σ_2 satisfy (3.50) (with no boundedness assumption on the functions σ and σ_2). Then, for any $T > 0$ and $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, the solution map $(Z_0, \mathbb{X}) \mapsto Z$ in (3.51) is locally Lipschitz.*

More explicitly, given any $M_0, M, D < \infty$, if we assume that

$$\max \{ \|\nabla \sigma\|_{\infty}, \|\nabla^2 \sigma\|_{\infty}, \|\nabla^3 \sigma\|_{\infty}, \|\nabla \sigma_2\|_{\infty}, \|\nabla^2 \sigma_2\|_{\infty} \} \leq D, \quad (3.52)$$

and we consider starting points $Z_0, \bar{Z}_0 \in \mathbb{R}^d$ and rough paths $\mathbb{X}, \bar{\mathbb{X}} \in \mathcal{C}^{\alpha}$ with

$$\max \{ |\sigma(Z_0)|, |\sigma_2(Z_0)|, |\sigma(\bar{Z}_0)|, |\sigma_2(\bar{Z}_0)| \} \leq M_0, \quad (3.53)$$

$$\max \{ \|\mathbb{X}^1\|_{\alpha}, \|\mathbb{X}^2\|_{2\alpha}, \|\bar{\mathbb{X}}^1\|_{\alpha}, \|\bar{\mathbb{X}}^2\|_{2\alpha} \} \leq M, \quad (3.54)$$

then the corresponding solutions $Z = (Z_s)_{s \in [0, T]}$, $\bar{Z} = (\bar{Z}_s)_{s \in [0, T]}$ of (3.18) satisfy

$$\begin{aligned} & \|Z - \bar{Z}\|_{\infty, \tau} + \|\delta Z - \delta \bar{Z}\|_{\alpha, \tau} + \|Z^{[2]} - \bar{Z}^{[2]}\|_{2\alpha, \tau} \\ & \leq \mathfrak{C}'_M |Z_0 - \bar{Z}_0| + 30 M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}). \end{aligned} \quad (3.55)$$

provided τ satisfies $0 < \tau \wedge T \leq \hat{\tau}'$ for a suitable $\hat{\tau}' = \hat{\tau}'_{\alpha, T, D, M_0, M} > 0$, where we set

$$\mathfrak{C}'_M := 16 \{(\|\nabla \sigma\|_{\infty} + \|\nabla \sigma_2\|_{\infty}) M + 1\} \leq 32 (D M + 1).$$

Proof. It is convenient to define the constant

$$\mathfrak{C}'_M := (\|\nabla \sigma\|_{\infty} + \|\nabla \sigma_2\|_{\infty}) M \leq 2 D M. \quad (3.56)$$

Let Z and $\bar{Z}$ be two solutions of (3.18) with respective routh paths $\mathbb{X}$ and $\bar{\mathbb{X}}$. Defining $Y := Z - \bar{Z}$ and $Y^{[2]} := Z^{[2]} - \bar{Z}^{[2]}$, see (3.24), we rewrite our goal (3.55) as

$$\begin{aligned} \|Y\|_{\infty, \tau} + \|\delta Y\|_{\alpha, \tau} + \|Y^{[2]}\|_{2\alpha, \tau} & \leq 16 (\mathfrak{C}'_M + 1) |Y_0| \\ & + 30 M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}). \end{aligned} \quad (3.57)$$

Throughout the proof we use the shorthand

$$\varepsilon := (\tau \wedge T)^{\alpha} \quad (3.58)$$

and we write *for ε small enough* to mean *for all $0 < \varepsilon < \varepsilon_0$, for a suitable ε_0 depending on α, T, M_0, M, D* . We claim that the following estimates hold for δY and $Y^{[2]}$:

$$\|\delta Y\|_{\alpha, \tau} \leq \mathfrak{C}'_M \|Y\|_{\infty, \tau} + 2 M_0 \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \varepsilon \|Y^{[2]}\|_{2\alpha, \tau}, \quad (3.59)$$

$$\|Y^{[2]}\|_{2\alpha, \tau} \leq \mathfrak{C}'_M \|Y\|_{\infty, \tau} + 2 M_0 \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} + \varepsilon \|Y^{[3]}\|_{3\alpha, \tau}, \quad (3.60)$$

and, moreover, *for ε small enough* the following estimate holds for $Y^{[3]} := Z^{[3]} - \bar{Z}^{[3]}$:

$$\varepsilon \|Y^{[3]}\|_{3\alpha, \tau} \leq \|Y\|_{\infty, \tau} + M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}) + \|\delta Y\|_{\alpha, \tau} + \frac{1}{4} \|Y^{[2]}\|_{\alpha, \tau}. \quad (3.61)$$

It is now elementary (but tedious) to deduce our goal (3.57). Plugging (3.61) into (3.60) we obtain $\|Y^{[2]}\|_{2\alpha, \tau} \leq (\dots) + \frac{1}{4} \|Y^{[2]}\|_{2\alpha, \tau}$ which yields $\|Y^{[2]}\|_{2\alpha, \tau} \leq \frac{4}{3} (\dots)$ (since $\|Y^{[2]}\|_{2\alpha, \tau} < \infty$ by Lemma 3.8). Making $(\dots)$ explicit, we get

$$\|Y^{[2]}\|_{2\alpha, \tau} \leq 2 (\mathfrak{C}'_M + 1) \|Y\|_{\infty, \tau} + 4 M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}) + 2 \|\delta Y\|_{\alpha, \tau} \quad (3.62)$$

which plugged into (3.59) yields, *for ε small enough* (it suffices that $\varepsilon \leq \frac{1}{4}$),

$$\|\delta Y\|_{\alpha, \tau} \leq 3 (\mathfrak{C}'_M + 1) \|Y\|_{\infty, \tau} + 6 M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}), \quad (3.63)$$

and looking back at (3.62) we obtain

$$\|Y^{[2]}\|_{2\alpha, \tau} \leq 8 (\mathfrak{C}'_M + 1) \|Y\|_{\infty, \tau} + 16 M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}), \quad (3.64)$$

so that, overall,

$$\begin{aligned} \|Y\|_{\infty, \tau} + \|\delta Y\|_{\alpha, \tau} + \|Y^{[2]}\|_{2\alpha, \tau} & \leq 12 (\mathfrak{C}'_M + 1) \|Y\|_{\infty, \tau} \\ & + 22 M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}). \end{aligned} \quad (3.65)$$

It only remains to make $\|Y\|_{\infty,\tau}$ explicit. Since $\|Y\|_{\infty,\tau} \leq |Y_0| + 3\varepsilon \|\delta Y\|_{\alpha,\tau}$ by (2.5), for ε small enough (more precisely for $\varepsilon \leq \frac{1}{36(\mathfrak{c}'_M + 1)}$) we can bound

$$(\mathfrak{c}'_M + 1) \|Y\|_{\infty,\tau} \leq (\mathfrak{c}'_M + 1) |Y_0| + \frac{1}{12} \|\delta Y\|_{\alpha,\tau}, \quad (3.66)$$

which inserted into (3.63) yields

$$\|\delta Y\|_{\alpha,\tau} \leq 4(\mathfrak{c}'_M + 1) |Y_0| + 8M_0 (\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}).$$

Plugging this into (3.66), and then (3.66) into (3.65), we obtain our goal (3.57).

It remains to prove (3.59), (3.60) and (3.61). We first state some useful bounds that will be used repeatedly. Recalling (3.52) and (3.28)-(3.30), let us define

$$\bar{\tau} = \bar{\tau}_{\alpha,D,M} := \frac{1}{\{4(K_{3\alpha} + 3)(2(D^2 + D)(M^2 + M) + 1)\}^{1/\alpha}}, \quad (3.67)$$

By the a priori estimate (3.29) we can then bound

$$\text{for } \varepsilon = (\tau \wedge T)^\alpha \leq \bar{\tau}^\alpha: \quad \|\delta Z\|_{\alpha,\tau} + \|Z^{[2]}\|_{2\alpha,\tau} \leq 4M_0 M, \quad (3.68)$$

hence

$$\max\{\|\delta\sigma(Z)\|_{\alpha,\tau}, \|\delta\sigma_2(Z)\|_{\alpha,\tau}\} \leq \max\{\|\nabla\sigma\|_\infty, \|\nabla\sigma_2\|_\infty\} \|\delta Z\|_{\alpha,\tau} \leq 4M_0 \mathfrak{c}'_M, \quad (3.69)$$

which implies that, by (2.5) and for ε small enough,

$$\max\{\|\sigma(Z)\|_{\infty,\tau}, \|\sigma_2(Z)\|_{\infty,\tau}\} \leq M_0 + 3\varepsilon 4M_0 \mathfrak{c}'_M \leq 2M_0.$$

We record the following simple bound, for any Lipschitz function f ,

$$\|f(Z) - f(\bar{Z})\|_{\infty,\tau} \leq \|\nabla f\|_\infty \|Z - \bar{Z}\|_{\infty,\tau} = \|\nabla f\|_\infty \|Y\|_{\infty,\tau}. \quad (3.70)$$

We will also use a number of times the elementary estimate, for $a, b, \bar{a}, \bar{b} \in \mathbb{R}$,

$$|ab - \bar{a}\bar{b}| = |ab - a\bar{b} + a\bar{b} - \bar{a}\bar{b}| \leq |a| |b - \bar{b}| + |\bar{b}| |a - \bar{a}|. \quad (3.71)$$

We can now prove (3.59). Since $\delta Y_{st} = \delta Z_{st} - \delta \bar{Z}_{st} = \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma(\bar{Z}_s) \bar{\mathbb{X}}_{st}^1 + Y_{st}^{[2]}$, see (3.24) for Z and $\bar{Z}$, by (2.7) and (3.53)-(3.54) we get, applying (3.71),

$$\begin{aligned} \|\delta Y\|_{\alpha,\tau} &\leq \|\sigma(Z)\|_{\infty,\tau} \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha + \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty,\tau} \|\bar{\mathbb{X}}^1\|_\alpha + \|Y^{[2]}\|_{\alpha,\tau} \\ &\leq 2M_0 \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha + \|\sigma(Z) - \sigma(\bar{Z})\|_{\infty,\tau} M + \varepsilon \|Y^{[2]}\|_{2\alpha,\tau}, \end{aligned}$$

because $\|Y^{[2]}\|_{\alpha,\tau} \leq \varepsilon \|Y^{[2]}\|_{2\alpha,\tau}$ by (2.6) (recall the definition (3.58) of ε). Applying (3.70) with $f = \sigma$ and recalling $\mathfrak{c}'_M$ from (3.56), we obtain (3.59).

The proof of (3.60) is similar. Since $Z_{st}^{[3]} = Z_{st}^{[2]} - \sigma_2(Z_s) \mathbb{X}_{st}^2$ and similarly for $\bar{Z}^{[3]}$, see (3.24), we can write $Y_{st}^{[2]} = Z_{st}^{[2]} - \bar{Z}_{st}^{[2]} = \sigma_2(Z_s) \mathbb{X}_{st}^2 - \sigma_2(\bar{Z}_s) \bar{\mathbb{X}}_{st}^2 + Y_{st}^{[3]}$, therefore

$$\begin{aligned} \|Y^{[2]}\|_{2\alpha,\tau} &\leq \|\sigma_2(Z)\|_{\infty,\tau} \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} + \|\sigma_2(Z) - \sigma_2(\bar{Z})\|_{\infty,\tau} \|\bar{\mathbb{X}}^2\|_{2\alpha} + \|Y^{[3]}\|_{2\alpha,\tau} \\ &\leq 2M_0 \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} + \|\sigma_2(Z) - \sigma_2(\bar{Z})\|_{\infty,\tau} M + \varepsilon \|Y^{[3]}\|_{3\alpha,\tau}, \end{aligned}$$

since $\|Y^{[3]}\|_{2\alpha,\tau} \leq \varepsilon \|Y^{[3]}\|_{3\alpha,\tau}$ by (2.6). Applying (3.70) for $f = \sigma_2$ we obtain (3.60).

We finally prove (3.61). Since $Y_{st}^{[3]} = Z_{st}^{[3]} - \bar{Z}_{st}^{[3]} = o(t-s)$, see (3.19), the weighted Sewing Bound (1.42) yields

$$\|Y^{[3]}\|_{3\alpha,\tau} \leq K_{3\alpha} \|\delta Y^{[3]}\|_{3\alpha,\tau}, \quad (3.72)$$

hence we can focus on $\delta Y^{[3]} = \delta Z^{[3]} - \delta \bar{Z}^{[3]}$. Let us recall (3.26): for $0 \leq s \leq u \leq t \leq T$

$$\delta Z_{sut}^{[3]} = \underbrace{(\sigma(Z_u) - \sigma(Z_s) - \sigma_2(Z_s) \mathbb{X}_{su}^1)}_{B_{su}} \mathbb{X}_{ut}^1 + \delta \sigma_2(Z)_{su} \mathbb{X}_{ut}^2,$$

and analogously for $\delta \bar{Z}^{[3]}$ and $\bar{B}_{su}$, therefore by (3.71) and (3.21) we obtain

$$\begin{aligned} \|\delta Y^{[3]}\|_{3\alpha,\tau} &\leq \|B\|_{2\alpha,\tau} \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha + \|B - \bar{B}\|_{2\alpha,\tau} \|\bar{\mathbb{X}}^1\|_{\alpha,\tau} \\ &\quad + \|\delta \sigma_2(Z)\|_{\alpha,\tau} \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} + \|\delta \sigma_2(Z) - \delta \sigma_2(\bar{Z})\|_{\alpha,\tau} \|\bar{\mathbb{X}}^2\|_{2\alpha}. \end{aligned} \quad (3.73)$$

It remains to estimate the four terms in the RHS: in view of (3.72), relation (3.61) is proved if we show that, *for ε small enough*,

$$\varepsilon K_{3\alpha} \|B\|_{2\alpha,\tau} \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha \leq M_0 \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha, \quad (3.74)$$

$$\varepsilon K_{3\alpha} \|B - \bar{B}\|_{2\alpha,\tau} \|\bar{\mathbb{X}}^1\|_{\alpha,\tau} \leq \frac{1}{2} (\|Y\|_{\infty,\tau} + \|\delta Y\|_{\alpha,\tau}) + \frac{1}{4} \|Y^{[2]}\|_{2\alpha,\tau}, \quad (3.75)$$

$$\varepsilon K_{3\alpha} \|\delta \sigma_2(Z)\|_{\alpha,\tau} \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} \leq M_0 \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}, \quad (3.76)$$

$$\varepsilon K_{3\alpha} \|\delta \sigma_2(Z) - \delta \sigma_2(\bar{Z})\|_{\alpha,\tau} \|\bar{\mathbb{X}}^2\|_{2\alpha} \leq \frac{1}{2} (\|Y\|_{\infty,\tau} + \|\delta Y\|_{\alpha,\tau}). \quad (3.77)$$

We first deal with (3.76) and (3.77), then we focus on (3.74) and (3.75).

Proving (3.76) is very simple: since $\|\delta \sigma_2(Z)\|_{\alpha,\tau} \leq 4 M_0 \mathbf{c}'_M$ by (3.69), we see that (3.76) holds *for ε small enough*. To prove (3.77), note that by (2.51) we have

$$\|\delta \sigma(Z) - \delta \sigma(\bar{Z})\|_{(\gamma-1)\alpha,\tau} \leq \|\nabla \sigma\|_\infty \|\delta Y\|_{\alpha,\tau} + 4 M_0 M [\sigma]_{C^{\gamma-1}} \|Y\|_{\infty,\tau}.$$

Applying (3.54) and (3.68) we obtain

$$\|\delta \sigma_2(Z) - \delta \sigma_2(\bar{Z})\|_{\alpha,\tau} \|\bar{\mathbb{X}}^2\|_{2\alpha} \leq \|\nabla \sigma_2\|_\infty M \|\delta Y\|_{\alpha,\tau} + e^{\frac{T}{\tau}} \|\nabla^2 \sigma_2\|_\infty 8 M_0 M^2 \|Y\|_{\infty,\tau},$$

which shows that (3.77) holds *for ε small enough*.

Let us now prove (3.74). By (3.22) we have, for $0 \leq s \leq t \leq T$,

$$B_{st} = \underbrace{\nabla \sigma(Z_s) Z_{st}^{[2]}}_{E_{st}} + \underbrace{\int_0^1 [(\nabla \sigma(Z_s + r \delta Z_{st}) - \nabla \sigma(Z_s)) \delta Z_{st}] dr}_{F_{st}} \quad (3.78)$$

and similarly for $\bar{E}_{st}$ and $\bar{F}_{st}$. In particular, recalling (3.68), we get

$$\begin{aligned} \|B\|_{2\alpha,\tau} &\leq \|\nabla \sigma\|_\infty \|Z^{[2]}\|_{2\alpha,\tau} + \|\nabla^2 \sigma\|_\infty \|\delta Z\|_{\alpha,\tau}^2 \\ &\leq \|\nabla \sigma\|_\infty 4 M_0 M + \|\nabla^2 \sigma\|_\infty (4 M_0 M)^2, \end{aligned}$$

hence we see that (3.74) holds *for ε small enough*.

We finally prove (3.75), which is a bit tedious. In view of (3.78), we first consider

$$E_{st} - \bar{E}_{st} = (\nabla \sigma(Z_s) - \nabla \sigma(\bar{Z}_s)) Z_{st}^{[2]} + \nabla \sigma(\bar{Z}_s) (Z_{st}^{[2]} - \bar{Z}_{st}^{[2]}).$$

Applying (2.9) with $H = Z^{[2]}$ and $\bar{\tau}$ from (3.67), we obtain

$$\|E - \bar{E}\|_{2\alpha, \tau} \leq \|\nabla\sigma(Z) - \nabla\sigma(\bar{Z})\|_{\infty, \tau} e^{\frac{T}{\bar{\tau}}} \|Z^{[2]}\|_{2\alpha, \bar{\tau}} + \|\nabla\sigma\|_{\infty} \|Y^{[2]}\|_{2\alpha, \tau}.$$

By (3.70) with $f = \nabla\sigma$ and the a priori estimate (3.68) we obtain

$$\|E - \bar{E}\|_{2\alpha, \tau} \leq \|\nabla^2\sigma\|_{\infty} \|Y\|_{\infty, \tau} e^{\frac{T}{\bar{\tau}}} 4M_0M + \|\nabla\sigma\|_{\infty} \|Y^{[2]}\|_{2\alpha, \tau}. \quad (3.79)$$

We then consider $F_{st} - \bar{F}_{st}$. By (2.19), for $0 \leq r \leq 1$ we can estimate

$$\begin{aligned} & |(\nabla\sigma(Z_s + r\delta Z_{st}) - \nabla\sigma(Z_s)) - (\nabla\sigma(\bar{Z}_s + r\delta\bar{Z}_{st}) - \nabla\sigma(\bar{Z}_s))| |\delta Z_{st}| \\ & \leq \|\nabla^2\sigma\|_{\infty} |\delta Y_{st}| |\delta Z_{st}| + \|\nabla^3\sigma\|_{\infty} \max_{0 \leq u \leq 1} \{(1-u)|Y_s| + u|Y_t|\} |\delta Z_{st}|^2, \end{aligned}$$

as well as

$$|\nabla\sigma(Z_s + r\delta Z_{st}) - \nabla\sigma(Z_s)| |\delta Z_{st} - \delta\bar{Z}_{st}| \leq \|\nabla^2\sigma\|_{\infty} |\delta Z_{st}| |\delta Y_{st}|.$$

We can then estimate $F_{st} - \bar{F}_{st}$ from (3.78) as in (3.71): applying (2.9) twice with $H = \delta Z$ and $H = (\delta Z)^2$, always with $\bar{\tau}$ from (3.67), and recalling (3.68), we obtain

$$\begin{aligned} \|F - \bar{F}\|_{2\alpha, \tau} & \leq 2\|\nabla^2\sigma\|_{\infty} \|\delta Y\|_{\alpha, \tau} e^{\frac{T}{\bar{\tau}}} \|\delta Z\|_{\alpha, \bar{\tau}} + \|\nabla^3\sigma\|_{\infty} \|Y\|_{\infty, \tau} e^{\frac{T}{\bar{\tau}}} \|\delta Z\|_{\alpha, \bar{\tau}}^2 \\ & \leq e^{\frac{T}{\bar{\tau}}} \{8M_0M \|\nabla^2\sigma\|_{\infty} \|\delta Y\|_{\alpha, \tau} + (4M_0M)^2 \|\nabla^3\sigma\|_{\infty} \|Y\|_{\infty, \tau}\}. \end{aligned} \quad (3.80)$$

Since $\|B - \bar{B}\|_{2\alpha, \tau} \leq \|E - \bar{E}\|_{2\alpha, \tau} + \|F - \bar{F}\|_{2\alpha, \tau}$ in view of (3.78), we see by (3.79) and (3.80) that (3.75) holds for ε small enough. The proof is complete. $\square$

3.8. GLOBAL EXISTENCE AND UNIQUENESS

Let us suppose that $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is of class C^3 with $\|\nabla\sigma\|_{\infty} + \|\nabla\sigma_2\|_{\infty} < +\infty$.

THEOREM 3.12. *Let $\alpha > \frac{1}{3}$. If $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is of class C^3 with $\|\nabla\sigma\|_{\infty} + \|\nabla\sigma_2\|_{\infty} < +\infty$ then for every $z_0 \in \mathbb{R}^k$ and $T > 0$ there is a unique solution $(Z_t)_{t \in [0, T]}$ to (3.19) such that $Z_0 = z_0$.*

Proof. By Theorem 3.10 we have at most one solution. We now construct a solution on an arbitrary finite interval $[0, T]$, arguing as in the proof of Theorem 2.15. We define $\Lambda \subseteq [0, T]$ as the set of all s such that there is a solution $(Z_t)_{t \in [0, s]}$ to (3.19). By Proposition 3.6, Λ is an open subset of $[0, T]$ and contains 0. By the a priori estimates of Theorem 3.9, Λ is a closed subset of $[0, T]$. Therefore $\Lambda = [0, T]$. $\square$

3.9. MILSTEIN SCHEME AND LOCAL EXISTENCE

In this section we prove the local existence result of Proposition 3.6, under the assumption that σ, σ_2 are of class C^1 and uniformly Lipschitz. To construct a solution to (3.10), we set $t_i := \frac{i}{n}$, $i \geq 0$, and for a given $y_0 \in \mathbb{R}^k$

$$y_{t_{i+1}} = y_{t_i} + \sigma(y_{t_i}) \mathbb{X}_{t_i t_{i+1}}^1 + \sigma_2(y_{t_i}) \mathbb{X}_{t_i t_{i+1}}^2, \quad i \geq 0.$$

We set $D := \max \{1, \|\nabla \sigma\|_\infty, \|\nabla \sigma_2\|_\infty\}$, $\mathbb{T} := \{t_i : t_i \leq T\}$ and

$$\begin{aligned}\delta y_{t_i t_j} &:= y_{t_j} - y_{t_i}, \\ \|\delta y\|_\alpha^\mathbb{T} &:= \sup_{0 < i < j \leq nT} \frac{|y_{t_j} - y_{t_i}|}{|t_j - t_i|^\alpha}, \\ A_{t_i t_j} &:= \sigma(y_{t_i}) \mathbb{X}_{t_i t_j}^1 + \sigma_2(y_{t_i}) \mathbb{X}_{t_i t_j}^2.\end{aligned}$$

The main technical estimate is the following

LEMMA 3.13. *Let $M > 0$. There exists $T_{M,D,\alpha} > 0$ such that, for all $T \in (0, T_{M,D,\alpha})$ and $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha,d}$ such that $\|\mathbb{X}^1\|_\alpha + \|\mathbb{X}^2\|_{2\alpha} \leq M$, we have*

$$\begin{aligned}\|\delta y\|_\alpha^\mathbb{T} &\leq 5M(|\sigma(y_0)| + |\sigma_2(y_0)|), \\ \|\delta y - A\|_{3\alpha}^\mathbb{T} &\lesssim_{M,D,\alpha} (|\sigma(y_0)| + |\sigma_2(y_0)|).\end{aligned}$$

Proof. Let us set $R_{t_i t_j} := \delta y_{t_i t_j} - A_{t_i t_j}$. By the definitions, $R_{t_i t_{i+1}} = 0$. Then we can apply the discrete Sewing bound (Theorem 1.18) to R on $\mathbb{T} := \{\frac{i}{n} : i \leq nT\}$ and we obtain

$$\|R\|_{3\alpha}^\mathbb{T} \leq C_{3\alpha} \|\delta R\|_{3\alpha}^\mathbb{T}, \quad C_{3\alpha} = 2^{3\alpha} \sum_{n \geq 1} \frac{1}{n^{3\alpha}}.$$

Now, analogously to (3.26), since $\delta R = -\delta A$,

$$\delta R_{t_i t_j t_k} = \underbrace{-(\sigma(y_{t_j}) - \sigma(y_{t_i}) - \sigma_2(y_{t_i}) \mathbb{X}_{t_i t_j}^1) \mathbb{X}_{t_j t_k}^1}_{B_{ij}} - \underbrace{(\sigma_2(y_{t_i}) - \sigma_2(y_{t_j})) \mathbb{X}_{t_j t_k}^2}_{C_{ij}},$$

so that

$$\|\delta R\|_{3\alpha}^\mathbb{T} \leq M(\|B\|_{2\alpha}^\mathbb{T} + \|C\|_\alpha^\mathbb{T}).$$

We set

$$H_{t_i t_j} := \delta y_{t_i t_j} - \sigma(y_{t_i}) \mathbb{X}_{t_i t_j}^1,$$

and by (3.23) we obtain

$$\begin{aligned}B_{t_i t_j} &= \sigma(y_{t_j}) - \sigma(y_{t_i}) - \sigma_2(y_{t_i}) \mathbb{X}_{t_i t_j}^1 = \\ &= \underbrace{\int_0^1 (\sigma_2(y_{t_i} + u \delta y_{t_i t_j}) - \sigma_2(y_{t_i})) \mathbb{X}_{t_i t_j}^1 du}_{E_{ij}} + \underbrace{\int_0^1 \nabla \sigma(y_{t_i} + u \delta y_{t_i t_j}) du H_{t_i t_j}}_{F_{ij}} \\ &\quad - \underbrace{\int_0^1 \nabla \sigma(y_{t_i} + u \delta y_{t_i t_j}) (\sigma(y_{t_i} + u \delta y_{t_i t_j}) - \sigma(y_{t_i})) \mathbb{X}_{t_i t_j}^1 du}_{G_{ij}}.\end{aligned}$$

First

$$\|E\|_{2\alpha}^\mathbb{T} \leq \|\nabla \sigma_2\|_\infty \|\delta y\|_\alpha^\mathbb{T} \|\mathbb{X}^1\|_\alpha \leq DM \|\delta y\|_\alpha^\mathbb{T}.$$

Similarly

$$\|G\|_{2\alpha}^\mathbb{T} \leq \|\nabla \sigma\|_\infty^2 \|\delta y\|_\alpha^\mathbb{T} \|\mathbb{X}^1\|_\alpha \leq D^2 M \|\delta y\|_\alpha^\mathbb{T}.$$

By the definition of $R_{t_i t_j}$

$$\begin{aligned} |H_{t_i t_j}| &\leq |R_{t_i t_j}| + |\sigma_2(y_{t_i}) \mathbb{X}_{t_i t_j}^2| \\ &\leq [T^\alpha \|R\|_{3\alpha}^\mathbb{T} + (|\sigma_2(y_0)| + T^\alpha \|\nabla \sigma_2\|_\infty \|\delta y\|_\alpha^\mathbb{T}) \|\mathbb{X}^2\|_{2\alpha}] |t_j - t_i|^{2\alpha} \\ &\leq (T^\alpha \|R\|_{3\alpha}^\mathbb{T} + M |\sigma_2(y_0)| + T^\alpha D M \|\delta y\|_\alpha^\mathbb{T}) |t_j - t_i|^{2\alpha}. \end{aligned}$$

Therefore

$$\begin{aligned} \|F\|_{2\alpha}^\mathbb{T} &\leq D \|H\|_{2\alpha}^\mathbb{T} \\ &\leq D (T^\alpha \|R\|_{3\alpha}^\mathbb{T} + M |\sigma_2(y_0)| + T^\alpha D M \|\delta y\|_\alpha^\mathbb{T}). \end{aligned}$$

Finally

$$\begin{aligned} \|B\|_{2\alpha}^\mathbb{T} &\leq \|E\|_{2\alpha}^\mathbb{T} + \|F\|_{2\alpha}^\mathbb{T} + \|G\|_{2\alpha}^\mathbb{T} \\ &\leq D [M |\sigma_2(y_0)| + T^\alpha \|R\|_{3\alpha}^\mathbb{T} + D M (2 + T^\alpha) \|\delta y\|_\alpha^\mathbb{T}]. \end{aligned}$$

Analogously

$$\|C\|_{2\alpha}^\mathbb{T} \leq D \|\delta y\|_\alpha^\mathbb{T}.$$

Therefore

$$\|R\|_{3\alpha}^\mathbb{T} \leq C_{3\alpha} D M (M |\sigma_2(y_0)| + T^\alpha \|R\|_{3\alpha}^\mathbb{T} + [1 + D M (2 + T^\alpha)] \|\delta y\|_\alpha^\mathbb{T}).$$

If $T^\alpha C_{3\alpha} D M \leq \frac{1}{2}$ then

$$\|R\|_{3\alpha}^\mathbb{T} \leq 2C_{3\alpha} D M (M |\sigma_2(y_0)| + [1 + D M (2 + T^\alpha)] \|\delta y\|_\alpha^\mathbb{T}). \quad (3.81)$$

We set

$$L(y) := 2C_{3\alpha} D M (M |\sigma_2(y_0)| + [1 + D M (2 + T^\alpha)] \|\delta y\|_\alpha^\mathbb{T})$$

Now we obtain by (3.81)

$$\begin{aligned} \|\delta y\|_\alpha^\mathbb{T} &\leq \|R\|_\alpha^\mathbb{T} + \|A\|_\alpha^\mathbb{T} \\ &\leq T^{2\alpha} L(y) + (|\sigma(y_0)| + |\sigma_2(y_0)| + 2 D T^\alpha \|\delta y\|_\alpha^\mathbb{T}) M. \end{aligned}$$

If we assume also that $2 D M T^\alpha \leq \frac{1}{2}$, we obtain

$$\|\delta y\|_\alpha \leq 2 T^{2\alpha} L(y) + 2 M (|\sigma(y_0)| + |\sigma_2(y_0)|).$$

By the definition of $L(y)$, if furthermore $2C_{3\alpha} D M [1 + D M (2 + T^\alpha)] T^{2\alpha} \leq \frac{1}{2}$, we obtain finally

$$\begin{aligned} \|\delta y\|_\alpha^\mathbb{T} &\leq 5 M (|\sigma(y_0)| + |\sigma_2(y_0)|), \\ L(y) &\leq 12 C_{3\alpha} D M^2 [1 + D M (2 + T^\alpha)] (|\sigma(y_0)| + |\sigma_2(y_0)|) =: K, \end{aligned}$$

and by (3.81)

$$\|\delta y - A\|_{3\alpha}^\mathbb{T} \leq K.$$

The proof is complete. $\square$

Proof of Proposition 3.6. Arguing as in Theorem 2.16 we obtain the result of local existence for equation (3.19) of Proposition 3.6. $\square$

CHAPTER 4

STOCHASTIC DIFFERENTIAL EQUATIONS

In this chapter we connect the *rough difference equations (RDE)* discussed in the previous chapter, see (3.18), with the classical *stochastic differential equations (SDE)* $dY_t = \sigma(Y_t) dB_t$ driven by a Brownian motion B . Indeed, both RDE and SDE are ways to make sense of the ill-posed differential equation $\dot{Y}_t = \sigma(Y_t) \dot{B}_t$.

We fix a time horizon $T > 0$ and two dimensions $k, d \in \mathbb{N}$. Let $B = (B_t)_{t \in [0, T]}$ be a d -dimensional Brownian motion (with continuous paths) relative to a filtration $(\mathcal{F}_t)_{t \in [0, T]}$, defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$. We fix a sufficiently regular function $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ and we consider a solution $Y = (Y_t)_{t \in [0, T]}$ of the SDE

$$dY_t = \sigma(Y_t) dB_t \quad \text{i.e.} \quad Y_t = Y_0 + \int_0^t \sigma(Y_s) dB_s, \quad t \geq 0, \quad (4.1)$$

where the stochastic integral is in the Ito sense. We always fix a version of Y with continuous paths (we recall that the Ito integral is a continuous local martingale).

We want to show that Y solves a *rough difference equation* driven by the *rough path* $\mathbb{B} = (\mathbb{B}^1, \mathbb{B}^2)$ (see Definition 3.2) defined by

$$\mathbb{B}_{st}^1 := B_t - B_s, \quad \mathbb{B}_{st}^2 := \int_s^t (B_r - B_s) \otimes dB_r, \quad 0 \leq s \leq t \leq T, \quad (4.2)$$

where the stochastic integral is in the Ito sense. More explicitly, for $i, j \in \{1, \dots, d\}$

$$(\mathbb{B}_{st}^1)^i := B_t^i - B_s^i, \quad (\mathbb{B}_{st}^2)^{ij} := \int_s^t (B_r^i - B_s^i) dB_r^j, \quad (4.3)$$

where we write $B_t = (B_t^1, \dots, B_t^d)$, so that $\mathbb{B}^1: [0, T]_{\leq}^2 \rightarrow \mathbb{R}^d$ and $\mathbb{B}^2: [0, T]_{\leq}^2 \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$.

Our first main result is that $(\mathbb{B}^1, \mathbb{B}^2)$ is indeed a rough path over B .

THEOREM 4.1. (ITO ROUGH PATH) *Almost surely, $\mathbb{B} := (\mathbb{B}^1, \mathbb{B}^2)$ is an α -rough path over B (see Definition 3.2) for any $\alpha \in]\frac{1}{3}, \frac{1}{2}[$, namely it satisfies a.s.*

$$\begin{aligned} \delta \mathbb{B}_{sut}^2 &:= \mathbb{B}_{st}^2 - \mathbb{B}_{su}^2 - \mathbb{B}_{ut}^2 = \mathbb{B}_{su}^1 \otimes \mathbb{B}_{ut}^1, \\ |\mathbb{B}_{st}^1| &\lesssim |t - s|^\alpha, \quad |\mathbb{B}_{st}^2| \lesssim |t - s|^{2\alpha}. \end{aligned} \quad (4.4)$$

Our second main result is that, under suitable assumptions, the solution Y of the SDE (4.1) solves the RDE (3.18) driven by the Ito rough path $\mathbb{X} = \mathbb{B}$.

THEOREM 4.2. (SDE & RDE) *If $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is of class C^2 , then almost surely any solution $Y = (Y_t)_{t \in [0, T]}$ of the SDE (4.1) is also a solution of the RDE*

$$\delta Y_{st} = \sigma(Y_s) \mathbb{B}_{st}^1 + \sigma_2(Y_s) \mathbb{B}_{st}^2 + o(t - s), \quad 0 \leq s \leq t \leq T. \quad (4.5)$$

(We recall that $\sigma_2(\cdot) := \nabla \sigma(\cdot) \sigma(\cdot)$ is defined in (3.5).)

If $\sigma(\cdot)$ is of class C^3 and, furthermore, $\sigma(\cdot)$ and $\sigma_2(\cdot)$ are globally Lipschitz, i.e. $\|\nabla \sigma\|_\infty + \|\nabla \sigma_2\|_\infty < \infty$, then almost surely both the SDE (4.1) and the RDE (4.5) admit a unique solution $Y = (Y_t)_{t \in [0, T]}$ and these solutions coincide.

The key tool we exploit in this chapter is a *local expansion of stochastic integrals*, see Theorem 4.3 in the next Section 4.1. The proofs of Theorems 4.1 and 4.2 are direct consequences of this result, see Section 4.2.

In Sections 4.3 and 5.1 we discuss useful generalizations of the SDE (4.1), where we add a drift and we allow for stochastic integration in the Stratonovich sense, which leads to generalized versions of Theorems 4.1 and 4.2.

In Section 5.2 we present the celebrated result by Wong-Zakai on the limit of solutions of the SDE (4.1) with a regularized Brownian motion (via convolution).

Finally, Section 4.4 is devoted to a far-reaching generalization of Kolmogorov's continuity criterion, which leads to the proof of Theorem 4.3 in Section 4.5.

4.1. LOCAL EXPANSION OF STOCHASTIC INTEGRALS

We recall that $B = (B_t)_{t \in [0, T]}$ is a d -dimensional Brownian motion. Let $h = (h_t)_{t \in [0, T]}$ be a stochastic process with values in $\mathbb{R}^k \otimes (\mathbb{R}^d)^*$. We assume that h is adapted and has continuous paths, in particular $\int_0^T |h_s|^2 ds < \infty$, hence the Itô integral

$$I_t := I_0 + \int_0^t h_r dB_r \quad (4.6)$$

is well-defined as a local martingale. It is a classical result that the stochastic process $I = (I_t)_{t \in [0, T]}$ admits a version with continuous paths, which we always fix.

We now state the main technical result of this chapter, proved in Section 4.5 below, which connects the regularity of h to the regularity of I .

THEOREM 4.3. (LOCAL EXPANSION OF STOCHASTIC INTEGRALS) *Let $h: [0, T] \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ be an adapted process with a.s. continuous paths. Fix any $\alpha \in]0, \frac{1}{2}[$ and recall $(\mathbb{B}^1, \mathbb{B}^2)$ from (4.2).*

1. *Almost surely I is of class $\mathcal{C}^\alpha$, i.e.*

$$|I_t - I_s| \lesssim (t - s)^\alpha, \quad \forall 0 \leq s \leq t \leq T. \quad (4.7)$$

(We recall that the implicit constant in the relation $\lesssim$ is random.)

2. *Assume that, almost surely, $|\delta h_{sr}| \lesssim (r - s)^\beta$ for some $\beta \in]0, 1]$ (i.e. h is of class $\mathcal{C}^\beta$). Then, almost surely,*

$$|\delta I_{st} - h_s \mathbb{B}_{st}^1| = \left| \int_s^t \delta h_{sr} dB_r \right| \lesssim (t - s)^{\alpha + \beta}, \quad \forall 0 \leq s \leq t \leq T. \quad (4.8)$$

3. *Assume that, almost surely, $|\delta h_{sr} - h_s^1 \mathbb{B}_{sr}^1| \lesssim (r - s)^{\eta + \alpha}$, for some adapted process $h^1 = (h_t^1)_{t \in [0, T]}$ of class $\mathcal{C}^\eta$ with $\eta \in]0, 1]$. Then, almost surely,*

$$\begin{aligned} |\delta I_{st} - h_s \mathbb{B}_{st}^1 - h_s^1 \mathbb{B}_{st}^2| &= \left| \int_s^t (\delta h_{sr} - h_s^1 \mathbb{B}_{sr}^1) dB_r \right| \\ &\lesssim (t - s)^{\eta + 2\alpha}, \quad \forall 0 \leq s \leq t \leq T. \end{aligned} \quad (4.9)$$

The proof of Theorem 4.3 is postponed to Section 4.5.

4.2. BROWNIAN ROUGH PATH AND SDES

In this section we exploit Theorem 4.3 to prove Theorems 4.1 and 4.2.

Proof. (OF THEOREM 4.1) We need to verify that $\mathbb{B} = (\mathbb{B}^1, \mathbb{B}^2)$ satisfies the Chen relation (3.13) and the analytic bounds (3.14).

The Chen relation $\delta \mathbb{B}_{sut}^2 = \mathbb{B}_{su}^1 \otimes \mathbb{B}_{ut}^1$ for $0 \leq s \leq u \leq t \leq T$ holds by (4.3):

$$\begin{aligned} \delta(\mathbb{B}^2)_{sut}^{ij} &= (\mathbb{B}^2)_{st}^{ij} - (\mathbb{B}^2)_{su}^{ij} - (\mathbb{B}^2)_{ut}^{ij} \\ &= \int_s^t (B_r^i - B_s^i) dB_r^j - \int_s^u (B_r^i - B_s^i) dB_r^j - \int_u^t (B_r^i - B_u^i) dB_r^j \\ &= \int_u^t (B_u^i - B_s^i) dB_r^j = (B_u^i - B_s^i) \int_u^t 1 dB_r^j = (B_u^i - B_s^i)(B_t^j - B_u^j), \end{aligned}$$

by the properties of the Itô integral and the fact that the times $s \leq u \leq t$ are ordered.

The first analytic bound $|\mathbb{B}_{st}^1| \lesssim |t - s|^\alpha$ for $\alpha \in]0, \frac{1}{2}[$ is a well-known almost sure property of Brownian motion, which also follows from Theorem 4.3, applying (4.7) with $h \equiv 1$. Finally, the second analytic bound $|\mathbb{B}_{st}^2| \lesssim |t - s|^{2\alpha}$ is also a consequence of Theorem 4.3: it suffices to apply (4.8) with $h_s := B_s$ and $\beta = \alpha$. $\square$

Proof. (THEOREM 4.2) We first prove the second part of the statement.

- When σ is globally Lipschitz ($\|\nabla \sigma\|_\infty < +\infty$), it is a classical result that for the SDE (4.1) there is existence of strong solutions and pathwise uniqueness.
- When σ is of class C^3 , by Theorem 3.10 there is uniqueness of solutions for the RDE (3.19), and if both σ and σ_2 are globally Lipschitz ($\|\nabla \sigma\|_\infty < +\infty$ and $\|\nabla \sigma_2\|_\infty < +\infty$) there is also existence of solutions, by Theorem 3.12.

Therefore we only need to prove the first part of the statement: we assume that σ is of class C^2 and we show that given a solution $Y = (Y_t)_{t \in [0, T]}$ of the SDE (4.1), almost surely Y is also a solution to the RDE (4.5).

Since Y is solution to (4.1), recalling (4.2) we can write

$$\begin{aligned} \delta Y_{st} - \sigma(Y_s) \mathbb{B}_{st}^1 - \sigma_2(Y_s) \mathbb{B}_{st}^2 &= \int_s^t (\sigma(Y_r) - \sigma(Y_s)) dB_r - \sigma_2(Y_s) \int_s^t (B_r - B_s) dB_r \\ &= \int_s^t (\delta \sigma(Y)_{sr} - \sigma_2(Y_s) \mathbb{B}_{sr}^1) dB_r. \end{aligned}$$

Let us fix $\alpha \in]0, \frac{1}{2}[$. We prove below that, almost surely,

$$|\delta \sigma(Y)_{st} - \sigma_2(Y_s) \mathbb{B}_{st}^1| \lesssim (t - s)^{2\alpha}, \quad \forall 0 \leq s \leq t \leq T. \quad (4.10)$$

This means that the assumptions of part 3 of Theorem 4.3 are satisfied by $h_r = \sigma(Y_r)$ and $h_r^1 = \sigma_2(Y_r)$ with $\eta = \alpha$: applying (4.9) we then obtain, almost surely,

$$|\delta Y_{st} - \sigma(Y_s) \mathbb{B}_{st}^1 - \sigma_2(Y_s) \mathbb{B}_{st}^2| \lesssim (t - s)^{3\alpha}.$$

If we fix $\alpha > \frac{1}{3}$, this shows that Y is indeed a solution of the RDE (4.5).

It remains to prove (4.10). By Itô's formula and (4.1) we have, for $0 \leq s < t \leq T$,

$$\begin{aligned}
 \sigma(Y_t) &= \sigma(Y_s) + \int_s^t \sum_{a=1}^k \partial_a \sigma(Y_r) dY_r^a + \frac{1}{2} \int_s^t \sum_{a,b=1}^k \partial_{ab} \sigma(Y_r) d\langle Y^a, Y^b \rangle_r \\
 &= \sigma(Y_s) + \int_s^t \sum_{a=1}^k \partial_a \sigma(Y_r) \sum_{c=1}^d \sigma_c^a(Y_r) dB_r^c + \\
 &\quad + \underbrace{\int_s^t \frac{1}{2} \sum_{a,b=1}^k \sum_{c=1}^d \partial_{ab} \sigma(Y_r) \sigma_c^a(Y_r) \sigma_c^b(Y_r) dr}_{p(Y_r)} \\
 &= \sigma(Y_s) + \int_s^t \sigma_2(Y_r) dB_r + \int_s^t p(Y_r) dr,
 \end{aligned} \tag{4.11}$$

therefore

$$\delta \sigma(Y)_{st} - \sigma_2(Y_s) \mathbb{B}_{st}^1 = \int_s^t (\sigma_2(Y_r) - \sigma_2(Y_s)) dB_r + \int_s^t p(Y_r) dr.$$

To prove (4.10), we show that both integrals in the RHS are $O((t-s)^{2\alpha})$.

- Since σ is of class C^2 and Y has continuous paths, the random function $r \mapsto p(Y_r)$ is continuous, hence bounded for $r \in [0, T]$, therefore

$$\left| \int_s^t p(Y_r) dr \right| \lesssim (t-s) \lesssim (t-s)^{2\alpha}, \quad \forall 0 \leq s \leq t \leq T.$$

- Almost surely Y is of class $\mathcal{C}^\alpha$, thanks to (4.7) from Theorem 4.3 and (4.1). Since σ_2 is of class C^1 , hence locally Lipschitz, $r \mapsto \sigma_2(Y_r)$ is of class $\mathcal{C}^\alpha$ too. Applying (4.8) from Theorem 4.3 with $\beta = \alpha$ we then obtain, almost surely,

$$\left| \int_s^t (\sigma_2(Y_r) - \sigma_2(Y_s)) dB_r \right| \lesssim (t-s)^{2\alpha}, \quad \forall 0 \leq s \leq t \leq T.$$

This completes the proof. □

4.3. SDE WITH A DRIFT

It is natural to consider the SDE (4.1) with a non-zero drift term:

$$\begin{aligned}
 dY_t &= b(Y_t) dt + \sigma(Y_t) dB_t \quad \text{i.e.} \\
 Y_t &= Y_0 + \int_0^t b(Y_s) ds + \int_0^t \sigma(Y_s) dB_s, \quad t \geq 0,
 \end{aligned} \tag{4.12}$$

where $b: \mathbb{R}^k \rightarrow \mathbb{R}^k$ and $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ are given and we recall that $B = (B_t)_{t \geq 0}$ is a d -dimensional Brownian motion. We can generalize Theorem 4.2 as follows.

THEOREM 4.4. (SDE & RDE WITH DRIFT) *If $\sigma(\cdot)$ is of class C^2 and $b(\cdot)$ is continuous, then almost surely any solution $Y = (Y_t)_{t \in [0, T]}$ of the SDE (4.12) is also a solution of the RDE*

$$\delta Y_{st} = b(Y_s)(t - s) + \sigma(Y_s) \mathbb{B}_{st}^1 + \sigma_2(Y_s) \mathbb{B}_{st}^2 + o(t - s), \quad 0 \leq s \leq t \leq T. \quad (4.13)$$

If $\sigma(\cdot)$ and $b(\cdot)$ are of class C^3 and, furthermore, $\sigma(\cdot)$, $\sigma_2(\cdot)$ and $b(\cdot)$ are globally Lipschitz, i.e. $\|\nabla \sigma\|_\infty + \|\nabla \sigma_2\|_\infty + \|\nabla b\|_\infty < \infty$, then almost surely the SDE (4.12) and the RDE (4.13) have the same unique solution $Y = (Y_t)_{t \in [0, T]}$.

Proof. We cast the generalized SDE (4.12) in the “usual framework” by adding a component to the driving noise B , i.e. we define $\tilde{B}: [0, T] \rightarrow \mathbb{R}^d \times \mathbb{R}$ by

$$\tilde{B}_t := (B_t, t) = (B_t^1, \dots, B_t^d, t), \quad t \in [0, T],$$

and accordingly we define $\tilde{\sigma}: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^{d+1})^*$ by

$$\tilde{\sigma}(\cdot) \tilde{b} := \sigma(\cdot) b + b(\cdot) t \quad \text{for} \quad \tilde{b} = (b, t) \in \mathbb{R}^d \times \mathbb{R},$$

that is $\tilde{\sigma}(\cdot)_j^i = \sigma(\cdot)_j^i \mathbb{1}_{\{j \leq d\}} + b(\cdot)^i \mathbb{1}_{\{j = d+1\}}$. We can then rewrite the SDE (4.12) as

$$dY_t = \tilde{\sigma}(Y_t) d\tilde{B}_t \quad \text{i.e.} \quad Y_t = Y_0 + \int_0^t \tilde{\sigma}(Y_s) d\tilde{B}_s, \quad t \geq 0. \quad (4.14)$$

We next extend the Ito rough path $\mathbb{B} = (\mathbb{B}^1, \mathbb{B}^2)$ from (4.2), defining

$$\tilde{\mathbb{B}}_{st}^1 := \tilde{B}_t - \tilde{B}_s = \begin{pmatrix} \mathbb{B}_{st}^1 \\ t - s \end{pmatrix}, \quad (4.15)$$

$$\tilde{\mathbb{B}}_{st}^2 := \int_s^t (\tilde{B}_r - \tilde{B}_s) \otimes d\tilde{B}_r = \begin{pmatrix} \mathbb{B}_{st}^2 & \int_s^t (B_r - B_s) dr \\ \int_s^t (r - s) dB_r & \int_s^t (r - s) dr = \frac{(t - s)^2}{2} \end{pmatrix}. \quad (4.16)$$

One can show that $\tilde{\mathbb{B}} = (\tilde{\mathbb{B}}^1, \tilde{\mathbb{B}}^2)$ is a rough path over $\tilde{B}$, following closely the proof of Theorem 4.1. Indeed, if we fix $\alpha \in]0, \frac{1}{2}[$, we have almost surely $B \in C^\alpha$, hence

$$\left| \int_s^t (B_r - B_s) dr \right| \lesssim (t - s)^{\alpha+1}, \quad \left| \int_s^t (r - s) dB_r \right| \lesssim (t - s)^{\alpha+1}. \quad (4.17)$$

We can now write the RDE which generalizes (4.5):

$$\delta Y_{st} = \tilde{\sigma}(Y_s) \tilde{\mathbb{B}}_{st}^1 + \tilde{\sigma}_2(Y_s) \tilde{\mathbb{B}}_{st}^2 + o(t - s). \quad (4.18)$$

Interestingly, plugging the definitions of $\tilde{\mathbb{B}}$ and $\tilde{\sigma}$ into (4.18) we do not obtain (4.13), because the components of $\tilde{\mathbb{B}}_{st}^2$ other than $\mathbb{B}_{st}^2$ are missing in (4.13), see (4.16). The point is that these components can be absorbed in the reminder $o(t - s)$, see (4.17), hence the RDE (4.18) and (4.13) are fully equivalent.

To complete the proof, we are left with comparing the SDE (4.14) with the RDE (4.18). This can be done following the very same arguments as in the proof of Theorem 4.2. The details are left to the reader. $\square$

Remark 4.5. The strategy of adding the drift term as an additional component of the driving noise, as in the proof of Theorem 4.4, suffers from a technical limitation, namely *we are forced to use the same regularity exponent α for all components*, due to Definition 3.2 of rough paths. This prevents us from exploiting the additional regularity of the drift term: for instance, in the second part of Theorem 4.4, the assumption that $b(\cdot)$ is of class C^3 could be removed, because the “driving noise” t is smooth and the classical theory of ordinary differential equations applies.

A natural solution would be to generalize Definition 3.2, allowing rough paths to have a different regularity exponent for each component. The key results can be generalized to this setting, but for simplicity we refrain from pursuing this path.

4.4. A REFINED KOLMOGOROV CRITERION

In this section we prepare the ground for the proof of Lemmas 4.10 and 4.11 in Section 4.5 below, which are the main technical tools in the proof of Theorem 4.3. We suppose without loss of generality that $T = 1$, namely our processes are defined on the interval $[0, 1]$. Define the set $\mathbb{D}$ of dyadic points in $[0, 1]$ by

$$\mathbb{D} := \bigcup_{k \geq 0} D_k, \quad \text{where} \quad D_k := \left\{ d_i^k := \frac{i}{2^k} \right\}_{0 \leq i \leq 2^k}. \quad (4.19)$$

Given $d, \tilde{d} \in \mathbb{D}$, we write $d \rightarrow \tilde{d}$ if and only if $\tilde{d}$ is consecutive to d in some layer D_k of $\mathbb{D}$, that is $d = d_i^k$ and $\tilde{d} = d_{i+1}^k$, for some $k \geq 0$ and $0 \leq i \leq 2^k - 1$.

Remarkably, in order to prove relation (4.35), *it is enough to have a suitable control on $R_{d, \tilde{d}}$ for consecutive points $d \rightarrow \tilde{d}$* (together with a global control on δR), as the next result shows. This turns out to be at the heart of the Kolmogorov continuity criterion, but we stress that it is a deterministic statement.

THEOREM 4.6. (KOLMOGOROV CRITERION: DETERMINISTIC PART) *Given a function $A: \mathbb{D}_{<}^2 \rightarrow \mathbb{R}$, for $0 < \rho < \gamma$ we define the constants*

$$Q_\gamma := \sup_{d, \tilde{d} \in \mathbb{D}: d \rightarrow \tilde{d}} \frac{|A_{d, \tilde{d}}|}{|\tilde{d} - d|^\gamma}, \quad (4.20)$$

$$K_{\rho, \gamma} := \sup_{\substack{0 \leq s < u < t \leq 1 \\ s, u, t \in \mathbb{D}}} \frac{|\delta A_{s, u, t}|}{\min(u - s, t - u)^\rho |t - s|^{\gamma - \rho}}. \quad (4.21)$$

Then there is a constant $C_{\rho, \gamma} < \infty$ such that

$$|A_{st}| \leq C_{\rho, \gamma} (Q_\gamma + K_{\rho, \gamma}) |t - s|^\gamma, \quad \forall (s, t) \in \mathbb{D}_{<}^2. \quad (4.22)$$

A key tool for Theorem 4.6 is the next result, proved at this end of this section, which ensures the existence of suitable *short paths* in $\mathbb{D}$.

LEMMA 4.7. (DYADIC PATHS) *For any $s, t \in \mathbb{D}$ with $s < t$, there are integers $n, m \geq 1$ and a path of $(m + n + 1)$ points in $\mathbb{D}$ which leads from s to t , labelled as follows:*

$$s = s_m < \dots < s_1 < s_0 = t_0 < t_1 < \dots < t_n = t, \quad (4.23)$$

with the property that for all $i \in \{0, \dots, m - 1\}$ and $j \in \{0, \dots, n - 1\}$

$$s_{i+1} \rightarrow s_i, \quad t_j \rightarrow t_{j+1}; \quad |s_i - s_{i+1}| < \frac{|t - s|}{2^i}, \quad |t_{j+1} - t_j| < \frac{|t - s|}{2^j}. \quad (4.24)$$

Proof of Theorem 4.6. Fix $s, t \in \mathbb{D}$ with $s < t$. We use Lemma 4.7 with the same notation. By the definition of δA , we write

$$A_{st} = A_{st_0} + A_{t_0t} + \delta A_{s,t_0,t}.$$

In the case $m \geq 2$, we can develop A_{st_0} as follows (recall that $s = s_m$ and $s_0 = t_0$):

$$A_{st_0} = \sum_{i=0}^{m-1} A_{s_{i+1}s_i} + \sum_{i=0}^{m-2} \delta A_{s,s_{i+1},s_i}.$$

Similarly, when $n \geq 2$, we develop

$$A_{t_0t} = \sum_{j=0}^{n-1} A_{t_j t_{j+1}} + \sum_{j=0}^{n-2} \delta A_{t_j, t_{j+1}, t},$$

so that

$$\begin{aligned} A_{st} &= \underbrace{\sum_{i=0}^{m-1} A_{s_{i+1}s_i} + \sum_{j=0}^{n-1} A_{t_j t_{j+1}}}_{\Xi_1} + \\ &\quad + \underbrace{\delta A_{s,t_0,t} + \sum_{i=0}^{m-2} \delta A_{s,s_{i+1},s_i} + \sum_{j=0}^{n-2} \delta A_{t_j, t_{j+1}, t}}_{\Xi_2}. \end{aligned} \quad (4.25)$$

By the definition of Q_γ , for any $d \rightarrow \tilde{d}$ we can bound

$$|A_{d\tilde{d}}| \leq Q_\gamma |\tilde{d} - d|^\gamma.$$

By Lemma 4.7, this bound applies to any couple (s_{i+1}, s_i) and (t_j, t_{j+1}) . Then we can estimate Ξ_1 in (4.25) as follows, exploiting the bounds in (4.24):

$$\begin{aligned} &Q_\gamma \left\{ \sum_{i=0}^{m-1} |s_i - s_{i+1}|^\gamma + \sum_{j=0}^{n-1} |t_{j+1} - t_j|^\gamma \right\} \leq \\ &\leq Q_\gamma \left\{ \sum_{i=0}^{\infty} (2^{-i})^\gamma + \sum_{j=0}^{\infty} (2^{-j})^\gamma \right\} |t - s|^\gamma = \\ &= Q_\gamma \left\{ \frac{2}{1 - 2^{-\gamma}} \right\} |t - s|^\gamma, \end{aligned}$$

which agrees with (4.22). On the other hand, thanks to (4.21) and (4.24),

$$|\delta A_{s,s_{i+1},s_i}| \leq K_{\rho,\gamma} \left(\frac{|t-s|}{2^i} \right)^\rho |t-s|^{\gamma-\rho} = K_{\rho,\gamma} 2^{-i\rho} |t-s|^\gamma$$

and similarly for $\delta A_{t_j, t_{j+1}, t}$, so that the term Ξ_2 can be bounded above by

$$K_{\rho,\gamma} |t-s|^\gamma \left(1 + \sum_{i=0}^{m-2} 2^{-i\rho} + \sum_{j=0}^{n-2} 2^{-j\rho} \right) \leq K_{\rho,\gamma} |t-s|^\gamma \left(1 + \frac{2}{1 - 2^{-\rho}} \right).$$

This completes the proof of (4.22). $\square$

As a simple consequence of Theorem 4.6, we show that suitable moment conditions ensure the finiteness of the constant Q_γ in (4.20), as in the classical Kolmogorov criterion.

PROPOSITION 4.8. (KOLMOGOROV CRITERION: PROBABILISTIC PART) *Let $A = (A_{st})_{(s,t) \in \mathbb{D}_{<}^2}$ be a stochastic process which satisfies the following bound, for some γ_0 , $p, c \in (0, \infty)$:*

$$\mathbb{E}[|A_{st}|^p] \leq c|t - s|^{p\gamma_0}, \quad \forall (s, t) \in \mathbb{D}_{<}^2.$$

Then, for any value of γ such that

$$\gamma < \gamma_0 - \frac{1}{p}, \quad (4.26)$$

the random variable $Q_\gamma = Q_\gamma(A)$ defined in (4.20) is in L^p :

$$\mathbb{E}[|Q_\gamma|^p] \leq \frac{c}{1 - 2^{1-p(\gamma_0-\gamma)}} < \infty. \quad (4.27)$$

In particular, a.s. $Q_\gamma < +\infty$.

Proof. By definition of Q_γ in (4.20), bounding the supremum with a sum we can write

$$|Q_\gamma|^p \leq \sum_{d, \tilde{d} \in \mathbb{D}: d \rightarrow \tilde{d}} \left(\frac{|A_{d, \tilde{d}}|}{|\tilde{d} - d|^\gamma} \right)^p = \sum_{k \geq 0} \sum_{i=0}^{2^k-1} \frac{|A_{d_i^k d_{i+1}^k}|^p}{|d_{i+1}^k - d_i^k|^{p\gamma}}.$$

Let us write $\gamma = \gamma_0 - \frac{1+\epsilon}{p}$, for some $\epsilon > 0$. Since $d_{i+1}^k - d_i^k = \frac{1}{2^k}$ we have

$$\begin{aligned} \mathbb{E}[|Q_\gamma|^p] &\leq \sum_{k \geq 0} \sum_{i=0}^{2^k-1} c |d_{i+1}^k - d_i^k|^{p(\gamma_0-\gamma)} \\ &\leq \sum_{k \geq 0} \sum_{i=0}^{2^k-1} \frac{c}{2^{(1+\epsilon)k}} = \sum_{k \geq 0} \frac{c}{2^{\epsilon k}} = \frac{c}{1 - 2^{-\epsilon}} < \infty. \end{aligned}$$

The proof is complete. $\square$

Remark 4.9. Given a stochastic process $(X_t)_{t \in \mathbb{D}}$ defined on dyadic times, if we apply Theorem 4.6 and Proposition 4.8 to $(A_{st} := \delta X_{st} = X_t - X_s)_{(s,t) \in \mathbb{D}_{<}^2}$ we obtain the classical Kolmogorov continuity criterion. Note that in this case $K_{\rho, \gamma} = 0$ because $\delta A = 0$.

Proof of Lemma 4.7. We refer to Figure 4.1 for a graphical representation. Given $s, t \in \mathbb{D}$ with $s < t$, since $0 < t - s \leq 1$, we can define $\ell \geq 1$ as the unique integer such that

$$\frac{1}{2^\ell} < t - s \leq \frac{1}{2^{\ell-1}}. \quad (4.28)$$

We now take the smallest $k \in \{0, \dots, 2^\ell - 1\}$ for which $d_k^\ell > s$ and define

$$s_0 := t_0 := d_k^\ell.$$

The definition of k guarantees that $d_k^\ell < t$, because if $d_k^\ell \geq t$ then $\frac{k}{2^\ell} - s \geq t - s > \frac{1}{2^\ell}$ and this would violate the minimality of k .

Note that $0 < d_k^\ell - s \leq d_k^\ell - d_{k-1}^\ell = \frac{1}{2^\ell}$ and $0 < t - d_k^\ell < t - s$, by (4.28), therefore

$$0 < s_0 - s < \frac{1}{2^{\ell-1}}, \quad 0 < t - t_0 < \frac{1}{2^{\ell-1}}. \quad (4.29)$$

Since both $s_0 - s \in \mathbb{D}$ and $t - t_0 \in \mathbb{D}$, for suitable integers $m \geq 1$ and $n \geq 1$ we have

$$s_0 - s = \frac{1}{2^{q_1}} + \frac{1}{2^{q_2}} + \dots + \frac{1}{2^{q_m}}, \quad t - t_0 = \frac{1}{2^{r_1}} + \frac{1}{2^{r_2}} + \dots + \frac{1}{2^{r_n}},$$

where $q_m > q_{m-1} > \dots > q_1 \geq \ell$ and $r_n > \dots > r_1 \geq \ell$. We can thus write

$$\begin{aligned} s &= s_0 - \frac{1}{2^{q_1}} - \frac{1}{2^{q_2}} - \dots - \frac{1}{2^{q_m}}, \\ t &= t_0 + \frac{1}{2^{r_1}} + \frac{1}{2^{r_2}} + \dots + \frac{1}{2^{r_n}}. \end{aligned}$$

We can finally define

$$\begin{aligned} s_i &:= s_0 - \frac{1}{2^{q_1}} - \frac{1}{2^{q_2}} - \dots - \frac{1}{2^{q_i}} \quad \text{for } i = 1, \dots, m, \\ t_j &:= t_0 + \frac{1}{2^{r_1}} + \frac{1}{2^{r_2}} + \dots + \frac{1}{2^{r_j}} \quad \text{for } j = 1, \dots, n. \end{aligned}$$




Figure 4.1. An instance of Lemma 4.7 with $s = \frac{5}{32}$ and $t = \frac{11}{16}$. Note that $\ell = 1$ (because $\frac{1}{2^1} < |t - s| = \frac{17}{32} \leq \frac{1}{2^0}$, cf. (4.28)) and $s_0 = t_0 = \frac{1}{2}$. The points $t_1, \dots, t_n$ are built iteratively: first take the largest $\frac{1}{2^{r_1}}$ (i.e. the smallest r_1) such that $t_1 := t_0 + \frac{1}{2^{r_1}} \leq t$; if $t_1 < t$, then take the largest $\frac{1}{2^{r_2}}$ such that $t_2 := t_1 + \frac{1}{2^{r_2}} \leq t$; and so on, until $t_n = t$. Similarly for $s_1, \dots, s_m$.

Since q_i and r_j are strictly increasing integers with $q_1 \geq \ell$ and $r_1 \geq \ell$, we have the bounds $q_i \geq \ell + (i - 1)$ and $r_j \geq \ell + (j - 1)$, for all $i \in \{0, \dots, m - 1\}$ and $j \in \{0, \dots, n - 1\}$, hence

$$\begin{aligned} |s_i - s_{i+1}| &= \frac{1}{2^{q_{i+1}}} \leq \frac{1}{2^i} \frac{1}{2^\ell} < \frac{|t - s|}{2^i}, \\ |t_{j+1} - t_j| &= \frac{1}{2^{r_{j+1}}} \leq \frac{1}{2^j} \frac{1}{2^\ell} < \frac{|t - s|}{2^j}. \end{aligned}$$

having used (4.28). This proves the bounds in (4.24).

We note that, for any integer $r \geq \ell$, we have the inclusion $D_\ell \subseteq D_r$. Then, given any $x \in D_\ell$, we have that $x \in D_r$, hence $x \rightarrow x + 2^{-r}$. Since $t_0 = d_k^\ell \in D_\ell$ and $r_1 \geq \ell$, this shows that $t_0 \rightarrow t_1 = t_0 + 2^{-r_1}$. Proceeding inductively, we have $t_j \rightarrow t_{j+1} = t_j + 2^{-r_{j+1}}$. A similar argument applies to the points s_i and completes the proof of (4.24). $\square$

4.5. PROOF OF THEOREM 4.3

In this section we prove the three assertions of Theorem 4.3.

Proof of the first assertion of Theorem 4.3. We want to prove that for any $\alpha \in (0, \frac{1}{2})$, a.s. I is α -Hölder continuous, namely there is an a.s. finite random constant C such that

$$|\delta I_{st}| \leq C |t - s|^\alpha, \quad \forall 0 \leq s \leq t \leq T. \quad (4.30)$$

First observation: if the claim holds under the stronger assumption $|h| \leq c$ almost surely, for some deterministic $c < \infty$, then we can deduce the general result by localization. Indeed, if we only assume that $\sup_{[0,T]} |h| < \infty$ a.s., we can define for $n \in \mathbb{N}$ the stopping times

$$\tau_n := \inf \{t \in [0, T] : |h_t| > n\}.$$

Let us define

$$h_s^{(n)} := h_{s \wedge \tau_n}, \quad I_t^{(n)} := \int_0^t h_s^{(n)} dB_s.$$

Note that $\sup_{[0,T]} |h^{(n)}| \leq n$ by the definition of τ_n . Then

$$|\delta I_{st}^{(n)}| \leq C^{(n)} |t - s|^\alpha, \quad \forall 0 \leq s < t \leq T, \quad (4.31)$$

for a suitable a.s. finite random constant $C^{(n)}$. Let us define the events

$$A_n := \{\tau_n = \infty\} = \left\{ \sup_{[0,T]} |h| \leq n \right\}$$

and note that $h = h^{(n)}$ on A_n . By the locality property of the stochastic integral, $I = I^{(n)}$ a.s. on A_n ^{4.1}.

Note that $A := \bigcup_{n \in \mathbb{N}} A_n = \{\sup_{[0,T]} |h| < \infty\}$, hence $\mathbb{P}(A) = 1$. If we define $C := C^{(n)}$ on $A_n \setminus A_{n-1}$ (with $A_0 := \emptyset$) and $C := \infty$ on A^c , we have $C < \infty$ a.s. and relation (4.7) holds.

Second observation: if relation (4.30) holds for all s, t in a (deterministic) dense subset $\mathbb{D} \subseteq [0, T]$, then it holds for all $s, t \in [0, T]$, because δI_{st} is a continuous function of (s, t) .

In conclusion, the proof is reduced to showing (4.30) only for $s, t \in \mathbb{D}$, under the assumption that $\sup_{[0,T]} |h| \leq c < \infty$ almost surely. Suppose that this is the case and set $A_{st} := \delta I_{st}$, $0 \leq s \leq t \leq T$. Here $\delta A = 0$ and therefore the constant $K_{\rho, \gamma}$ in (4.21) is equal to zero for any $0 < \rho < \gamma$. It remains to estimate Q_α using Proposition 4.8.

By the BDG inequality of Proposition 4.12, for any $p \geq 2$

$$\mathbb{E}[|\delta I_{st}|^p] \leq c_p \mathbb{E} \left[\left(\int_s^t h_u^2 du \right)^{\frac{p}{2}} \right] \leq C_p |t - s|^{\frac{p}{2}}.$$

Then Proposition 4.8 applies with $\gamma_0 = \frac{1}{2}$ and any $\alpha = \gamma_0 - \frac{1}{p} \in (0, \frac{1}{2})$ for p sufficiently large. By Theorem 4.6, we obtain (4.30) and the proof is complete. $\square$

For $0 \leq s \leq t \leq T$ we define the (random) continuous function

$$R_{st} := I_t - I_s - h_s(B_t - B_s) = \int_s^t \delta h_{sr} dB_r. \quad (4.32)$$

We recall that a.s. $B \in \mathcal{C}^\alpha$ for every $\alpha < \frac{1}{2}$.

Proof of the second assertion of Theorem 4.3. Let $\alpha < \frac{1}{2}$. We want to show that, if a.s. $h \in \mathcal{C}^\beta$, for some $\beta \in (0, 1]$, then there is an a.s. finite random constant C such that

$$|R_{st}| \leq C |t - s|^{\alpha + \beta}, \quad \forall 0 \leq s \leq t \leq T. \quad (4.33)$$

^{4.1} We mean that $I^{(n)}$ and I are indistinguishable on A_n : for a.e. $\omega \in A_n$ one has $I_t^{(n)}(\omega) = I_t(\omega)$ for all $t \in [0, 1]$ (we recall that we always fix continuous versions of the stochastic integrals).

First observation: if the claim holds under the stronger assumption $\|\delta h\|_\beta \leq c$ almost surely, for some deterministic $c < \infty$, then we can deduce the general result by localization. Indeed, if we only assume that $\|\delta h\|_\beta < \infty$ a.s., we can define for $n \in \mathbb{N}$ the stopping times

$$\tau_n := \inf \{t \in [0, 1] : \|\delta h\|_{\beta, [0, t]} > n\},$$

where $\|\delta h\|_{\alpha, [0, t]}$ is the Hölder semi-norm of h restricted to $[0, t]$ (equivalently, the Hölder semi-norm of $s \mapsto h_{s \wedge t}$ on the whole interval $s \in [0, 1]$). Let us define

$$h_s^{(n)} := h_{s \wedge \tau_n}, \quad I_t^{(n)} := \int_0^t h_s^{(n)} dB_s, \quad R_{st}^{(n)} := I_t^{(n)} - I_s^{(n)} - h_s^{(n)}(B_t - B_s).$$

Note that $\|\delta h^{(n)}\|_\beta \leq n$, by definition of τ_n . (Indeed, $\|\delta h\|_{\beta, [0, t]} \leq n$ for all $t < \tau_n$, which means that $|h(r) - h(s)| \leq n|r - s|^\beta$ for all $r, s \in [0, \tau_n]$; then, by continuity, $|h(r) - h(s)| \leq n|r - s|^\beta$ for all $r, s \in [0, \tau_n]$, which means that $\|\delta h\|_{\beta, [0, \tau_n]} = \|\delta h^{(n)}\|_\beta \leq n$). Then

$$|R_{st}^{(n)}| \leq C^{(n)}|t - s|^{\alpha+\beta}, \quad \forall 0 \leq s < t \leq T, \quad (4.34)$$

for a suitable a.s. finite random constant $C^{(n)}$. Let us define the events

$$A_n := \{\tau_n = \infty\} = \{\|\delta h\|_\alpha \leq n\}$$

and note that $h = h^{(n)}$ on A_n . By the locality property of the stochastic integral, $I = I^{(n)}$ a.s. on A_n ,^{4.2} hence also $R = R^{(n)}$ a.s. on A_n . Redefining $C^{(n)} = \infty$ on the exceptional set $\{R = R^{(n)}\}^c$, we get by (4.34)

$$\text{on the event } A_n: \quad |R_{st}| \leq C^{(n)}|t - s|^{\alpha+\beta}, \quad \forall 0 \leq s < t \leq T.$$

Note that $A := \bigcup_{n \in \mathbb{N}} A_n = \{\|\delta h\|_\beta < \infty\}$, hence $\mathbb{P}(A) = 1$. If we define $C := C^{(n)}$ on $A_n \setminus A_{n-1}$ (with $A_0 := \emptyset$) and $C := \infty$ on A^c , we have $C < \infty$ a.s. and relation (4.8) holds.

Second observation: if relation (4.33) holds for all s, t in a (deterministic) dense subset $\mathbb{D} \subseteq [0, 1]$, then it holds for all $s, t \in [0, 1]$, because R_{st} is a continuous function of (s, t) .

In conclusion, the proof is reduced to showing (4.33) only for $s, t \in \mathbb{D}$, under the assumption that $\|\delta h\|_\beta \leq c < \infty$. This technical result is formulated in the separate Lemma 4.10. $\square$

LEMMA 4.10. *Let $0 < \alpha < \frac{1}{2}$ and $0 < \beta \leq 1$. Assume that $\mathbb{E}[\|\delta h\|_\beta^p] < \infty$ for all $p > 0$. Then there is an a.s. finite random constant C such that*

$$|R_{st}| \leq C|t - s|^{\alpha+\beta}, \quad \forall s, t \in \mathbb{D} \quad \text{with } s \leq t. \quad (4.35)$$

Equivalently, a.s. $R \in C_2^{\alpha+\beta}$.

Proof. We apply Theorem 4.6 to the (random) function $A(s, t) = R_{st}$, with $\gamma = \alpha + \beta$ and $\rho = \alpha \wedge \beta$. Then relation (4.22) yields (4.35). It remains to show that a.s. $Q_{\alpha+\beta} < \infty$ and $K_{\rho, \alpha+\beta} < \infty$.

^{4.2.} We mean that $I^{(n)}$ and I are indistinguishable on A_n : for a.e. $\omega \in A_n$ one has $I_t^{(n)}(\omega) = I_t(\omega)$ for all $t \in [0, 1]$ (we recall that we always fix continuous versions of the stochastic integrals).

We recall that R_{st} is defined in (4.32). In particular, for $s < u < t$

$$\delta R_{sut} = R_{st} - R_{su} - R_{ut} = (h_u - h_s)(B_t - B_u).$$

Then by (4.21), a.s.

$$K_{\rho, \alpha + \beta}(R) \leq \|\delta h\|_\beta \|\delta B\|_\alpha \sup_{0 \leq s < u < t \leq 1} \frac{|u - s|^\beta |t - u|^\alpha}{\min(u - s, t - u)^{\alpha \wedge \beta} |t - s|^{\alpha \vee \beta}}.$$

By our assumption that $\|\delta h\|_\beta \in L^p$ and by the fact that B is a Brownian motion, it follows that $\|\delta h\|_\beta \|\delta B\|_\alpha < \infty$ a.s., hence it only remains to show that the constant defined by the supremum is bounded above by 1. However, this constant equals

$$\sup_{a, b > 0, a + b = 1} \frac{a^\alpha b^\beta}{(a \wedge b)^{\alpha \wedge \beta}} = \sup_{a, b > 0, a + b = 1} \left(\frac{ab}{(a \wedge b)} \right)^{\alpha \wedge \beta} a^{\alpha - \alpha \wedge \beta} b^{\beta - \alpha \wedge \beta} \leq 1.$$

We want now to estimate $Q_{\alpha + \beta}(R)$. We note that, for fixed $s < t$, we have a.s. $R_{st} = \int_s^t (h_u - h_s) dB_u$. By the Burkholder-Davies-Gundy inequality, see Proposition 4.12, for any $p > 2$ there is a universal constant c_p such that

$$\begin{aligned} \mathbb{E}[|R_{st}|^p] &\leq c_p \mathbb{E} \left[\left(\int_s^t (h_u - h_s)^2 du \right)^{\frac{p}{2}} \right] \\ &\leq c_p \mathbb{E} \left[\|\delta h\|_\beta^p \left(\int_s^t (u - s)^{2\beta} du \right)^{\frac{p}{2}} \right] \\ &\leq c_p \mathbb{E}[\|\delta h\|_\beta^p] (t - s)^{p(\beta + \frac{1}{2})}. \end{aligned}$$

By Proposition 4.8, we have $Q_\gamma < \infty$ a.s. for any $\gamma < \beta + \frac{1}{2} - \frac{1}{p}$. Plugging $\gamma = \alpha + \beta$ we get $\alpha < \frac{1}{2} - \frac{1}{p}$, which is satisfied for p large enough, since $\alpha < \frac{1}{2}$. $\square$

Next, we suppose that there exists another adapted process $h^1 = (h_t^1)_{t \in [0, T]}$ with values in $\mathbb{R}^k \otimes (\mathbb{R}^d)^*$ such that a.s.

$$|\delta h_{st} - h_s^1 \mathbb{B}_{st}^1| \lesssim |t - s|^{\eta + \alpha}.$$

Then we define

$$\begin{aligned} \hat{R}_{st} &:= R_{st} - h_s^1 \mathbb{B}_{st}^2 = \delta I_{st} - h_s \mathbb{B}_{st}^1 - h_s^1 \mathbb{B}_{st}^2 \\ &= \int_s^t (\delta h_{sr} - h_s^1 \mathbb{B}_{sr}^1) dB_r, \end{aligned} \tag{4.36}$$

where $\mathbb{B}^2$ is defined in (4.2). Then the third assertion of Theorem 4.3 follows with the same localisation argument as for the second one and from the following

LEMMA 4.11. *Assume that $\mathbb{E}[\|\delta h^1\|_\eta^p + \|\delta h - h^1 \mathbb{B}^1\|_{\eta + \alpha}^p] < \infty$, for some $\alpha \in (0, \frac{1}{2})$ and for all $p > 0$. Then there is an a.s. finite random constant C such that*

$$|\hat{R}_{st}| \leq C |t - s|^{\eta + 2\alpha}, \quad \forall s, t \in \mathbb{D} \quad \text{with } s \leq t. \tag{4.37}$$

Equivalently, a.s. $\hat{R} \in C_2^{\eta + 2\alpha}$.

Proof. We apply Theorem 4.6 to the (random) function $A(s, t) = R_{st}$, with $\gamma = \alpha + \beta$ and $\rho = \alpha \wedge \eta$. Then

$$\delta \hat{R}_{sut} = (\delta h_{su} - h_s^1 \mathbb{B}_{su}^1) \mathbb{B}_{ut}^1 + \delta h_{su}^1 \mathbb{B}_{ut}^2.$$

Now

$$\begin{aligned} K_{\rho, \eta+2\alpha}(\hat{R}) &\leq \|\delta h - h^1 \mathbb{B}^1\|_{\eta+\alpha} \|\mathbb{B}^1\|_{\alpha} \sup_{0 \leq s < u < t \leq 1} \frac{|u-s|^{\eta+\alpha} |t-u|^{\alpha}}{\min(u-s, t-u)^{\rho} |t-s|^{\eta+2\alpha-\rho}} \\ &\quad + \|\delta h^1\|_{\eta} \|\mathbb{B}^2\|_{2\alpha} \sup_{0 \leq s < u < t \leq 1} \frac{|u-s|^{\eta} |t-u|^{2\alpha}}{\min(u-s, t-u)^{\rho} |t-s|^{\eta+2\alpha-\rho}}. \end{aligned}$$

We note that the first supremum is equal to

$$\sup_{a, b > 0, a+b=1} \frac{a^{\eta+\alpha} b^{\alpha}}{(a \wedge b)^{\rho}} \leq \sup_{a, b > 0, a+b=1} \left(\frac{ab}{a \wedge b} \right)^{\alpha \wedge \eta} a^{\alpha \vee \eta} b^{\alpha - \alpha \wedge \eta} \leq 1,$$

while the second supremum is equal to

$$\sup_{a, b > 0, a+b=1} \frac{a^{\eta} b^{2\alpha}}{(a \wedge b)^{\rho}} \leq \sup_{a, b > 0, a+b=1} \left(\frac{ab}{a \wedge b} \right)^{\alpha \wedge \eta} a^{\eta - \alpha \wedge \eta} b^{2\alpha - \alpha \wedge \eta} \leq 1.$$

Now by (4.36)

$$\begin{aligned} \mathbb{E}[|\hat{R}_{st}|^p] &\leq \mathbb{E}\left[\left(\int_s^t (\delta h_{su} - h_s^1 \mathbb{B}_{su}^1)^2 du\right)^{\frac{p}{2}}\right] \\ &\leq c_p \mathbb{E}\left[\|\delta h - h^1 \mathbb{B}^1\|_{\eta+\alpha}^p \left(\int_s^t (u-s)^{2(\eta+\alpha)} du\right)^{\frac{p}{2}}\right] \\ &\leq c_p \mathbb{E}[\|\delta h - h^1 \mathbb{B}^1\|_{\eta+\alpha}^p] (t-s)^{p(\eta+\alpha+\frac{1}{2})}. \end{aligned}$$

By Proposition 4.8, we have $Q_{\gamma} < \infty$ a.s. for any $\gamma < \eta + \alpha + \frac{1}{2} - \frac{1}{p}$. Plugging $\gamma = \eta + 2\alpha$ we get $\alpha < \frac{1}{2} - \frac{1}{p}$, which is satisfied for p large enough, since $\alpha < \frac{1}{2}$. $\square$

Finally, we give a proof of (half of) Burkholder-Davies-Gundy inequality for $p \geq 2$.

PROPOSITION 4.12. *For all $p \geq 2$ there is a constant $c_p < \infty$ such that for all $0 \leq s \leq t \leq T$*

$$\mathbb{E}\left[\left|\int_s^t y_u dB_u\right|^p\right] \leq c_p \mathbb{E}\left[\left(\int_s^t y_u^2 du\right)^{\frac{p}{2}}\right]$$

for any progressively measurable process such that $\mathbb{P}$ -a.s. $\int_0^1 y_u^2 du < \infty$.

Proof. To simplify the notation we set $s = 0$ and $m_t := \int_0^t y_u dB_u$.

First we make the additional assumptions that $\mathbb{E}[\int_0^1 y_u^2 du] < \infty$ and m is bounded by some deterministic constant. By the Itô formula applied to m_t , we get

$$d|m_t|^p = p|m_t|^{p-1} \text{sgn}(m_t) y_t dB_t + \frac{p(p-1)}{2} |m_t|^{p-2} y_t^2 dt.$$

In general $(\int_0^t |m_u|^{p-1} \text{sgn}(m_u) y_u dB_u)_t$ is a local martingale, but under our additional assumptions it is a true martingale with zero expectation, because $\mathbb{E}[\int_0^1 |m_u|^{2(p-1)} y_u^2 du] < \infty$ (recall that m is bounded). Consequently

$$\mathbb{E}[|m_t|^p] = \frac{p(p-1)}{2} \mathbb{E}\left[\int_0^t |m_u|^{p-2} y_u^2 du\right].$$

If we set $|\bar{m}_t| := \sup_{u \leq t} |m_u|$, we obtain by Hölder

$$\begin{aligned} \mathbb{E}[|m_t|^p] &\leq \frac{p(p-1)}{2} \mathbb{E} \left[|\bar{m}_t|^{p-2} \int_0^t y_u^2 du \right] \\ &\leq \frac{p(p-1)}{2} \mathbb{E}[|\bar{m}_t|^p]^{1-\frac{2}{p}} \mathbb{E} \left[\left(\int_0^t y_u^2 du \right)^{\frac{p}{2}} \right]^{\frac{2}{p}}. \end{aligned} \quad (4.38)$$

Since $(|m_t|)_{t \geq 0}$ is submartingale bounded in L^p with continuous trajectories, by Doob L^p inequality we have: $\mathbb{E}[|\bar{m}_t|^p] \leq (\frac{p}{p-1})^p \mathbb{E}[|m_t|^p]$. Plugging the above in (4.38) we conclude:

$$\mathbb{E} \left[\left| \int_0^t y_u dB_u \right|^p \right] \leq c_p \mathbb{E} \left[\left(\int_0^t y_u^2 du \right)^{\frac{p}{2}} \right].$$

As far as the general case is concerned, let us define

$$\tau^n = \inf \{ t \geq 0 : |m_t| > n \} \wedge \inf \left\{ t \geq 0 : \int_0^t y_u^2 du > n \right\}$$

Note that $\mathbb{P}$ -a.s. τ^n is a non decreasing sequence of stopping times, with $\tau^n = \infty$ for n large enough. We denote $y_t^n := y \mathbb{1}_{[0, \tau^n]}(t)$ and $m_t^n := \int_0^t y_u^n dB_u$. By construction, y^n and m^n satisfy our additional assumptions. Since $m_t^n = m_{t \wedge \tau^n}$ a.s., we have

$$\begin{aligned} \mathbb{E} \left[\left| \int_0^{t \wedge \tau^n} y_u dB_u \right|^p \right] &\leq c_p \mathbb{E} \left[\left(\int_0^t y_u^2 \mathbb{1}_{[0, \tau^n]}(u) du \right)^{\frac{p}{2}} \right] \\ &\leq c_p \mathbb{E} \left[\left(\int_0^t y_u^2 du \right)^{\frac{p}{2}} \right]. \end{aligned}$$

Finally we notice that by Fatou's Lemma

$$\begin{aligned} \mathbb{E} \left[\left| \int_0^t y_u dB_u \right|^p \right] &= \mathbb{E} \left[\liminf_{n \rightarrow \infty} \left| \int_0^{t \wedge \tau^n} y_u dB_u \right|^p \right] \\ &\leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\left| \int_0^{t \wedge \tau^n} y_u dB_u \right|^p \right] \\ &\leq c_p \mathbb{E} \left[\left(\int_0^t y_u^2 du \right)^{\frac{p}{2}} \right]. \end{aligned}$$

The proof is complete. □

CHAPTER 5

WONG-ZAKAI

5.1. ITÔ VERSUS STRATONOVICH

We recall that $B = (B_t)_{t \in [0, T]}$ is a Brownian motion in $\mathbb{R}^d$. Given the Itô rough path $\mathbb{B} = (\mathbb{B}^1, \mathbb{B}^2)$ over B constructed in Theorem 4.2, see (4.2), we can define a new rough path $\bar{\mathbb{B}} = (\bar{\mathbb{B}}^1, \bar{\mathbb{B}}^2)$ over B , called the *Stratonovich rough path*, given by

$$\bar{\mathbb{B}}_{st}^1 := \mathbb{B}_{st}^1, \quad \bar{\mathbb{B}}_{st}^2 := \mathbb{B}_{st}^2 + \frac{t-s}{2} \text{Id}_{\mathbb{R}^d}, \quad \forall 0 \leq s \leq t \leq T, \quad (5.1)$$

that is

$$(\bar{\mathbb{B}}_{st}^2)^{ij} := (\mathbb{B}_{st}^2)^{ij} + \frac{t-s}{2} \mathbb{1}_{\{i=j\}} = \begin{cases} \frac{(B_t^i - B_s^i)^2}{2} & \text{if } i = j, \\ \int_s^t (B_r^i - B_s^i) dB_r^j & \text{if } i \neq j. \end{cases} \quad (5.2)$$

The fact that $\bar{\mathbb{B}}$ is an α -rough path over B , for any $\alpha \in]\frac{1}{3}, \frac{1}{2}[$, is a consequence of Theorem 4.1 (note that $\bar{\mathbb{B}}_{st}^2 = \mathbb{B}_{st}^2 + \delta f_{st}$ with $f_t = \frac{t}{2} \text{Id}_{\mathbb{R}^d}$, hence $\delta \bar{\mathbb{B}}^2 = \delta \mathbb{B}^2$).

Remark 5.1. (STRATONOVICH INTEGRAL) If $X, Y: [0, T] \rightarrow \mathbb{R}$ are continuous semimartingales, the Stratonovich integral of X with respect to Y is defined by

$$\int_0^t X_s \circ dY_s := \int_0^t X_s dY_s + \frac{1}{2} \langle X, Y \rangle_t, \quad t \in [0, T], \quad (5.3)$$

where $\int_0^t X_s dY_s$ is the Itô integral and $\langle \cdot, \cdot \rangle$ is the quadratic covariation. For Brownian motion B on $\mathbb{R}^d$ we have $\langle B^i, B^j \rangle_t = t \mathbb{1}_{\{i=j\}}$, hence recalling (4.2) we see that

$$\bar{\mathbb{B}}_{st}^2 := \int_s^t \bar{\mathbb{B}}_{sr}^1 \otimes \odot dB_r, \quad 0 \leq s \leq t \leq T. \quad (5.4)$$

This explains why we call $\bar{\mathbb{B}} = (\bar{\mathbb{B}}^1, \bar{\mathbb{B}}^2)$ the Stratonovich rough path.

Let us consider now the Stratonovich version of the SDE (4.12):

$$dY_t = b(Y_t) dt + \sigma(Y_t) \circ dB_t, \quad \text{i.e.}$$

$$Y_t = Y_0 + \int_0^t b(Y_s) ds + \int_0^t \sigma(Y_s) \circ dB_s, \quad t \geq 0, \quad (5.5)$$

where $b: \mathbb{R}^k \rightarrow \mathbb{R}^k$ and $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ are given. This equation can be recast in the Itô form by the conversion rule (5.3): since the martingale part of $(\sigma(Y_t))_{t \geq 0}$ is $(\int_0^t \sigma_2(Y_s) dB_s)_{t \geq 0}$ by the Itô formula, see (4.11), we obtain

$$Y_t = Y_0 + \int_0^t \left(b(Y_s) + \frac{1}{2} \text{Tr}_{\mathbb{R}^d} [\sigma_2(Y_s)] \right) ds + \int_0^t \sigma(Y_s) dB_s, \quad t \geq 0.$$

This is precisely the SDE (4.12) with a different drift $\hat{b}(\cdot) := b(\cdot) + \frac{1}{2} \text{Tr}_{\mathbb{R}^d}[\sigma_2(\cdot)]$.

As an immediate corollary of Theorem 4.4, we obtain the following result.

THEOREM 5.2. (STRATONOVICH SDE & RDE) *If $\sigma(\cdot)$ is of class C^2 and $b(\cdot)$ is continuous, then almost surely any solution $Y = (Y_t)_{t \in [0, T]}$ of the Stratonovich SDE (5.5) is also a solution of the following RDE, for $0 \leq s \leq t \leq T$:*

$$\begin{aligned} \delta Y_{st} &= b(Y_s)(t-s) + \sigma(Y_s) \bar{\mathbb{B}}_{st}^1 + \sigma_2(Y_s) \bar{\mathbb{B}}_{st}^2 + o(t-s) \\ &= \left(b(Y_s) + \frac{1}{2} \text{Tr}_{\mathbb{R}^d}[\sigma_2(Y_s)] \right) (t-s) + \sigma(Y_s) \mathbb{B}_{st}^1 + \sigma_2(Y_s) \mathbb{B}_{st}^2 + o(t-s). \end{aligned} \quad (5.6)$$

If $\sigma(\cdot)$, $\sigma_2(\cdot)$, $b(\cdot)$ are of class C^3 and, furthermore, $\sigma(\cdot)$, $\sigma_2(\cdot)$, $b(\cdot)$ are globally Lipschitz, i.e. $\|\nabla \sigma\|_\infty + \|\nabla \sigma_2\|_\infty + \|\nabla b\|_\infty < \infty$, then almost surely the SDE (5.5) and the RDE (5.6) have the same unique solution $Y = (Y_t)_{t \in [0, T]}$.

In conclusion, if the coefficients $b(\cdot)$ and $\sigma(\cdot)$ are sufficiently regular, the Itô equation (4.12) can be reinterpreted as the RDE

$$\delta Y_{st} = b(Y_s)(t-s) + \sigma(Y_s) \mathbb{B}_{st}^1 + \sigma_2(Y_s) \mathbb{B}_{st}^2 + o(t-s), \quad 0 \leq s \leq t \leq T,$$

while the Stratonovich equation (5.5) can be reinterpreted as the RDE

$$\delta Y_{st} = b(Y_s)(t-s) + \sigma(Y_s) \bar{\mathbb{B}}_{st}^1 + \sigma_2(Y_s) \bar{\mathbb{B}}_{st}^2 + o(t-s), \quad 0 \leq s \leq t \leq T.$$

In other words, rough paths allow to describe the Itô and the Stratonovich SDEs as *the same equation* where only the second level of the rough path has been changed. This shows that, in a sense, the *relevant noise* for a SDE is not only the Brownian path $(B_t)_{t \geq 0}$, but rather the rough path $\mathbb{B}$ or $\bar{\mathbb{B}}$.

5.2. WONG-ZAKAI

In this section we show the following application of the previous results. We consider a family $(\rho_\varepsilon)_{\varepsilon > 0}$ of compactly supported mollifiers on $\mathbb{R}$, namely $\rho: \mathbb{R} \rightarrow [0, \infty)$ is smooth, compactly supported in $[-1, 1]$, satisfies $\int_{\mathbb{R}} \rho(x) dx = 1$ and we set

$$\rho_\varepsilon(x) := \frac{1}{\varepsilon} \rho\left(\frac{x}{\varepsilon}\right), \quad \varepsilon > 0, \quad x \in \mathbb{R}. \quad (5.7)$$

(We do not assume that ρ is even.) We consider a d -dimensional two-sided Brownian motion $(B_t)_{t \in \mathbb{R}}$, namely a Gaussian centered process with values in $\mathbb{R}^d$ such that

$$B_0 = 0, \quad \mathbb{E}[B_s^i B_t^j] = \mathbb{1}_{(i=j)} \mathbb{1}_{(st \geq 0)} (|s| \wedge |t|),$$

which is equivalent to say that $(B_t)_{t \geq 0}$ and $(B_{-t})_{t \geq 0}$ are two independent d -dimensional Brownian motions.

We consider the following problem: we define a regularization of $(B_t)_{t \geq 0}$ by

$$B_t^\varepsilon := (\rho_\varepsilon * B)_t = \int_{\mathbb{R}} \rho_\varepsilon(u) B_{t-u} du, \quad t \geq 0, \quad (5.8)$$

and we consider the integral equation (3.3) controlled by B^ε , namely

$$Z_t^\varepsilon = Z_0 + \int_0^t \sigma(Z_s^\varepsilon) \dot{B}_s^\varepsilon ds, \quad 0 \leq t \leq T. \quad (5.9)$$

It is easy to check that $(B_t^\varepsilon)_{t \geq 0}$ converges to $(B_t)_{t \geq 0}$ as $\varepsilon \downarrow 0$ uniformly for $t \in [0, T]$ (and even in C^α for any $\alpha < \frac{1}{2}$; see below). Then we want to understand whether $(Z_t^\varepsilon)_{t \geq 0}$ also converges, and especially to which limit.

This question has a very natural answer in the context of rough paths. We define the *canonical rough path* over B^ε (see section 8.7 below for more on this notion):

$$\mathbb{B}_{st}^{\varepsilon,1} := B_t^\varepsilon - B_s^\varepsilon, \quad \mathbb{B}_{st}^{\varepsilon,2} := \int_s^t \mathbb{B}_{su}^{\varepsilon,1} \otimes \dot{B}_u^\varepsilon du, \quad 0 \leq s \leq t. \quad (5.10)$$

Then we can prove the following result.

THEOREM 5.3. (WONG-ZAKAI) *As $\varepsilon \downarrow 0$, $\mathbb{B}^\varepsilon$ converges in probability to the Stratonovich rough path $\bar{\mathbb{B}}$, see (5.1), namely for any $\alpha < \frac{1}{2}$*

$$\|\mathbb{B}^{\varepsilon,1} - \bar{\mathbb{B}}^1\|_\alpha + \|\mathbb{B}^{\varepsilon,2} - \bar{\mathbb{B}}^2\|_{2\alpha} \xrightarrow{\varepsilon \downarrow 0} 0 \quad \text{in probability.} \quad (5.11)$$

The convergence holds almost surely along sequences $(\varepsilon_n)_n$ that converge to 0 exponentially fast.

Moreover let $(Z_t^\varepsilon)_{t \in [0, T]}$ be the solution to the controlled equation

$$Z_t^\varepsilon = Z_0 + \int_0^t \sigma(Z_s^\varepsilon) \dot{B}_s^\varepsilon ds, \quad t \geq 0.$$

Assume that $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^$ is of class C^3 , with $\|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty + \|\nabla^3 \sigma\|_\infty + \|\nabla \sigma_2\|_\infty + \|\nabla^2 \sigma_2\|_\infty < +\infty$. Then, for any $\alpha \in]0, \frac{1}{2}[$, we have $Z^\varepsilon \rightarrow Z$ in probability in $C^\alpha([0, T]; \mathbb{R}^k)$ as $\varepsilon \downarrow 0$, where Z is the unique solution to the Stratonovich SDE*

$$Z_t = Z_0 + \int_0^t \sigma(Z_s) \circ dB_s = Z_0 + \int_0^t \sigma(Z_s) dB_s + \frac{1}{2} \int_0^t \text{Tr}_{\mathbb{R}^d}[\sigma_2(Z_s)] ds.$$

Proof. Fix $\alpha \in (\frac{1}{3}, \frac{1}{2})$. Let $\mathbb{B}^\varepsilon$ be the canonical smooth rough path associated with B^ε as in (3.9). Suppose we have proved that $\mathbb{B}^\varepsilon$ converges to $\bar{\mathbb{B}}$ as in (5.11). By Proposition 3.5, the solution Z^ε to the controlled equation (5.9) is equal to the (unique by Theorem 3.10) solution to the rough finite difference equation (3.19) associated with the α -rough path $\mathbb{B}^\varepsilon$. In the notation (3.51), we have $Z^\varepsilon = \Phi(Z_0, \mathbb{B}^\varepsilon)$, and by Theorem 5.2 we have $Z = \Phi(Z_0, \bar{\mathbb{B}})$. By the continuity result Theorem 3.11 we obtain that $Z^\varepsilon = \Phi(Z_0, \mathbb{B}^\varepsilon) \rightarrow \Phi(Z_0, \bar{\mathbb{B}}) = Z$ a.s. as $\varepsilon \downarrow 0$.

It remains now to prove (5.11). We first observe that by (5.8)

$$\mathbb{B}_{st}^{\varepsilon,1} = \int_{\mathbb{R}} \rho_\varepsilon(u) \delta B_{s-u, t-u} du. \quad (5.12)$$

Let us fix $\alpha < \frac{1}{2}$ and set $\|\delta B\|_\alpha := \|\delta B\|_{\alpha, [-1, T+1]}$, so that $|\delta B_{ab}| \leq \|\delta B\|_\alpha (b-a)^\alpha$ for all $-1 \leq a < b \leq T+1$. Then, uniformly for $\varepsilon \in (0, 1)$ and $0 \leq s < t \leq T$, we can bound

$$|\mathbb{B}_{st}^{\varepsilon,1}| \leq \|\delta B\|_\alpha (t-s)^\alpha. \quad (5.13)$$

We can write similarly

$$\mathbb{B}_{st}^{\varepsilon,1} - \bar{\mathbb{B}}_{st}^1 = \int_{\mathbb{R}} \rho_{\varepsilon}(u) (\delta B_{s-u,t-u} - \delta B_{st}) du, \quad (5.14)$$

hence for any $\alpha' \in]\alpha, \frac{1}{2}[$ we can estimate, by the triangle inequality,

$$|\mathbb{B}_{st}^{\varepsilon,1} - \bar{\mathbb{B}}_{st}^1| \leq 2 \|\delta B\|_{\alpha'} (t-s)^{\alpha'}.$$

At the same time, since $\delta B_{st} - \delta B_{s-u,t-u} = \delta B_{t-u,t} - \delta B_{s-u,s}$, we can also bound

$$|\mathbb{B}_{st}^{\varepsilon,1} - \bar{\mathbb{B}}_{st}^1| \leq 2 \|\delta B\|_{\alpha'} \int_{\mathbb{R}} \rho_{\varepsilon}(u) u^{\alpha'} du \leq 2 \|\delta B\|_{\alpha'} \varepsilon^{\alpha'}$$

because ρ_{ε} is supported in $[-\varepsilon, \varepsilon]$. Overall, we have shown that

$$\begin{aligned} \|\mathbb{B}^{\varepsilon,1} - \bar{\mathbb{B}}^1\|_{\alpha,[0,T]} &\leq 2 \|\delta B\|_{\alpha',[-1,T+1]} \sup_{0 \leq s < t \leq T} \frac{(t-s)^{\alpha'} \wedge \varepsilon^{\alpha'}}{(t-s)^{\alpha}} \\ &= 2 \|\delta B\|_{\alpha',[-1,T+1]} \varepsilon^{\alpha'-\alpha} \xrightarrow{\varepsilon \downarrow 0} 0, \quad \forall \alpha < \alpha' \end{aligned} \quad (5.15)$$

(for the equality, consider separately $t-s > \varepsilon$ and $t-s \leq \varepsilon$). We stress that the previous arguments are *pathwise*. Since $\|\delta B\|_{\alpha',[-1,T+1]} < \infty$ *almost surely* for any $\alpha' < \frac{1}{2}$, it follows that $\|\mathbb{B}^{\varepsilon,1} - \bar{\mathbb{B}}^1\|_{\alpha} \rightarrow 0$ *almost surely* for any $\alpha < \frac{1}{2}$.

To complete the proof of (5.11), it remains to show that $\|\mathbb{B}^{\varepsilon,2} - \bar{\mathbb{B}}^2\|_{2\alpha} \rightarrow 0$ in probability as $\varepsilon \downarrow 0$. We distinguish $(\mathbb{B}^{\varepsilon,2} - \bar{\mathbb{B}}^2)^{ij}$ for $i = j$ (diagonal terms) and for $i \neq j$ (off-diagonal terms, in case $d > 1$). To lighten notation, we fix $i \neq j$ and abbreviate $X = B^i$ and $Y = B^j$, which are *independent Brownian motions*.

Diagonal terms are easy: by (5.10) and integration by parts (since X^{ε} is smooth)

$$(\mathbb{B}^{\varepsilon,2})_{st}^{ii} = \int_s^t (X_u^{\varepsilon} - X_s^{\varepsilon}) \dot{X}_u^{\varepsilon} du = \frac{(X_t^{\varepsilon} - X_s^{\varepsilon})^2}{2}.$$

Similarly $(\bar{\mathbb{B}}^2)_{st}^{ii} = \frac{(X_t - X_s)^2}{2}$ by definition (5.2) of Stratonovich Brownian motion. Since $(\delta X_{st}^{\varepsilon})^2 - (\delta X_{st})^2 = 2 \delta X_{st} \delta(X^{\varepsilon} - X)_{st} + (\delta(X^{\varepsilon} - X)_{st})^2$, by what we already proved on $\|(\mathbb{B}^{\varepsilon,1} - \bar{\mathbb{B}}^1)^i\|_{\alpha} = \|\delta(X^{\varepsilon} - X)\|_{\alpha}$, see (5.15), we have *almost surely*

$$\|(\mathbb{B}^{\varepsilon,2} - \bar{\mathbb{B}}^2)^{ii}\|_{2\alpha} \leq \|\delta X\|_{\alpha} \|\delta(X^{\varepsilon} - X)\|_{\alpha} + \frac{\|\delta(X^{\varepsilon} - X)\|_{\alpha}^2}{2} \xrightarrow{\varepsilon \downarrow 0} 0.$$

We next turn to off-diagonal terms $(\mathbb{B}^{\varepsilon,2} - \bar{\mathbb{B}}^2)^{ij} = L^{\varepsilon} - L$, where we set

$$L_{st} := \int_s^t \delta X_{sw} dY_w, \quad L_{st}^{\varepsilon} := \int_s^t \delta X_{su}^{\varepsilon} dY_u^{\varepsilon} = \int_s^t \delta X_{su}^{\varepsilon} \dot{Y}_u^{\varepsilon} du. \quad (5.16)$$

The core of the proof is the following second moment bound, that we prove below.

PROPOSITION 5.4. (SECOND MOMENT BOUND) *For all $\varepsilon > 0$, $s < t$ we have*

$$\mathbb{E}[(L_{st}^{\varepsilon} - L_{st})^2] \leq 10 (t-s)^2 \min \left\{ 1, \frac{\varepsilon}{t-s} \right\}. \quad (5.17)$$

We derive from (5.17) a bound for moments of order $p \geq 2$ exploiting a key property known as *hypercontractivity*, that we state in the special case which is relevant for us. The proof is given below.

PROPOSITION 5.5. (HYPERCONTRACTIVITY) *Consider the stochastic integral*

$$W := \int_{-\infty}^{\infty} \left(\int_{-\infty}^t g(s, t) dX_s \right) dY_t \quad (5.18)$$

for a deterministic function $g \in L^2(\mathbb{R}^2 \rightarrow \mathbb{R})$. Then the following bound holds:

$$\forall p \in [2, \infty): \quad \mathbb{E}[|W|^p] \leq c_p^2 \mathbb{E}[W^2]^{\frac{p}{2}}, \quad (5.19)$$

with $c_p := \mathbb{E}[|\mathcal{N}(0, 1)|^p] < \infty$.

We can now apply (5.19) to $L_{st}^\varepsilon - L_{st}$, which is of the form (5.18) (see (5.24) below): plugging (5.17) into (5.19) we obtain

$$\mathbb{E}[|L_{st}^\varepsilon - L_{st}|^p] \leq 10 c_p^2 (t-s)^p \min \left\{ 1, \left(\frac{\varepsilon}{t-s} \right)^{\frac{p}{2}} \right\}. \quad (5.20)$$

Since $\min \{1, x\} \leq x^\kappa$ for all $x \geq 0$ and $\kappa \in [0, 1]$, it follows that

$$\forall \kappa \in (0, 1]: \quad \mathbb{E}[|L_{st}^\varepsilon - L_{st}|^p] \leq 10 c_p^2 (t-s)^{p(1-\frac{\kappa}{2})} \varepsilon^{p\frac{\kappa}{2}}. \quad (5.21)$$

We now fix $\alpha < \frac{1}{2}$ and exploit Theorem 4.6 for $A_{st} := L_{st}^\varepsilon - L_{st}$ with $\rho = \alpha$ and $\gamma = 2\alpha$. We need to control the *random constants* $Q_{2\alpha}$ and $K_{\alpha, 2\alpha}$ from (4.20)-(4.21).

- For $Q_{2\alpha}$ we apply Proposition 4.8 with $\gamma_0 = 1 - \frac{\kappa}{2}$: if we take $\kappa > 0$ small and $p \geq 2$ large, so that (4.26) is satisfied, by (5.21) and (4.27) we get

$$\mathbb{E}[Q_{2\alpha}^p] \leq \mathfrak{C} \varepsilon^{p\frac{\kappa}{2}} \quad \text{with} \quad \mathfrak{C} = \mathfrak{C}_{p, \alpha, \kappa} := \frac{10 c_p^2}{1 - 2^{1-p(1-2\alpha-\frac{\kappa}{2})}}. \quad (5.22)$$

This implies that $Q_{2\alpha} \rightarrow 0$ in probability as $\varepsilon \downarrow 0$, and even *almost surely* along sequences $\varepsilon = \varepsilon_n \downarrow 0$ which vanish exponentially fast.

- For $K_{\alpha, 2\alpha}$ we note that, by the Chen relation,

$$\delta A_{sut} = \delta X_{su}^\varepsilon \delta Y_{ut}^\varepsilon - \delta X_{su} \delta Y_{ut} = \delta X_{su}^\varepsilon (\delta Y_{ut}^\varepsilon - \delta Y_{ut}) + \delta Y_{ut} (\delta X_{su}^\varepsilon - \delta X_{su}),$$

therefore by (5.13) and (5.15), if we fix any $\alpha' \in (\alpha, \frac{1}{2})$, we can bound

$$\begin{aligned} K_{\alpha, 2\alpha} &\leq \|\delta X^\varepsilon\|_\alpha \|\delta(Y^\varepsilon - Y)\|_\alpha + \|\delta Y\|_\alpha \|\delta(X^\varepsilon - X)\|_\alpha \\ &\leq 2 (\|\delta X\|_\alpha \|\delta Y\|_{\alpha'} + \|\delta Y\|_\alpha \|\delta X\|_{\alpha'}) \varepsilon^{\alpha' - \alpha}. \end{aligned} \quad (5.23)$$

This shows that $K_{\alpha, 2\alpha} \rightarrow 0$ almost surely as $\varepsilon \downarrow 0$.

We can finally apply (4.22) to conclude that, by (5.22) and (5.23),

$$\|(\mathbb{B}^{\varepsilon, 2} - \bar{\mathbb{B}}^2)^{ij}\|_{2\alpha} = \|L^\varepsilon - L\|_{2\alpha} \leq C_{\alpha, 2\alpha} (Q_{2\alpha} + K_{\alpha, 2\alpha}) \xrightarrow[\varepsilon \downarrow 0]{} 0 \quad \text{in probability.}$$

This completes the proof of (5.11). $\square$

Proof of Proposition 5.4. Recalling that $X^\varepsilon = \rho_\varepsilon * X$ and $Y^\varepsilon = \rho_\varepsilon * Y$, an integration by parts for the stochastic (Wiener) integral yields for $s < t$

$$\begin{aligned}\delta X_{st}^\varepsilon &:= \int_{\mathbb{R}} (\rho_\varepsilon(t-v) - \rho_\varepsilon(s-v)) X_v dv = \int_{\mathbb{R}} \left(\int_{s-v}^{t-v} \rho_\varepsilon(r) dr \right) dX_v, \\ \dot{Y}_\varepsilon(t) &:= \int_{\mathbb{R}} (\rho_\varepsilon)'(t-w) Y_w dw = \int_{\mathbb{R}} \rho_\varepsilon(t-w) dY_w.\end{aligned}$$

Recalling the definition (5.16) of L_{st} and L_{st}^ε , we can write

$$L_{st}^\varepsilon - L_{st} = \iint (g_\varepsilon^{(s,t)}(v, w) - \mathbb{1}_{(s \leq v \leq w \leq t)}) dX_v dY_w, \quad (5.24)$$

where we set

$$\begin{aligned}g_\varepsilon^{(s,t)}(v, w) &:= \int_s^t \rho_\varepsilon(u-w) \left(\int_{s-v}^{u-v} \rho_\varepsilon(r) dr \right) du \\ &= \int \mathbb{1}_{(s \leq r \leq u \leq t)} \rho_\varepsilon(r-v) \rho_\varepsilon(u-w) dr du.\end{aligned}$$

Since $0 \leq g_\varepsilon^{(s,t)}(v, w) \leq 1$ (recall that $\rho_\varepsilon(\cdot)$ is a probability density), it follows that

$$\begin{aligned}\mathbb{E}[(L_{st}^\varepsilon - L_{st})^2] &= \iint (g_\varepsilon^{(s,t)}(v, w) - \mathbb{1}_{(s \leq v \leq w \leq t)})^2 dv dw \\ &\leq \iint |g_\varepsilon^{(s,t)}(v, w) - \mathbb{1}_{(s \leq v \leq w \leq t)}| dv dw.\end{aligned} \quad (5.25)$$

To estimate this integral, we give a probabilistic representation of $g_\varepsilon(v, w)$: denoting by Q_1 and Q_2 two independent random variables with density $\rho(\cdot)$, since $\rho_\varepsilon(\cdot - v)$ and $\rho_\varepsilon(\cdot - w)$ are the densities of $\varepsilon Q_1 + v$ and $\varepsilon Q_2 + w$, we can write

$$g_\varepsilon^{(s,t)}(v, w) = \mathbb{P}(s \leq \varepsilon Q_1 + v \leq \varepsilon Q_2 + w \leq t).$$

Writing $v = s + a(t-s)$ and $w = s + b(t-s)$, for new variables a, b , we note that

$$g_\varepsilon^{(s,t)}(s + a(t-s), s + b(t-s)) = g_\delta^{(0,1)}(a, b) \quad \text{with} \quad \delta := \frac{\varepsilon}{t-s}.$$

A change of variables in the integral (5.25) then yields

$$\mathbb{E}[(L_{st}^\varepsilon - L_{st})^2] \leq (t-s)^2 \iint |g_\delta^{(0,1)}(a, b) - \mathbb{1}_{(0 \leq a \leq b \leq 1)}| da db.$$

Looking at our goal (5.17), it only remains to show that

$$\iint |g_\delta^{(0,1)}(a, b) - \mathbb{1}_{(0 \leq a \leq b \leq 1)}| da db \leq 10 \min\{1, \delta\}. \quad (5.26)$$

We define the subset

$$D := \{(a, b) \in \mathbb{R}^2: 0 \leq a \leq b \leq 1\}$$

so that we can write

$$g_\delta^{(0,1)}(a, b) = \mathbb{P}(0 \leq \delta Q_1 + a \leq \delta Q_2 + b \leq 1) = \mathbb{E}[\mathbb{1}_{D-\delta Q}(a, b)] \quad \text{with } Q := (Q_1, Q_2).$$

We can express the integral in (5.26) as

$$\begin{aligned} \iint |g_\delta^{(0,1)}(a, b) - \mathbb{1}_{(0 \leq a \leq b \leq 1)}| da db &= \mathbb{E} \left[\int_{\mathbb{R}^2} |\mathbb{1}_{D-\delta Q}(z) - \mathbb{1}_D(z)| dz \right] \\ &= \mathbb{E}[|(D - \delta Q) \Delta D|] \end{aligned}$$

where $|\cdot|$ denotes Lebesgue measure in $\mathbb{R}^2$ and $A \Delta B := (A \cap B^c) \cup (A^c \cap B)$ is the symmetric difference between sets. Note that $z \in (D - y) \Delta D$ means that either $z \in D$ but $z + y \in D^c$, or $z \in D^c$ but $z + y \in D$, and in both cases $\text{dist}(z, \partial D) \leq |y|$, where ∂D is the boundary of D . In other terms, for any $y \in \mathbb{R}^2$ we have the inclusion

$$(D - y) \Delta D \subseteq \{z \in \mathbb{R}^2: \text{dist}(z, \partial D) \leq |y|\}.$$

Since ∂D is a triangle with perimeter $2 + \sqrt{2}$, the area of $\{z \in \mathbb{R}^2: \text{dist}(z, \partial D) \leq |y|\}$ is bounded above by $2(2 + \sqrt{2})|y|$, hence

$$\mathbb{E}[|(D - \delta Q) \Delta D|] \leq 2(2 + \sqrt{2}) \mathbb{E}[|\delta Q|] \leq 2(2 + \sqrt{2}) \sqrt{2} \delta,$$

because $|Q| = \sqrt{Q_1^2 + Q_2^2} \leq \sqrt{2}$ (we recall that $\rho(\cdot)$ is supported in $[-1, 1]$, hence $|Q_1|, |Q_2| \leq 1$). Since $2(2 + \sqrt{2})\sqrt{2} \leq 10$, the proof of (5.26) is completed. $\square$

Proof of Proposition 5.5. By (5.18) we can write $W = \int_{-\infty}^{\infty} h(t) dY_t$ where $h(t) = h(X, t) := \int_{-\infty}^t g(s, t) dX_s$ depends only on X . Since X and Y are independent, it follows that W is a Gaussian random variable conditionally on X , as a Wiener integral. Recalling that $c_p := \mathbb{E}[|\mathcal{N}(0, 1)|^p]$, we can thus write

$$\mathbb{E}[|W|^p | X] = c_p \mathbb{E}[W^2 | X]^{\frac{p}{2}},$$

We now denote by $E = C(\mathbb{R}, \mathbb{R})$ the standard path space for X and Y , so that can write $W = f(X, Y)$ for a suitable measurable function $f: E \times E \rightarrow \mathbb{R}$. Denoting by μ the law of X , i.e. the two-sided Wiener measure, Fubini's theorem yields

$$\mathbb{E}[W^2 | X] = \mathbb{E}[f(x, Y)^2]_{x=X} = (\|f(x, \cdot)\|_{L^2(\mu(dy))}^2)_{x=X},$$

hence

$$\mathbb{E}[|W|^p] = c_p \mathbb{E} \left[\mathbb{E}[|W|^2 | X]^{\frac{p}{2}} \right] = c_p (\| \|f(x, \cdot)\|_{L^2(\mu(dy))} \|_{L^p(\mu(dx))})^p.$$

We now apply the *Minkowski integral inequality* (see Remark 5.6 below), which states that for $p \geq 2$ switching the two norms yields an upper bound:

$$\begin{aligned} \mathbb{E}[|W|^p] &\leq c_p (\| \|f(x, \cdot)\|_{L^p(\mu(dx))} \|_{L^2(\mu(dy))})^p \\ &= c_p \left(\left\| \mathbb{E}[|f(X, y)|^p]^{\frac{1}{p}} \right\|_{L^2(\mu(dy))} \right)^p. \end{aligned} \quad (5.27)$$

We finally observe that $f(X, y)$ is a Gaussian random variable, i.e. $W = f(X, Y)$ is Gaussian conditionally on Y (because $W = \int \tilde{h}(s) dX_s$ with $\tilde{h}(t) := \int_s^\infty g(s, t) dY_t$ is a Wiener integral conditionally on Y , by independence of X and Y). It follows that

$$\mathbb{E}[|f(X, y)|^p] = c_p \mathbb{E}[f(X, y)^2]^{\frac{p}{2}} = c_p (\|f(x, y)\|_{L^2(\mu(dx))})^p. \quad (5.28)$$

Plugging (5.28) into (5.27) we obtain (5.19), since

$$\| \|f(x, y)\|_{L^2(\mu(dx))}\|_{L^2(\mu(dy))} = \mathbb{E}[W^2]^{1/2}$$

by Fubini's theorem. $\square$

Remark 5.6. (MINKOWSKI'S INTEGRAL INEQUALITY) Given σ -finite measure spaces (E, μ) and (F, ν) and a measurable function $f: E \times F \rightarrow \mathbb{R}$, Minkowski's integral inequality states that for any $0 < q \leq p \leq \infty$

$$\| \|f(x, y)\|_{L^q(E, \mu(dx))}\|_{L^p(F, \nu(dy))} \leq \| \|f(x, y)\|_{L^p(F, \nu(dy))}\|_{L^q(E, \mu(dx))}. \quad (5.29)$$

For $q = p$ this holds an equality, as a consequence of Fubini's theorem. If $q < p$, the proof goes as follows: if the left-hand side of (5.29) is equal to zero, there is nothing to prove; if it is not, then raising it to power p gives, by Fubini's theorem,

$$\begin{aligned} \int_F \left(\int_E |f|^q d\mu \right)^{\frac{p}{q}} d\nu &= \int_F \left[\int_E |f|^q \left(\int_E |f|^q d\mu \right)^{\frac{p}{q}-1} d\mu \right] d\nu \\ &= \int_E \left[\int_F |f|^q \left(\int_E |f|^q d\mu \right)^{\frac{p}{q}-1} d\nu \right] d\mu \\ &\leq \int_E \left[\left(\int_F |f|^p d\nu \right)^{\frac{q}{p}} \left\{ \int_F \left(\int_E |f|^q d\mu \right)^{\frac{p-q}{q} \frac{p}{p-q}} d\nu \right\}^{\frac{p-q}{p}} \right] d\mu \\ &= \left\{ \int_F \left(\int_E |f|^q d\mu \right)^{\frac{p}{q}} d\nu \right\}^{\frac{p-q}{p}} \int_E \left(\int_F |f|^p d\nu \right)^{\frac{q}{p}} d\mu \end{aligned}$$

where we have used the Hölder inequality on (F, ν) with conjugated exponents $\frac{p}{q}$ and $\frac{p}{p-q}$. The first term in the last line is the left-hand side raised to power $\frac{p-q}{p}$: dividing by such term (which is not zero by assumption) we obtain (5.29).

Note that for $q = 1$ we have additionally, since $|\int_E f d\mu| \leq \int_E |f| d\mu$,

$$\left[\int_F \left| \int_E f d\mu \right|^p d\nu \right]^{\frac{1}{p}} \leq \int_E \left| \int_F |f|^p d\nu \right|^{\frac{1}{p}} d\mu.$$

In the special case $E = \{1, 2\}$ with $\mu = \delta_1 + \delta_2$, if we set $f_i(\cdot) := f(i, \cdot)$, then for $p \geq 1$ we recover the usual Minkowski inequality $\|f_1 + f_2\|_{L^p} \leq \|f_1\|_{L^p} + \|f_2\|_{L^p}$.

Part II

Rough Integration

CHAPTER 6

THE SEWING LEMMA

We fix throughout the chapter a time horizon $T > 0$ and two continuous functions $X, Y: [0, T] \rightarrow \mathbb{R}$. In this setting the *integral*

$$\int_0^T Y_r dX_r \tag{6.1}$$

can be defined as $\int_0^T Y_r \dot{X}_r dr$ if X is differentiable or, more generally, as a Lebesgue integral if X is of bounded variation, so that dX is a signed measure. The key question we want to address is: *how to define the integral when X does not have such regularity?* This is an example of a more general problem: given a distribution (generalized function) $\dot{X}$ and a non-smooth function Y , how to define their product $Y\dot{X}$?

A motivation is given by $X = B$ with $(B_t)_{t \geq 0}$ a Brownian motion. In this special case, one can use probability theory to answer the question and define the integral in (6.1), but one sees that there are several possible definitions: for example Itô, Stratonovich, etc.

In this book, we are going to present the alternative answer provided by the theory of Rough Paths, originally introduced by Terry Lyons. This theory yields a robust construction of the integral in (6.1) and sheds a new “pathwise” light on stochastic integration.

The approach we follow is based on the *Sewing Lemma*, to which this chapter is devoted. In particular, we will show in Chapter 7 that the integral in (6.1) has a canonical definition (*Young integral*) when Y and X are Hölder continuous, under a constraint on their Hölder exponents. Going beyond this constraint requires Rough Paths, which will be studied in Chapter 8.

6.1. LOCAL APPROXIMATION

If X is of class C^1 , we can define the integral function

$$I_t := \int_0^t Y_r \dot{X}_r dr, \quad t \in [0, T].$$

Then we have $I_0 = 0$ and for $0 \leq s \leq t \leq T$

$$I_t - I_s - Y_s(X_t - X_s) = \int_s^t (Y_r - Y_s) \dot{X}_r dr = o(t - s) \tag{6.2}$$

as $t - s \rightarrow 0$, because $\dot{X}$ is bounded and $|Y_r - Y_s| = o(1)$ as $|r - s| \rightarrow 0$. Thus the integral function I_t satisfies

$$I_0 = 0, \quad I_t - I_s = Y_s(X_t - X_s) + o(t - s), \quad 0 \leq s \leq t \leq T. \quad (6.3)$$

Remarkably, *the relation (6.3) characterizes $(I_t)_{t \in [0, T]}$* . Indeed, if I^1 and I^2 satisfy (6.3) with the same functions X, Y , their difference $\Delta := I^1 - I^2$ satisfies

$$|\Delta_t - \Delta_s| = o(t - s), \quad 0 \leq s \leq t \leq T,$$

which implies $\frac{d}{dt}\Delta_t \equiv 0$ and then $\Delta_t = \Delta_0 = I_0^1 - I_0^2 = 0$ by (6.3). This simple result deserves to be stated in a separate

LEMMA 6.1. *Given any pair of functions $X, Y: [0, T] \rightarrow \mathbb{R}$, there can be at most one function $I: [0, T] \rightarrow \mathbb{R}$ satisfying (6.3).*

The formulation (6.3) is interesting also because the derivative $\dot{X}$ of X does not appear. Therefore, if we can find a function $I: [0, T] \rightarrow \mathbb{R}$ which satisfies (6.3), such a function is *unique* and we can take it as a *definition* of the integral (6.1).

We will see in Section 7.1 that this program can be accomplished when X and Y satisfy suitable Hölder regularity assumptions. In order to get there, in the next sections we will look at a more general problem.

6.2. A GENERAL PROBLEM

Let us generalise the problem (6.3). We define $A: [0, T]_{\leq}^2 \rightarrow \mathbb{R}$ by setting for $0 \leq s \leq t \leq T$

$$A_{st} := Y_s(X_t - X_s). \quad (6.4)$$

We can then decouple (6.3) in two relations:

$$I_0 = 0, \quad I_t - I_s = A_{st} + R_{st}, \quad 0 \leq s \leq t \leq T, \quad (6.5)$$

$$R: [0, T]_{\leq}^2 \rightarrow \mathbb{R}, \quad R_{st} = o(t - s). \quad (6.6)$$

The general problem is, given a continuous $A: [0, T]_{\leq}^2 \rightarrow \mathbb{R}$, to find a pair of functions (I, R) satisfying (6.5)-(6.6). We call

- $A: [0, T]_{\leq}^2 \rightarrow \mathbb{R}$ the *germ*,
- $I: [0, T] \rightarrow \mathbb{R}$ the *integral*,
- $R: [0, T]_{\leq}^2 \rightarrow \mathbb{R}$ the *remainder*.

We are going to present conditions which allow to solve this problem.

Note that *we always have uniqueness*. Indeed, given (I^1, R^1) and (I^2, R^2) which solve (6.5)-(6.6) for the same A , by the same arguments which lead to Lemma 6.1 we have $\frac{d}{dt}(I_t^1 - I_t^2) \equiv 0$, hence $I^1 = I^2$ and then $R^1 = R^2$ by (6.5). We record this as

LEMMA 6.2. *Given any germ A , there can be at most one pair of functions (I, R) satisfying (6.5)-(6.6).*

6.3. AN ALGEBRAIC LOOK

We first focus on relation (6.5) alone. For a fixed germ A , this equation has infinitely many solutions (I, R) , because given *any* I we can simply *define* R so as to fulfill (6.5). Interestingly, all solutions admit an algebraic characterization in terms of R alone.

LEMMA 6.3. *Fix a function $A \in C_2$.*

1. *If a pair $(I, R) \in C_1 \times C_2$ satisfies (6.5), then R satisfies*

$$(\delta R)_{sut} = -(\delta A)_{sut}, \quad \forall 0 \leq s \leq u \leq t \leq T. \quad (6.7)$$

2. *Viceversa, given any function $R \in C_2$ which satisfies (6.7), if we set $I_t := A_{0t} + R_{0t}$, the pair $(I, R) \in C_1 \times C_2$ satisfies (6.5).*

Proof. Relation (6.5) clearly implies (6.7), simply because $\delta(\delta I) = 0$. Viceversa, given R satisfying (6.7), we can define $L_{st} := A_{st} + R_{st}$ so that

$$L_{st} - L_{su} - L_{ut} = 0.$$

Applying this formula to $(s', u', t') = (0, s, t)$, we obtain that $I_t := L_{0t}$ satisfies

$$I_t - I_s = L_{0t} - L_{0s} = L_{st} = A_{st} + R_{st}$$

and the proof is complete because $I_0 := L_{00} = A_{00} + R_{00} = 0$, which follows by (6.7) for $s = u = 0$. $\square$

We can now rephrase Lemma 6.3 as follows.

PROPOSITION 6.4. *Fix $A \in C_2$. Finding a pair $(I, R) \in C_1 \times C_2$ satisfying (6.5) is equivalent to finding $R \in C_2$ such that*

$$\delta R_{sut} = -\delta A_{sut}, \quad \forall 0 \leq s \leq u \leq t \leq T. \quad (6.8)$$

6.4. ENTERS ANALYSIS: THE SEWING LEMMA

So far we have analyzed (6.5). We now let (6.6) enter the game, i.e. we look for a pair of functions $(I, R) \in C_1 \times C_2$ which fulfills (6.5)-(6.6), given a (general) germ $A \in C_2$.

We stress that condition (6.6) is essential to ensure *uniqueness*: without it, equation (6.5) admits infinitely many solutions, as discussed before Lemma 6.3. When we couple (6.5) with (6.6), uniqueness is guaranteed by Lemma 6.2, but *existence* is no longer obvious. This is what we now focus on.

We start with a simple necessary condition.

LEMMA 6.5. *For (6.5)-(6.6) to admit a solution, it is necessary that the germ A satisfies*

$$|\delta A_{sut}| = o(t - s), \quad \text{for } 0 \leq s \leq u \leq t \leq T. \quad (6.9)$$

Proof. If (6.5) admits a solution, by Proposition 6.4 we have $|\delta A_{sut}| = |\delta R_{sut}|$. If furthermore R satisfies (6.6), we must have for $0 \leq s \leq u \leq t \leq T$

$$|\delta R_{sut}| \leq |R_{st}| + |R_{su}| + |R_{ut}| = o(t-s) + o(u-s) + o(t-u) = o(t-s). \quad \square$$

Remark 6.6. Choosing $u = s$ in (6.9) we obtain that $-A_{ss} = o(t-s)$, which means that $A_{ss} = 0$. Therefore a necessary condition for (6.5)-(6.6) to admit a solution is that A vanishes on the diagonal of $[0, T]_{\leq}^2$.

Remarkably, the necessary condition in Lemma 6.5 is close to being sufficient: it is enough to upgrade $o(t-s)$ in $O((t-s)^\eta)$ for some $\eta > 1$. This is the content of the celebrated *Sewing Lemma*, which we next present.

We have seen in the Sewing bound (Theorem 1.9) that any $R \in C_2$ such that $R_{st} = o(t-s)$ for $0 \leq s \leq t \leq T$ satisfies an a priori estimate $\|R\|_\eta \leq K_\eta \|\delta R\|_\eta$ for any $\eta > 1$. Of course, this estimate is only interesting if $\|\delta R\|_\eta < \infty$ for some $\eta > 1$. This property, that we call *coherence*, is at the heart of the celebrated Sewing Lemma (Gubinelli [2], Feyel-de La Pradelle [1]), as it provides a sufficient condition on the germ A for the solution of (6.5)-(6.6).

DEFINITION 6.7. (COHERENCE) *A germ $A \in C_2$ is called coherent if, for some $\eta > 1$, it satisfies $\delta A \in C_3^\eta$, i.e. $\|\delta A\|_\eta < \infty$. More explicitly:*

$$\exists \eta \in (1, \infty): \quad |\delta A_{sut}| \lesssim |t-s|^\eta, \quad 0 \leq s \leq u \leq t \leq T. \quad (6.10)$$

THEOREM 6.8. (SEWING LEMMA) *For any coherent germ $A \in C_2$ there exists a (unique) function $I: [0, T] \rightarrow \mathbb{R}$ such that $|A_{st} - \delta I_{st}| = o(t-s)$; equivalently, there exists a unique pair $(I, R) \in C_1 \times C_2$ such that*

$$I_0 = 0, \quad I_t - I_s = A_{st} + R_{st} \quad \text{with} \quad R_{st} = o(t-s). \quad (6.11)$$

- The “remainder” $R_{st} := \delta I_{st} - A_{st}$ satisfies the Sewing Bound:

$$\|R\|_\eta \leq K_\eta \|\delta A\|_\eta \quad \text{where} \quad K_\eta := (1 - 2^{1-\eta})^{-1}. \quad (6.12)$$

- The integral $I \in C_1$ is the limit of Riemann sums of the germ:

$$I_t := \lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} A_{t_i t_{i+1}} \quad (6.13)$$

along arbitrary partitions $\mathcal{P} = \{0 = t_0 < t_1 < \dots < t_k = t\}$ of $[0, t]$ with vanishing mesh $|\mathcal{P}| := \max_{i=0, \dots, k-1} |t_{i+1} - t_i| \rightarrow 0$ (we set $\#\mathcal{P} := k$).

The Sewing Lemma is a cornerstone of the theory of *Rough Paths*, to be introduced in Chapter 8. We will already see in Chapter 7 an interesting application to *Young integrals*. The (instructive) proof of Theorem 6.8 is postponed to Section 6.6.

Remark 6.9. For a fixed partition $\mathcal{P}$ of $[0, t]$ we have, by $\delta I_{st} = A_{st} + R_{st}$,

$$I_t = \sum_{i=0}^{\#\mathcal{P}-1} A_{t_i t_{i+1}} + \sum_{i=0}^{\#\mathcal{P}-1} R_{t_i t_{i+1}}.$$

Therefore, (6.13) is equivalent to

$$\lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} R_{t_i t_{i+1}} = 0$$

which is the reason why one wants the remainder R to be small close to the diagonal. The information $R_{st} = o(t - s)$ is not enough in general to obtain the existence of (I, R) , while the stronger estimate $|R_{st}| \lesssim |t - s|^\eta$ is sufficient.

6.5. THE SEWING MAP

Given a coherent germ A , by Theorem 6.8 we can find an integral I and a remainder R which solve (6.5)-(6.6). We now look closer at the remainder R .

LEMMA 6.10. *In the setting of Theorem 6.8, the remainder R is a function of δA : given two coherent germs A, A' with $\delta A = \delta A'$, the corresponding remainders R, R' coincide. Moreover, the map $\delta A \mapsto R$ is linear.*

Proof. By Proposition 6.4 we have $\delta(R - R') = \delta(A' - A) = 0$, hence $R - R' = \delta f$ for some $f \in C_1$ (see Remark 1.10). Both $|R_{st}|$ and $|R'_{st}|$ are $o(|t - s|)$ by (6.6), hence $|f_t - f_s| = o(|t - s|)$. Then f must be constant by Lemma 6.1 and therefore $R = R'$. Linearity of the map $\delta A \mapsto R$ is easy. $\square$

Since R is a function of δA , we introduce a specific notation for this map:

$$R = -\Lambda(\delta A)$$

where the minus sign is for later convenience.

Let us describe more precisely this map Λ . Throughout the following discussion, we fix arbitrarily $\eta \in (1, \infty)$.

- *Domain.* The map Λ is defined on δA for coherent germs A , see Definition 6.7. The domain of Λ is then $C_3^\eta \cap \delta C_2$, where we denote by $\delta C_2 \subseteq C_3$ the image of the space C_2 under the operator δ in (1.24).
- *Codomain.* The map Λ sends δA to $-R$, and we have $|R_{st}| \lesssim |t - s|^\eta$, see (6.12). A natural choice of codomain for Λ is then C_2^η .
- *Characterization.* In view of Proposition 6.4 and Lemma 6.2, the function $-R = \Lambda(\delta A)$ is characterized by the properties

$$\delta(-R) = \delta A, \quad |R_{st}| = o(t - s).$$

The second condition is already enforced by our choice C_2^η of codomain for Λ , which yields $|R_{st}| \lesssim |t - s|^\eta$ (with $\eta > 1$). The first relation can be rewritten as $\delta(\Lambda(B)) = B$ for all B in the domain of Λ , that is $\delta \circ \Lambda$ is the identity map.

In conclusion, we have proved the following result.

THEOREM 6.11. (SEWING MAP) *Let $\eta \in (1, \infty)$. There exists a unique map*

$$\Lambda: C_3^\eta \cap \delta C_2 \longrightarrow C_2^\eta,$$

called the Sewing Map, such that $\delta \circ \Lambda = \text{id}$ is the identity on $C_3^\eta \cap \delta C_2$.

- The map Λ is linear and satisfies

$$\|\Lambda(B)\|_\eta \leq K_\eta \|B\|_\eta \quad \forall B \in C_3^\eta \cap \delta C_2, \quad (6.14)$$

where K_η is the same constant as in (6.12).

- Given a coherent germ $A \in C_2$, i.e. such that $\delta A \in C_3^\eta$, the unique solution (I, R) of (6.5)-(6.6) is $R := -\Lambda(\delta A)$ and $I_t := A_{0t} + R_{0t}$.

6.6. PROOF OF THE SEWING LEMMA

We prove the Sewing Lemma, i.e. Theorem 6.8.

Proof. We fix a germ $A \in C_2$ with $\|\delta A\|_\eta < \infty$ for some $\eta > 1$ (we do *not* require $A_{ab} = o(b-a)$). Our goal is to build a function $I: [0, T] \rightarrow \mathbb{R}$ such that $|\delta I_{st} - A_{st}| = o(t-s)$. Uniqueness of I follows by Lemma 6.2, while the bound (6.12) follows by the Sewing Bound (1.27) applied to $R_{st} := \delta I_{st} - A_{st}$ (note that $\delta R = -\delta A$, because $\delta \circ \delta = 0$).

We fix $0 \leq s < t \leq T$. Given a partition $\mathcal{P} = \{s = t_0 < t_1 < \dots < t_m = t\}$ of $[s, t]$, let us define $I_{\mathcal{P}}(A) := \sum_{i=0}^{m-1} A_{t_i t_{i+1}}$ as in (1.21). The following bound holds:

$$|I_{\mathcal{P}}(A) - A_{st}| \leq C_\eta \|\delta A\|_\eta (t-s)^\eta \quad \text{with} \quad C_\eta := \sum_{n \geq 1} \frac{2^n}{n^\eta} < \infty, \quad (6.15)$$

as we showed in the proof of Theorem 1.18, see (1.48). Similarly, if $\mathcal{Q} \supseteq \mathcal{P}$ is another partition of $[s, t]$,

$$\begin{aligned} |I_{\mathcal{Q}}(A) - I_{\mathcal{P}}(A)| &\leq \sum_{i=0}^{\#\mathcal{P}-1} |I_{\mathcal{Q} \cap [t_i, t_{i+1}]}(A) - A_{t_i t_{i+1}}| \\ &\leq C_\eta \|\delta A\|_\eta \sum_{i=0}^{\#\mathcal{P}-1} (t_{i+1} - t_i)^\eta \\ &\leq C_\eta \|\delta A\|_\eta |\mathcal{P}|^{\eta-1} \sum_{i=0}^{\#\mathcal{P}-1} (t_{i+1} - t_i) \\ &\leq C_\eta \|\delta A\|_\eta T |\mathcal{P}|^{\eta-1} \end{aligned}$$

where we recall that $|\mathcal{P}| := \max_i (t_{i+1} - t_i)$. Finally, if $\mathcal{P}$ and $\mathcal{P}'$ are arbitrary partitions, setting $\mathcal{Q} := \mathcal{P} \cup \mathcal{P}'$ and applying the triangle inequality yields

$$|I_{\mathcal{P}'}(A) - I_{\mathcal{P}}(A)| \leq C_\eta \|\delta A\|_\eta T (|\mathcal{P}|^{\eta-1} + |\mathcal{P}'|^{\eta-1}).$$

This shows that the family $I_{\mathcal{P}}(A)$ is Cauchy as $|\mathcal{P}| \rightarrow 0$ (for every $\epsilon > 0$ there exists $\delta_\epsilon > 0$ such that $|\mathcal{P}|, |\mathcal{P}'| \leq \delta_\epsilon$ implies $|I_{\mathcal{P}'}(A) - I_{\mathcal{P}}(A)| \leq \epsilon$), hence it admits a limit as $|\mathcal{P}| \rightarrow 0$, that we call J_{st} .

We now define $I_t := J_{0t}$. We claim that

$$I_t - I_s = J_{st} \quad \text{for all } 0 \leq s < t \leq T.$$

Indeed, if we consider partitions $\mathcal{P}'$ on $[0, s]$ and $\mathcal{P}$ of $[s, t]$, then $\mathcal{P}'' := \mathcal{P} \cup \mathcal{P}'$ is a partition of $[0, t]$ such that $I_{\mathcal{P}''}(A) - I_{\mathcal{P}'}(A) = I_{\mathcal{P}}(A)$, and taking the limit of vanishing mesh we get $J_{0t} - J_{0s} = J_{st}$, that is the claim.

Finally, taking the limit of relation (6.15), since $I_{\mathcal{P}}(A) \rightarrow J_{st} = I_t - I_s$, we obtain our goal $|\delta I_{st} - A_{st}| \lesssim (t - s)^\eta = o(t - s)$. This completes the proof, since (6.13) holds by construction. $\square$

Remark 6.12. Taking the limit of (6.15) gives

$$|R_{st}| \leq C_\eta \|\delta A\|_\eta |t - s|^\eta, \quad R_{st} := \delta I_{st} - A_{st}, \quad 0 \leq s < t \leq T,$$

which is the bound (6.12) with K_η replaced by the worse constant C_η . This is because the estimate (6.15) holds for arbitrary partitions.

CHAPTER 7

THE YOUNG INTEGRAL

We can now come back to the problem that we discussed at the beginning of Chapter 6: given two continuous functions $X, Y: [0, T] \rightarrow \mathbb{R}$, how can we give a meaning to the integral $I_t = \int_0^t Y dX$ for $t \in [0, T]$?

A natural answer, recall (6.3), is to look for a function $I: [0, T] \rightarrow \mathbb{R}$ satisfying

$$I_0 = 0, \quad I_t - I_s = Y_s(X_t - X_s) + o(t - s), \quad 0 \leq s \leq t \leq T. \quad (7.1)$$

As an application of the *Sewing Lemma* (Theorem 6.8), we can show that such a function I exists (and is necessarily unique) when X and Y are Hölder functions of exponents $\alpha, \beta \in]0, 1]$ such that $\alpha + \beta > 1$. This leads to the notion of *Young integral*, to which this chapter is devoted.

Going beyond this setting, in order to treat the case $\alpha + \beta \leq 1$, will require the notion of *Rough Paths*, that we discuss in Chapter 8.

7.1. CONSTRUCTION OF THE YOUNG INTEGRAL

As we did in Chapter 6, it is convenient to rewrite (7.1) as follows: we look for a function $I: [0, T] \rightarrow \mathbb{R}$ satisfying

$$I_0 = 0, \quad I_t - I_s = A_{st} + R_{st} \quad \text{with} \quad R_{st} = o(t - s), \quad (7.2)$$

where the *germ* $A: [0, T]^2 \rightarrow \mathbb{R}$ is defined by

$$A_{st} = Y_s \delta X_{st} = Y_s(X_t - X_s). \quad (7.3)$$

This is the framework of the *Sewing Lemma*, see Theorem 6.8, for which we need to fulfill the *coherence condition* (6.10), that is $\|\delta A\|_\eta < \infty$ for some $\eta > 1$ (we use the norms introduced in (1.9)). Recalling that

$$\delta A_{sut} := A_{st} - A_{su} - A_{ut} = -\delta Y_{su} \delta X_{ut},$$

see (1.33), we can write for any $\alpha, \beta \in]0, 1]$

$$|\delta A_{sut}| = |Y_u - Y_s| |X_t - X_u| \implies \|\delta A\|_{\alpha+\beta} \leq \|\delta X\|_\alpha \|\delta Y\|_\beta. \quad (7.4)$$

As a consequence, it is natural to assume that $\|\delta X\|_\alpha < \infty$ and $\|\delta Y\|_\beta < \infty$ for $\alpha, \beta \in]0, 1]$ such that $\alpha + \beta > 1$.

We can now give a consistent definition of the integral $I_t = \int_0^t Y \, dX$, known as *Young integral*, when X and Y are suitable Hölder functions.

THEOREM 7.1. (YOUNG INTEGRAL) *Fix $\alpha, \beta \in]0, 1]$ with $\alpha + \beta > 1$. For every $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ there is a (necessarily unique) function $I: [0, T] \rightarrow \mathbb{R}$ which satisfies (7.1), i.e.*

$$I_0 = 0, \quad I_t - I_s = Y_s(X_t - X_s) + o(t - s). \quad (7.5)$$

The function I , called the Young integral, is also denoted by $I_t =: \int_0^t Y \, dX$.

The remainder $R_{st} := I_t - I_s - Y_s(X_t - X_s)$ satisfies the bound

$$\|R\|_{\alpha+\beta} \leq K_{\alpha+\beta} \|\delta X\|_\alpha \|\delta Y\|_\beta, \quad (7.6)$$

where $K_\eta := (1 - 2^{1-\eta})^{-1}$, see (6.12). This yields $I \in \mathcal{C}^\alpha$, more precisely

$$\|\delta I\|_\alpha \leq (\|Y\|_\infty + K_{\alpha+\beta} T^\beta \|\delta Y\|_\beta) \|\delta X\|_\alpha. \quad (7.7)$$

The Young integral $I = (I_t)_{t \in [0, T]}$, as a function of (X, Y) , is a continuous bilinear map $I: \mathcal{C}^\alpha \times \mathcal{C}^\beta \rightarrow \mathcal{C}^\alpha$.

Proof. Recalling (7.2)-(7.4), we have $\|\delta A\|_{\alpha+\beta} \leq \|\delta X\|_\alpha \|\delta Y\|_\beta < \infty$, that is $\delta A \in C_3^\eta$ with $\eta = \alpha + \beta > 1$, where the spaces C_k^η were defined in (1.10). By the *Sewing Lemma*, see Theorem 6.8, there exists a (unique) function I which satisfies (6.11) and (6.12), hence (7.5) and (7.6) hold.

In order to prove (7.7), we note that

$$\begin{aligned} \|\delta I\|_\alpha &\leq \|A\|_\alpha + \|R\|_\alpha \leq \|Y\|_\infty \|\delta X\|_\alpha + T^\beta \|R\|_{\alpha+\beta} \\ &\leq \|Y\|_\infty \|\delta X\|_\alpha + T^\beta K_{\alpha+\beta} \|\delta X\|_\alpha \|\delta Y\|_\beta. \end{aligned}$$

Recalling Remark 1.4, in particular (1.15), this bound implies that I is a continuous function of (X, Y) , as a map from $\mathcal{C}^\alpha \times \mathcal{C}^\beta$ to $\mathcal{C}^\alpha$.

We finally prove that the map $(X, Y) \mapsto I$ is bilinear: given $X, X' \in \mathcal{C}^\alpha$ and a fixed $Y \in \mathcal{C}^\beta$, if I satisfies (7.5) for (X, Y) and I' satisfies (7.5) for (X', Y) , then for any $a, b \in \mathbb{R}$ the function $\hat{I}_t := a I_t + b I'_t$ satisfies (7.5) for $(\hat{X} := aX + bX', Y)$. Linearity with respect to Y is proved similarly. $\square$

By (6.13) in Theorem 6.8, we also have that the Young integral (7.5) is the limit of Riemann sums

$$I_t = \lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} Y_{t_i} (X_{t_{i+1}} - X_{t_i})$$

along arbitrary partitions $\mathcal{P} = \{0 = t_0 < t_1 < \dots < t_k = t\}$ of $[0, t]$ with vanishing mesh $|\mathcal{P}| := \max_{i=0, \dots, k-1} |t_{i+1} - t_i| \rightarrow 0$ (with $\#\mathcal{P} := k$).

Remark 7.2. The setting of Theorem 7.1 provides a natural example of a germ $A_{st} := Y_s \delta X_{st}$ which is *not* in C_2^η for any $\eta > 1$ (excluding the trivial case when $Y \equiv 0$ on the intervals where X is not constant, hence $A \equiv 0$), but it satisfies $\delta A \in C_3^\eta$ with $\eta = \alpha + \beta > 1$.

Remark 7.3. (BEYOND YOUNG) It is natural to wonder what happens in Theorem 7.1 for $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta \leq 1$. In this case, *there might be no solution to (6.5)-(6.6)*, because the necessary condition (6.9) in Lemma 6.5 can fail. For a simple example, consider $X_t = t^\alpha$ and $Y_t = t^\beta$ for $t \in [0, T]$ and note that for $s = 0$ and $u = \frac{t}{2}$ we have by (1.33)

$$|\delta A_{sut}| = |\delta A_{0 \frac{t}{2} t}| = \left| \delta Y_{0 \frac{t}{2}} \right| \left| \delta X_{\frac{t}{2} t} \right| = \left(\frac{t}{2} \right)^\beta \left(t^\alpha - \left(\frac{t}{2} \right)^\alpha \right) \gtrsim t^{\alpha+\beta}, \quad (7.8)$$

which is not $o(t-s) = o(t)$ when $\alpha + \beta \leq 1$.

In order to define a notion of integral $I_t = \int_0^t Y_s dX_s$ when $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta \leq 1$, we need to relax condition (6.3), see Definition 8.1 below. This will lead to the notion of *Rough Paths*, described in Chapter 8.

7.2. ITÔ VERSUS YOUNG

Let $(B_t)_{t \in [0, T]}$ be a standard Brownian motion and $(h_t)_{t \in [0, T]}$ and adapted process. Let $\alpha, \beta \in (0, 1)$ such that $\alpha < \frac{1}{2}$ and $\alpha + \beta > 1$. We know that a.s. B is of class $\mathcal{C}^\alpha$, we suppose that a.s. h is of class $\mathcal{C}^\beta$. In this case we have

PROPOSITION 7.4. *Under the conditions above, the Young integral $I(B, h)$ constructed in (7.5) is a.s. equal to the Itô integral $(\int_0^t h_s dB_s)_{t \in [0, T]}$. In particular a.s. for all $t \in [0, T]$*

$$\int_0^t h_s dB_s = \lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} h_{t_i} (B_{t_{i+1}} - B_{t_i})$$

along arbitrary (random) partitions $\mathcal{P} = \{0 = t_0 < t_1 < \dots < t_k = t\}$ of $[0, t]$ with vanishing mesh $|\mathcal{P}| := \max_{i=0, \dots, k-1} |t_{i+1} - t_i| \rightarrow 0$ (with $\#\mathcal{P} := k$).

Proof. By (4.8) in Theorem 4.3 we know that the Itô integral $I_t := \int_0^t h_s dB_s$, $t \in [0, T]$, satisfies a.s.

$$|\delta I_{st} - h_s \delta B_{st}| \lesssim |t - s|^{\alpha+\beta}, \quad 0 \leq s \leq t \leq T.$$

Since $\alpha + \beta > 1$, I coincides a.s. with the Young integral $I(B, h)$. The convergence of the Riemann sums follows from (6.13). $\square$

7.3. INTEGRAL FORMULATION OF YOUNG EQUATIONS

In this section we explain why we call (2.4) a *Young equation*. In fact, we can interpret the finite difference equation (2.4) as an *integral equation*, using the Young integral of section 7.1.

PROPOSITION 7.5. *Let $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^k)$ with $\alpha > \frac{1}{2}$. Then Z satisfies (2.4) if and only if*

$$Z_t = Z_0 + \int_0^t \sigma(Z_s) dX_s, \quad t \in [0, T], \quad (7.9)$$

where the integral is in the Young sense.

Proof. We consider the germ $A_{st} := \sigma(Z_s) \delta X_{st}$, $0 \leq s \leq t \leq T$. By (7.4)

$$|\delta A_{sut}| = |\sigma(Z_u) - \sigma(Z_s)| |X_t - X_u| \implies \|\delta A\|_{2\alpha} \leq \|\nabla \sigma\|_\infty \|\delta X\|_\alpha \|\delta Z\|_\alpha.$$

Therefore we obtain that (2.4) is equivalent to (7.5) above. $\square$

In the case $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, this argument does not work and the Young integral is not adapted, since the germ $A_{st} := \sigma(Z_s) \delta X_{st}$ has the property $\delta A \in C_3^{2\alpha}$ with $2\alpha \leq 1$, so that the Sewing Lemma can not be applied. We have already seen in Chapter 3, see equation (3.19), how the germ A must be modified in this case. We are going to come back to this in the following chapters.

As an application of the estimates on the Young integral of Theorem 7.1, we want to give a local existence result for equation (2.4) which does not rely on compactness and which can be therefore used also in infinite dimension.

Let $Z_0 \in \mathbb{R}^k$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ be given, $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^d \otimes (\mathbb{R}^d)^*$ smooth and the unknown $Z: [0, T] \rightarrow \mathbb{R}^k$ is such that $\sigma(Z) \in \mathcal{C}^\alpha$ and $2\alpha > 1$, so that the right-hand side of (7.9) can be interpreted as a Young integral. We want now to show the following

THEOREM 7.6. (CONTRACTION FOR YOUNG DIFFERENTIAL EQUATIONS) *Let $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^d \otimes (\mathbb{R}^d)^*$ be of class C^2 with $\nabla \sigma$ and $\nabla^2 \sigma$ bounded. Let $\alpha \in]\frac{1}{2}, 1]$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ fixed. If $T > 0$ is small enough, then for any $Z_0 \in \mathbb{R}^k$ there exists a unique $Z \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^k)$ which satisfies (7.9).*

Proof. For all $f \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^k)$ we have

$$|\sigma(f_t) - \sigma(f_s)| \leq \|\nabla \sigma\|_\infty |f_t - f_s|$$

so that

$$\|\delta \sigma(f)\|_\alpha \leq \|\nabla \sigma\|_\infty \|\delta f\|_\alpha.$$

By (7.7) with $\alpha = \beta$ we obtain for all $f \in \mathcal{C}^\alpha$ satisfying (7.9)

$$\|\delta f\|_\alpha \leq (|\sigma(f_0)| + (1 + K_{2\alpha}) T^\alpha \|\nabla \sigma\|_\infty \|\delta f\|_\alpha) \|\delta X\|_\alpha$$

since

$$\|\sigma(f)\|_\infty \leq |\sigma(f_0)| + T^\alpha \|\delta \sigma(f)\|_\alpha.$$

Therefore, if T satisfies

$$T^\alpha \leq \frac{1}{2(1 + K_{2\alpha}) \|\nabla \sigma\|_\infty \|\delta X\|_\alpha}$$

then we have the following a priori estimate on solutions to (7.9)

$$\|\delta Z\|_\alpha \leq 2|\sigma(Z_0)| \|\delta X\|_\alpha.$$

We fix such T and we set $\mathcal{C}^\alpha(Z_0) := \{f \in \mathcal{C}^\alpha: f_0 = Z_0, \|\delta f\|_\alpha \leq 2|\sigma(Z_0)| \|\delta X\|_\alpha\}$. Then we define $\Lambda: \mathcal{C}^\alpha \rightarrow \mathcal{C}^\alpha$ given by

$$\Lambda(f) := h, \quad h_t := Z_0 + \int_0^t \sigma(f_s) dX_s, \quad t \in [0, T].$$

It is easy to see, arguing as above, that Λ acts on $\mathcal{C}^\alpha(Z_0)$, namely $\Lambda: \mathcal{C}^\alpha(Z_0) \rightarrow \mathcal{C}^\alpha(Z_0)$. Note that the map $\mathcal{C}^\alpha(Z_0) \times \mathcal{C}^\alpha(Z_0) \ni (a, b) \mapsto \|\delta a - \delta b\|_\alpha$ defines a distance on $\mathcal{C}^\alpha(Z_0)$ which induces the same topology as $\|\cdot\|_{\mathcal{C}^\alpha}$. We want to show that Λ is a contraction for this distance if T is small enough. By (7.7) we have for $\alpha = \beta$

$$\begin{aligned} \|\delta \Lambda(a) - \delta \Lambda(b)\|_\alpha &\leq (\|\sigma(a) - \sigma(b)\|_\infty + K_{2\alpha} T^\alpha \|\delta \sigma(a) - \delta \sigma(b)\|_\alpha) \|\delta X\|_\alpha \\ &\leq T^\alpha (1 + K_{2\alpha}) \|\delta X\|_\alpha \|\delta \sigma(a) - \delta \sigma(b)\|_\alpha. \end{aligned}$$

We now need to estimate $\|\delta \sigma(a) - \delta \sigma(b)\|_\alpha$. By Lemma 2.8

$$\|\delta \sigma(a) - \delta \sigma(b)\|_\alpha \leq \|\nabla \sigma\|_\infty \|\delta a - \delta b\|_\alpha + \|\nabla^2 \sigma\|_\infty (\|\delta a\|_\alpha + \|\delta b\|_\alpha) \|a - b\|_\infty.$$

Since, as usual, $\|a - b\|_\infty \leq T^\alpha \|\delta a - \delta b\|_\alpha$, we obtain

$$\|\delta \sigma(a) - \delta \sigma(b)\|_\alpha \leq (\|\nabla \sigma\|_\infty + T^\alpha \|\nabla^2 \sigma\|_\infty (\|\delta a\|_\alpha + \|\delta b\|_\alpha)) \|\delta a - \delta b\|_\alpha. \quad (7.10)$$

Therefore, for all $a, b \in \mathcal{C}^\alpha(Z_0)$

$$\|\delta \Lambda(a) - \delta \Lambda(b)\|_\alpha \leq C_T \|\delta a - \delta b\|_\alpha,$$

where $C_T := T^\alpha (1 + K_{2\alpha}) \|\delta X\|_\alpha (\|\nabla \sigma\|_\infty + T^\alpha \|\nabla^2 \sigma\|_\infty 4|\sigma(Z_0)| \|\delta X\|_\alpha)$. It is now enough to consider T small enough so that $C_T < 1$. $\square$

7.4. PROPERTIES OF THE YOUNG INTEGRAL

The Young integral $\int_0^t Y dX$, defined in Theorem 7.1, shares many properties with the classical Riemann-Lebesgue integral, that we now discuss. For an interval $[s, t] \subseteq [0, T]$ we will use the notation

$$I_t - I_s =: \int_s^t Y dX.$$

A elementary but useful observation is that $(X, Y) \rightarrow \int_s^t Y dX$ is a bilinear function. If the integrand $Y_u = c$ is constant for all $u \in [s, t]$, then $\int_s^t Y dX = c(X_t - X_s)$, which follows directly from (7.5). As a corollary, we obtain the following useful formula for the remainder.

LEMMA 7.7. *Let $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ for $\alpha, \beta \in]0, 1]$ with $\alpha + \beta > 1$ and let $I_t := \int_0^t Y_u dX_u$ be the Young integral, see Theorem 7.1. Then the remainder*

$$R_{st} := I_t - I_s - Y_s(X_t - X_s), \quad 0 \leq s \leq t \leq T,$$

admits the explicit formula

$$R_{st} = \int_s^t (Y_u - Y_s) dX_u, \quad 0 \leq s \leq t \leq T, \quad (7.11)$$

where the right hand side is a Young integral.

Proof. By linearity and the basic property mentioned above, we obtain

$$\int_s^t (Y_u - Y_s) dX_u = \int_s^t Y_u dX_u - \int_s^t Y_s dX_u = I_t - I_s - Y_s(X_t - X_s) = R_{st}. \quad \square$$

An important property is *integration by parts*, which follows by the uniqueness of the solution for the problem (6.5)-(6.6), recall Lemma 6.2.

PROPOSITION 7.8. (INTEGRATION BY PARTS) *Fix $\alpha, \beta \in]0, 1]$ with $\alpha + \beta > 1$. For all $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ the Young integral satisfies*

$$\int_0^t X dY + \int_0^t Y dX = X_t Y_t - X_0 Y_0. \quad (7.12)$$

Proof. Let us set $I'_t := \int_0^t X dY + \int_0^t Y dX$. By the property (7.5) we have

$$I'_t - I'_s = \underbrace{Y_s(X_t - X_s) + X_s(Y_t - Y_s)}_{A_{st}} + o(t - s).$$

Next we set $I''_t := X_t Y_t - X_0 Y_0$ and note that, by direct computation,

$$I''_t - I''_s = \underbrace{Y_s(X_t - X_s) + X_s(Y_t - Y_s)}_{A_{st}} + \underbrace{(X_t - X_s)(Y_t - Y_s)}_{R_{st}},$$

where $|R_{st}| \leq \|\delta X\|_\alpha \|\delta Y\|_\beta |t - s|^{\alpha+\beta} = o(t - s)$. By Lemma 6.2, for any germ A , there can be at most one function I which satisfies $\delta I_{st} = A_{st} + o(t - s)$ (6.5)-(6.6), hence $I' = I''$. $\square$

We next discuss the *chain rule*.

PROPOSITION 7.9. (CHAIN RULE) *Let $X \in \mathcal{C}^\alpha$ with $\alpha \in]\frac{1}{2}, 1]$. Let $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ be differentiable with $\varphi' \in \mathcal{C}^\gamma(\mathbb{R})$, for $\gamma \in]0, 1]$ such that $\gamma > \frac{1}{\alpha} - 1$ (a sufficient condition is that $\varphi \in C^2$). Then $\varphi'(X) = \varphi' \circ X \in \mathcal{C}^{\alpha\gamma}$ and*

$$\varphi(X_t) - \varphi(X_0) = \int_0^t \varphi'(X) dX, \quad (7.13)$$

where the right hand side is a Young integral.

Proof. It is easy to see that $\varphi'(X) \in \mathcal{C}^{\alpha\gamma}$, which implies that $\int_0^t \varphi'(X) dX$ is well-defined as a Young integral, since $\alpha + \alpha\gamma > 1$. By the definition (7.5) of the Young integral, proving (7.13) amounts to showing that

$$|\varphi(X_t) - \varphi(X_s) - \varphi'(X_s)(X_t - X_s)| \lesssim o(t - s).$$

By the classical Lagrange theorem, if, say, $X_t > X_s$, then

$$\varphi(X_t) - \varphi(X_s) - \varphi'(X_s)(X_t - X_s) = (\varphi'(\xi) - \varphi'(X_s))(X_t - X_s)$$

with $\xi \in]X_s, X_t[$. Since $\varphi' \in \mathcal{C}^\gamma$ and $X \in \mathcal{C}^\alpha$, it follows that

$$|\varphi(X_t) - \varphi(X_s) - \varphi'(X_s)(X_t - X_s)| \lesssim |X_t - X_s|^{\gamma+1} = o(t - s)$$

since $\gamma + 1 > \frac{1}{\alpha} \geq 1$. This completes the proof. $\square$

More generally, we have

COROLLARY 7.10. *In the same setting of Proposition 7.9, for all $s \leq t$*

$$\varphi(X_t) - \varphi(X_s) = \varphi'(X_s)(X_t - X_s) + \int_s^t (\varphi'(X_r) - \varphi'(X_s)) dX_r. \quad (7.14)$$

Proof. It is enough to note that, by (7.13),

$$\begin{aligned} \varphi(X_t) - \varphi(X_s) &= \int_s^t \varphi'(X_r) dX_r \\ &= \varphi'(X_s)(X_t - X_s) + \int_s^t (\varphi'(X_r) - \varphi'(X_s)) dX_r, \end{aligned}$$

where all integrals are in the Young sense. $\square$

In particular, for $X \in \mathcal{C}^\alpha$ with $\alpha > \frac{1}{2}$, we have

$$\frac{X_t^2}{2} - \frac{X_s^2}{2} = X_s(X_t - X_s) + \int_s^t (X_r - X_s) dX_r, \quad (7.15)$$

which can be rewritten as follows:

$$\int_s^t (X_r - X_s) dX_r = \frac{X_t^2}{2} - \frac{X_s^2}{2} - X_s(X_t - X_s) = \frac{(X_t - X_s)^2}{2}. \quad (7.16)$$

7.5. MORE ON HÖLDER SPACES

We discuss further properties of the Hölder spaces $\mathcal{C}^\alpha$ for $\alpha \in (0, 1)$ (excluding the case $\alpha = 1$ of Lipschitz functions). These will be useful in the next Section 7.6, when we discuss the uniqueness of the Young integral.

Let us denote by C^∞ the space of infinitely differentiable functions. We note that $C^\infty \subset \mathcal{C}^\alpha$ for every $\alpha \in (0, 1)$, but C^∞ is not dense in $\mathcal{C}^\alpha$.

THEOREM 7.11. *For any $\alpha \in (0, 1)$, the closure of C^∞ in $\mathcal{C}^\alpha$ is the subset $\mathcal{C}_0^\alpha$ defined by*

$$\mathcal{C}_0^\alpha := \{f: [0, T] \rightarrow \mathbb{R} : |f(t) - f(s)| = o(|t - s|^\alpha) \text{ uniformly as } |t - s| \rightarrow 0\}.$$

Remark 7.12. Note that $f \in \mathcal{C}_0^\alpha$ if and only if

$$\forall \epsilon > 0 \quad \exists \delta_\epsilon > 0: \quad |f(t) - f(s)| \leq \epsilon |t - s|^\alpha \quad \text{for } |t - s| \leq \delta_\epsilon, \quad (7.17)$$

which implies (exercise) that $C^1 \subset \mathcal{C}_0^\alpha \subset \mathcal{C}^\alpha$ for $\alpha \in (0, 1)$. It follows that the closure of C^1 in $\mathcal{C}^\alpha$ is again $\mathcal{C}_0^\alpha$, simply because $C^\infty \subset C^1 \subset \mathcal{C}_0^\alpha$.

Exercise 7.1. Prove that $C^1 \subset \mathcal{C}_0^\alpha$ and $\mathcal{C}_0^\alpha \subset \mathcal{C}^\alpha$ for $\alpha \in (0, 1)$ (inclusions are strict).

We stress that the subset $\mathcal{C}_0^\alpha$ is strictly included in $\mathcal{C}^\alpha$, but what is left out is not so large, in the following sense.

Exercise 7.2. Prove that $\mathcal{C}^{\alpha'} \subset \mathcal{C}_0^\alpha$ for $0 < \alpha < \alpha' < 1$ (the inclusion is strict).

The proof of Theorem 7.11, which we defer to Section 7.7, is based on the following classical approximation result (also proved in Section 7.7).

LEMMA 7.13. *For any continuous $f: [0, T] \rightarrow \mathbb{R}$ there is a sequence $f_n \in C^\infty$ such that $\|f_n - f\|_\infty \rightarrow 0$. One can take f_n with the same modulus of continuity as f , in the following sense: given an arbitrary function $h(\cdot)$,*

$$\begin{aligned} \text{if } & |f(t) - f(s)| \leq h(t-s) \quad \forall s, t \in [0, T], \\ \text{then } & |f_n(t) - f_n(s)| \leq h(t-s) \quad \forall s, t \in [0, T], \quad \forall n \in \mathbb{N}. \end{aligned} \quad (7.18)$$

It follows that $\|\delta f_n\|_\alpha \leq \|\delta f\|_\alpha$ for all $n \in \mathbb{N}$ and $\alpha \in (0, 1)$.

Remark 7.14. Lemma 7.13 holds with no change for functions $f: [0, T] \rightarrow R$, where R is an arbitrary Banach space. One only needs a notion of integral $\int_0^T f_s ds$ when f is continuous, and for this one can take the Riemann integral, i.e. the limit of Riemann sums $\sum_i f(t_i)(t_{i+1} - t_i)$ along partitions (t_i) of $[0, T]$ with vanishing mesh $\max_i |t_{i+1} - t_i| \rightarrow 0$ (one can check that such Riemann sums form a Cauchy family). This integral satisfies the key usual properties: $f \mapsto \int_0^T f_s ds$ is linear, $|\int_0^T f_s ds| \leq \int_0^T |f_s| ds$ and $\int_0^T f'_s ds = f_T - f_0$.

7.6. UNIQUENESS OF THE YOUNG INTEGRAL

Throughout this section we denote by I_t^{Young} the *Young integral* $I_t = \int_0^t Y dX$ built in Theorem 7.1. We want to compare it with the *classical integral*

$$I_t^{\text{classical}} := \int_0^t Y_u \dot{X}_u du$$

which is defined for continuous Y and continuously differentiable $X \in C^1$.

We remarked in (6.2)-(6.3) that $I_t^{\text{classical}}$ satisfies property (7.5), therefore $I_t^{\text{classical}}$ coincides with I_t^{Young} when $(X, Y) \in C^1 \times \mathcal{C}^\beta$, for any $\beta \in]0, 1]$. In other terms, the *Young integral is an extension of the classical integral*.

We can be more precise: by Theorem 7.1, for $\alpha, \beta \in]0, 1]$ with $\alpha + \beta > 1$, the Young integral $I^{\text{Young}} = (I_t^{\text{Young}})_{t \in [0, T]}$ is a continuous bilinear map from $\mathcal{C}^\alpha \times \mathcal{C}^\beta$ to $\mathcal{C}^\alpha$. This means that I^{Young} is a *continuous* extension of the classical integral $I^{\text{classical}}$ defined on $C^1 \times \mathcal{C}^\beta$. It would be tempting to state that it is the *unique* continuous extension, but *this is not true*, because $C^1 \subset \mathcal{C}^\alpha$ is not dense in $\mathcal{C}^\alpha$ (see Theorem 7.11 and Remark 7.12).

However we can characterize the Young integral as the unique continuous extension of the classical integral on $\mathcal{C}^\alpha \times \mathcal{C}^\beta$ for fixed α , provided we consider a slightly weaker notion of convergence on $\mathcal{C}^\alpha$.

DEFINITION 7.15. Fix $\alpha \in]0, 1]$. Given $f_n, f: [0, T] \rightarrow \mathbb{R}$, with $n \in \mathbb{N}$, we write

$$f_n \rightsquigarrow_\alpha f \quad \Longleftrightarrow \quad \|f_n - f\|_\infty \rightarrow 0 \quad \text{and} \quad \sup_{n \in \mathbb{N}} \|\delta f_n\|_\alpha < \infty. \quad (7.19)$$

In other terms, $f_n \rightsquigarrow_\alpha f$ if and only if $f_n \rightarrow f$ in the sup-norm and, moreover, the sequence f_n is bounded in $\mathcal{C}^\alpha$.

We leave it as an exercise to check some basic properties.

Exercise 7.3. Fix $\alpha \in]0, 1]$ and let $f_n, f: [0, T] \rightarrow \mathbb{R}$, with $n \in \mathbb{N}$. Prove the following.

1. If $f_n \rightsquigarrow_\alpha f$, then $f \in \mathcal{C}^\alpha$; more precisely $\|\delta f\|_\alpha \leq \sup_{n \in \mathbb{N}} \|\delta f_n\|_\alpha < \infty$.
2. If $f_n \rightsquigarrow_\alpha f$, then $f_n \rightarrow f$ in $\mathcal{C}^{\alpha'}$ for any $\alpha' < \alpha$, but not necessarily $f_n \rightarrow f$ in $\mathcal{C}^\alpha$.
3. If $f_n \rightsquigarrow_\alpha f$ and $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz, then $\varphi(f_n) \rightsquigarrow_\alpha \varphi(f)$.
4. In the definition (7.19) of $f_n \rightsquigarrow_\alpha f$, the uniform convergence $\|f_n - f\|_\infty \rightarrow 0$ can be replaced by pointwise convergence: $f_n(t) \rightarrow f(t)$ for every $t \in [0, T]$.

We can now provide the following characterization of the Young integral.

THEOREM 7.16. (CHARACTERIZATION OF THE YOUNG INTEGRAL, II) Fix $\alpha, \beta \in]0, 1]$ with $\alpha + \beta > 1$. The Young integral $I^{\text{Young}} = (I_t^{\text{Young}})_{t \in [0, T]}$ is the unique map $I: \mathcal{C}^\alpha \times \mathcal{C}^\beta \rightarrow \mathcal{C}^\alpha$ such that:

1. $I_t = I_t^{\text{classical}} = \int_0^t Y_u \dot{X}_u du$ for $X \in C^1$;
2. if $X_n \rightsquigarrow_\alpha X$ and $Y_n \rightsquigarrow_\beta Y$, we have $I(X_n, Y_n) \rightsquigarrow_\alpha I(X, Y)$.

Proof. We already know that the Young integral I^{Young} satisfies property 1. Let us show that it also satisfies property 2: given $X_n \rightsquigarrow_\alpha X$ and $Y_n \rightsquigarrow_\beta Y$, we need to prove that

$$I^{\text{Young}}(X_n, Y_n) \rightsquigarrow_\alpha I^{\text{Young}}(X, Y). \quad (7.20)$$

Let us fix $\alpha' < \alpha$, $\beta' < \beta$ such that we still have $\alpha' + \beta' > 1$. We know by Exercise 7.3 that $X_n \rightarrow X$ in $\mathcal{C}^{\alpha'}$ and $Y_n \rightarrow Y$ in $\mathcal{C}^{\beta'}$. Since the Young integral is a continuous bilinear operator $I^{\text{Young}}: \mathcal{C}^{\alpha'} \times \mathcal{C}^{\beta'} \rightarrow \mathcal{C}^{\beta'}$, we have the convergence $I^{\text{Young}}(X_n, Y_n) \rightarrow I^{\text{Young}}(X, Y)$ in $\mathcal{C}^{\alpha'}$, which implies

$$\|I^{\text{Young}}(X_n, Y_n) - I^{\text{Young}}(X, Y)\|_\infty \rightarrow 0.$$

To prove (7.20), it remains to observe that, by (7.7),

$$\sup_n \|I^{\text{Young}}(X_n, Y_n)\|_\alpha \leq \sup_n (\|Y_n\|_\infty + K_{\alpha+\beta} T^\alpha \|\delta Y_n\|_\beta) \|X_n\|_\alpha < \infty.$$

We next consider an operator $I: \mathcal{C}^\alpha \times \mathcal{C}^\beta \rightarrow \mathcal{C}^\alpha$ which satisfies properties 1 and 2 and we show that it must coincide with the Young integral I^{Young} . Given $X \in \mathcal{C}^\alpha$ and $Y \in \mathcal{C}^\beta$, by Lemma 7.13 we can construct a sequence $(X_n) \subset C^1$ with $\|X_n - X\|_\infty \rightarrow 0$ and $\|X_n\|_\alpha \leq \|X\|_\alpha$. By property 2 we have $I(X_n, Y) \rightsquigarrow_\alpha I(X, Y)$ and $I^{\text{Young}}(X_n, Y) \rightsquigarrow_\alpha I^{\text{Young}}(X, Y)$, which implies pointwise convergence: for any $t \in [0, T]$

$$I_t(X, Y) = \lim_n I_t(X_n, Y) \quad \text{and} \quad I_t^{\text{Young}}(X, Y) = \lim_n I_t^{\text{Young}}(X_n, Y).$$

By property 1 we have $I_t(X_n, Y) = I_t^{\text{Young}}(X_n, Y)$ for any n , hence

$$I_t(X, Y) = I_t^{\text{Young}}(X, Y) \quad \forall t \in [0, T],$$

which completes the proof. $\square$

Interestingly, it is also possible to characterize the Young integral as the unique continuous extension of $I^{\text{classical}}$, if we let the exponent α vary. Given $\alpha \in]0, 1[$, we define the space

$$\mathcal{C}^{>\alpha} := \bigcup_{\bar{\alpha} \in]\alpha, 1]} \mathcal{C}^{\bar{\alpha}}$$

and we agree that $f_n \rightarrow f$ in $\mathcal{C}^{>\alpha}$ if and only if $f_n \rightarrow f$ in $\mathcal{C}^{\bar{\alpha}}$ for some $\bar{\alpha} > \alpha$. The basic observation is that C^1 is dense in $\mathcal{C}^{>\alpha}$: for any $f \in \mathcal{C}^{>\alpha}$ we can find a sequence $f_n \in C^1$ such that $f_n \rightarrow f$ in $\mathcal{C}^{>\alpha}$: indeed, if $f \in \mathcal{C}^{\bar{\alpha}}$ with $\bar{\alpha} > \alpha$, by Exercise 7.2 we have $f \in \mathcal{C}_0^{\alpha'}$ for any $\alpha' \in]\alpha, \bar{\alpha}[$, then by Theorem 7.11 we can find $f_n \in C^\infty$ such that $f_n \rightarrow f$ in $\mathcal{C}^{\alpha'}$, hence $f_n \rightarrow f$ in $\mathcal{C}^{>\alpha}$.

If we fix $\alpha = 1 - \beta$, for $\beta \in]0, 1]$, the Young integral $I^{\text{Young}} = (I_t^{\text{Young}})_{t \in [0, T]}$ is a continuous map from $\mathcal{C}^{>(1-\beta)} \times \mathcal{C}^\beta$ to $\mathcal{C}^{>(1-\beta)}$, by Theorem 7.1.

These observations yield immediately the following result.

PROPOSITION 7.17. (CHARACTERIZATION OF THE YOUNG INTEGRAL, I) *Fix any $\beta \in]0, 1]$. The Young integral $I^{\text{Young}} = (I_t^{\text{Young}})_{t \in [0, T]}$, viewed as a map from $\mathcal{C}^{>(1-\beta)} \times \mathcal{C}^\beta$ to $\mathcal{C}^{>(1-\beta)}$, is the unique continuous extension of the classical integral $I^{\text{classical}} = (I_t^{\text{classical}})_{t \in [0, T]}$ defined on $C^1 \times \mathcal{C}^\beta$.*

Explicitly, I^{Young} is the unique map $I: \mathcal{C}^{>(1-\beta)} \times \mathcal{C}^\beta \rightarrow \mathcal{C}^{>(1-\beta)}$ such that:

- $I_t = I_t^{\text{classical}} = \int_0^t Y_u \dot{X}_u du$ for $X \in C^1$;
- if $(X_n, Y_n) \rightarrow (X, Y)$ in $\mathcal{C}^\alpha \times \mathcal{C}^\beta$, for some $\alpha > 1 - \beta$, then we have the convergence $I(X_n, Y_n) \rightarrow I(X, Y)$ in $\mathcal{C}^{\alpha'}$ for some $\alpha' > 1 - \beta$.

7.7. TWO TECHNICAL PROOFS

We give here the proof of Theorem 7.11 and Lemma 7.13.

Proof of Lemma 7.13. We extend $f: [0, T] \rightarrow \mathbb{R}$ to a function defined on the whole real line, by setting $f(t) = f(0)$ for $t < 0$ and $f(t) = f(T)$ for $t > T$.

Let us fix a C^∞ function $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ supported in $[-1, 1]$ with unit integral: $\int_{\mathbb{R}} \varphi(u) du = 1$. Note that $\varphi_n(t) := n\varphi(nt)$ is supported in $[-\frac{1}{n}, \frac{1}{n}]$ and also has unit integral: $\int_{\mathbb{R}} \varphi_n(u) du = 1$. We then define $f_n = \varphi_n * f$, that is

$$f_n(t) := \int_{\mathbb{R}} \varphi_n(t-u) f(u) du.$$

It is a classical result that $f_n \in C^\infty$ (we can differentiate inside the integral by dominated convergence, since f is bounded).

We next write

$$f_n(t) = \int_{\mathbb{R}} \varphi_n(u) f(t-u) du = \int_{\mathbb{R}} \varphi(v) f\left(t - \frac{v}{n}\right) dv,$$

which implies $\|f_n - f\|_\infty \leq \sup_{t \in \mathbb{R}, |u| \leq 1} \left| f\left(t - \frac{v}{n}\right) - f(t) \right|$ (since φ has unit integral), hence $\|f_n - f\|_\infty \rightarrow 0$. Property (7.18) is also directly checked. $\square$

Proof of Theorem 7.11. First we show that $\mathcal{C}_0^\alpha$ is closed in $\mathcal{C}^\alpha$: given f_n in $\mathcal{C}_0^\alpha$ and $f \in \mathcal{C}^\alpha$ such that $\|f_n - f\|_\alpha \rightarrow 0$, we need to show that $f \in \mathcal{C}_0^\alpha$, that is (7.17) holds. For $s < t$ and $n \in \mathbb{N}$ we can write, by the triangle inequality,

$$\frac{|f(t) - f(s)|}{(t-s)^\alpha} \leq \|\delta f - \delta f_n\|_\alpha + \frac{|f_n(t) - f_n(s)|}{(t-s)^\alpha}. \quad (7.21)$$

Fix $n = \bar{n}_\epsilon$ such that $\|\delta f_{\bar{n}_\epsilon} - \delta f\|_\alpha < \frac{\epsilon}{2}$. Since $f_{\bar{n}_\epsilon} \in \mathcal{C}_0^\alpha$, by (7.17) we can fix $\delta_\epsilon > 0$ such that for $|t - s| \leq \delta$ the last term in (7.21) is $\leq \frac{\epsilon}{2}$ and we are done.

It remains to show that, for any $f \in \mathcal{C}_0^\alpha$, there is a sequence $f_n \in C^\infty$ such that $\|f_n - f\|_\infty + \|\delta f_n - \delta f\|_\alpha \rightarrow 0$ (recall Remark 1.4). We define $f_n \in C^\infty$ as in Lemma 7.13, so we only need to show that $\|\delta f_n - \delta f\|_\alpha \rightarrow 0$.

Since $f \in \mathcal{C}_0^\alpha$, property (7.17) holds. The same property holds replacing for f_n , uniformly for $n \in \mathbb{N}$, thanks to relation (7.18). This means that for any $\epsilon > 0$, for all $0 \leq s < t \leq T$ with $|t - s| \leq \delta_\epsilon$, and for any $n \in \mathbb{N}$, we can write

$$\frac{|(f_n - f)(t) - (f_n - f)(s)|}{(t-s)^\alpha} \leq \frac{|f_n(t) - f_n(s)|}{(t-s)^\alpha} + \frac{|f(t) - f(s)|}{(t-s)^\alpha} \leq 2\epsilon.$$

If we fix $\bar{n}_\epsilon > 0$ such that $\|f_n - f\|_\infty \leq \epsilon (\delta_\epsilon)^\alpha$ for all $n \geq \bar{n}_\epsilon$, for $|t - s| > \delta_\epsilon$ we get

$$\frac{|(f_n - f)(t) - (f_n - f)(s)|}{(t-s)^\alpha} \leq \frac{2\|f_n - f\|_\infty}{(\delta_\epsilon)^\alpha} \leq \epsilon.$$

Altogether, the previous relations show that $\|\delta f_n - \delta f\|_\alpha \leq 2\epsilon$ for $n \geq \bar{n}_\epsilon$. This implies that $\|\delta f_n - \delta f\|_\alpha \rightarrow 0$. $\square$

CHAPTER 8

ROUGH PATHS

We have seen in Chapter 3 that it is possible to build a robust theory for a controlled equation of the form $\dot{Y}_t = \sigma(Y_t) \dot{X}_t$ with $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^\alpha$ for $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, provided we *choose* a function $\mathbb{X}^2: [0, T]^2 \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ satisfying for $0 \leq s \leq u \leq t \leq T$

$$\delta \mathbb{X}_{sut}^2 = \mathbb{X}_{su}^1 \otimes \mathbb{X}_{ut}^1, \quad |\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha},$$

see (3.13), where we denote $\mathbb{X}_{st}^1 := \delta X_{st}$, $0 \leq s \leq t \leq T$. In coordinates, the former identity means

$$(\delta \mathbb{X}^2)_{sut}^{ij} = \delta X_{su}^i \delta X_{ut}^j, \quad |(\mathbb{X}_{st}^2)^{ij}| \lesssim |t - s|^{2\alpha}, \quad i, j \in \{1, \dots, d\}. \quad (8.1)$$

In Section 3.2 we left the problem of the existence of such a function $\mathbb{X}^2$ open.

We recall that, for X of class C^1 , we have a natural choice for $\mathbb{X}^2$ given by

$$(\mathbb{X}_{st}^2)^{ij} := \int_s^t (X_r^i - X_s^i) \dot{X}_r^j dr, \quad 0 \leq s \leq t \leq T,$$

see (3.9). In Lemma 7.7 we saw that, for $\alpha > \frac{1}{2}$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$, the (uniquely defined) Young integral $I_t^{ij} := \int_0^t X^i dX^j$ satisfies

$$R_{st}^{ij} := I_t^{ij} - I_s^{ij} - X_s^i (X_t^j - X_s^j) = \int_s^t (X_r^i - X_s^i) dX_r^j, \quad |R_{st}^{ij}| \lesssim |t - s|^{2\alpha},$$

where the integral in the right-hand side is again of the Young type and $2\alpha > 1$.

There is a clear resemblance between the two last expressions, and indeed for $\alpha > \frac{1}{2}$ we show in Lemma 8.16 below that setting $(\mathbb{X}_{st}^2)^{ij} := R_{st}^{ij}$ we obtain (8.1) and this is the only possible choice.

If now $\alpha \leq \frac{1}{2}$, neither of these formulae is well-defined, because for $2\alpha \leq 1$ we are not in the setting of the Young integral. However, we have seen in Chapter 3 that the bound $|\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}$ is enough for the whole theory of existence, uniqueness and stability of the rough equation (3.19) to work, even if $2\alpha \leq 1$.

This suggests that, for every $i, j \in \{1, \dots, d\}$, the function $(\mathbb{X}_{st}^2)^{ij}$ can be interpreted as the remainder R^{ij} associated with an integral I^{ij} of (X^i, X^j) , where we *weaken* our requirements with respect to the Young integral, namely we only require that

$$I_t^{ij} - I_s^{ij} - X_s^i (X_t^j - X_s^j) = (\mathbb{X}_{st}^2)^{ij}, \quad |(\mathbb{X}_{st}^2)^{ij}| \lesssim |t - s|^{2\alpha},$$

and now $2\alpha \leq 1$. Therefore the choice of the rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ over X is equivalent to the choice of a *generalised integral* $I = \int_0^\cdot X \otimes dX \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d \otimes \mathbb{R}^d)$, and in this case $\mathbb{X}^2$ plays the role of a generalised remainder with respect to the germ $(s, t) \mapsto X_s \otimes (X_t - X_s)$.

In this chapter we explore these notions and explain them in greater detail.

8.1. INTEGRAL BEYOND YOUNG

Let us fix $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$. We saw in Theorem 7.3 that when $\alpha + \beta > 1$ we can define the integral $I_t = \int_0^t Y dX$ as the unique function which solves

$$I_0 = 0, \quad \delta I_{st} = Y_s \delta X_{st} + R_{st}, \quad R_{st} = o(t - s). \quad (8.2)$$

This was based on the observation that for the germ $A_{st} := Y_s \delta X_{st}$ we have

$$\delta A_{sut} = -\delta Y_{su} \delta X_{ut} \implies \|\delta A\|_{\alpha+\beta} \leq \|\delta X\|_\alpha \|\delta Y\|_\beta.$$

Therefore if $\eta := \alpha + \beta > 1$ we have $\|\delta A\|_\eta < \infty$, i.e. the germ A is coherent, see Definition 6.7, and the Sewing Lemma can be applied, see Theorem 6.8.

We now focus on the regime $\alpha + \beta \leq 1$. As we have already seen in (7.8) above, there exist germs A which allow *no function I solving (8.2)*. Indeed, we recall that choosing $X_t = t^\alpha$ and $Y_t = t^\beta$, $t \in [0, T]$, then the germ $A_{st} := Y_s \delta X_{st}$ satisfies $|\delta A_{0\frac{t}{2}}| \gtrsim t^{\alpha+\beta}$, see (7.8), and therefore the necessary condition (6.9) in Lemma 6.5 is not satisfied.

A solution is to relax the requirement $R_{st} = o(t - s)$ in (8.2), say to

$$\exists \eta \leq 1: \quad |R_{st}| \lesssim |t - s|^\eta. \quad (8.3)$$

Arguing as in Lemma 6.5, this would imply

$$|\delta R_{sut}| \lesssim |t - s|^\eta + |u - s|^\eta + |t - u|^\eta \lesssim |u - s|^\eta + |t - u|^\eta$$

since $\eta \leq 1$. On the other hand, by Proposition 6.4 we have $|\delta R_{sut}| = |\delta A_{sut}| \lesssim |u - s|^\beta |t - u|^\alpha$. Choosing $|u - s| = |t - u|$ shows that the best we can hope for in (8.3) is $\eta = \alpha + \beta$.

Summarizing, given $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta \leq 1$, it is natural to wonder whether there exists a function I which satisfies the following weakening of (8.2)

$$I_0 = 0, \quad \delta I_{st} = Y_s \delta X_{st} + R_{st}, \quad |R_{st}| \lesssim |t - s|^{\alpha+\beta}. \quad (8.4)$$

This would provide a “generalised notion of integral” $\int_0^\cdot Y dX$. This justifies the following

DEFINITION 8.1. *Fix $\alpha, \beta \in (0, 1)$ with $\alpha + \beta \leq 1$. Given $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$, if there exists a function $I: [0, T] \rightarrow \mathbb{R}$ which satisfies*

$$I_t - I_s = Y_s (X_t - X_s) + O(|t - s|^{\alpha+\beta}) \quad \text{uniformly as } |t - s| \rightarrow 0, \quad (8.5)$$

we say that I is a generalised integral of Y in dX .

We stress that this new definition of integral extends the previous one (8.2) for $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta > 1$, because the term $o(t - s)$ is actually $O(|t - s|^{\alpha+\beta})$ in this case, by the key estimate for the Young integral (or, equivalently, for the sewing map).

On the positive side, *there is always existence for (8.4) if $\alpha + \beta < 1$* . This is a non-trivial result, due (in a more general setting) to Lyons and Victoir. We state this as a separate result, which is a consequence of Proposition 8.5 below.

LEMMA 8.2. *Let $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta < 1$. There exists $(I, R) \in \mathcal{C}^\alpha \times C_2^{\alpha+\beta}$ satisfying (8.4).*

Remark 8.3. It is an easy observation that *uniqueness can not hold for (8.4)*. Indeed, given I which solves (8.4), any function of the form $I'_t := I_t + h_t - h_0$ with $h \in \mathcal{C}^{\alpha+\beta}$ still solves (8.4). As a matter of fact, *all solutions are of this form*, because given two solutions I, I' of (8.4), with corresponding R, R' , their difference $h := I' - I$ must satisfy $|\delta h_{st}| = |R'_{st} - R_{st}| \lesssim |t - s|^{\alpha+\beta}$.

Remark 8.4. An integral I as in Definition 8.1 is necessarily of class $\mathcal{C}^\alpha$ by (8.5).

We state now a result which implies Lemma 8.2 above.

PROPOSITION 8.5. (PARAINTEGRAL) *Fix $\alpha, \beta \in (0, 1)$ with $\alpha + \beta < 1$. There exists a (non unique) bilinear and continuous map $J_{\prec}: \mathcal{C}^\alpha \times \mathcal{C}^\beta \rightarrow \mathcal{C}_2^{\alpha+\beta}$ such that*

$$\|J_{\prec}(X, Y)\|_{\alpha+\beta} \leq C \| \delta X \|_{\alpha} \| \delta Y \|_{\beta}, \quad (8.6)$$

for a suitable $C = C(\alpha, \beta, T)$, with the property that, for all $s < u < t$,

$$\delta J_{\prec}(X, Y)_{sut} = \delta Y_{su} \delta X_{ut}. \quad (8.7)$$

The proof of Proposition 8.5 is postponed to Section 8.9 below.

Remark 8.6. Let $\alpha, \beta \in (0, 1)$ with $\alpha + \beta \leq 1$. Finding a generalised integral of Y in dX for $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ as in Definition 8.1 is equivalent to finding $R_{st} \in \mathcal{C}_2^{\alpha+\beta}$ such that

$$\delta R_{sut} = \delta Y_{su} \delta X_{ut}, \quad (8.8)$$

$$R \in \mathcal{C}_2^{\alpha+\beta}. \quad (8.9)$$

Indeed, if we define $A_{st} := Y_s \delta X_{st}$, relation (8.8) implies that $\delta(A + R) = 0$, hence there exists $I: [0, T] \rightarrow \mathbb{R}$ which satisfies $\delta I = A + R$, which is exactly relation (8.5).

By Proposition 8.5 and Remark 8.6, if $\alpha, \beta \in (0, 1)$ and $\alpha + \beta < 1$, any $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ admits an integral I as in Definition 8.1.

Remark 8.7. (THE ITÔ INTEGRAL AS A GENERALISED INTEGRAL) Fix a standard Brownian motion $(B_t)_{t \in [0, T]}$ and an adapted process $(h_t)_{t \in [0, T]}$ such that a.s. h is of class $\mathcal{C}^\beta$ with $\beta \in (0, 1)$. If $\alpha \in (0, \frac{1}{2})$ and $\alpha + \beta \leq 1$, then by (4.8) in Theorem 4.3 we have the a.s. estimate on the Itô integral

$$\left| \int_s^t (h_r - h_s) dB_r \right| \lesssim |t - s|^{\alpha+\beta}, \quad 0 \leq s \leq t \leq T.$$

Therefore in this setting the Itô integral $(\int_0^t h_r dB_r)_{t \in [0, T]}$ is a.s. a generalised integral of h in dB in the sense of Definition 8.1. Compare this with the case $\alpha + \beta > 1$ discussed in Proposition 7.4.

8.2. A NEGATIVE RESULT

We show that the usual integral $I(f, g) = \int_0^t g_s f'_s ds$, when $f \in C^1$, cannot be extended to a continuous operator on $\mathcal{C}^\alpha \times \mathcal{C}^\beta$, when $\alpha + \beta < 1$.

LEMMA 8.8. Set $[0, T] = [0, 1]$ and define, for $\alpha, \beta \in (0, 1)$,

$$g_n(t) := \frac{1}{n^\alpha} \cos(nt), \quad f_n(t) := \frac{1}{n^\beta} \sin(nt).$$

Then $g_n \rightsquigarrow_\alpha 0$ and $f_n \rightsquigarrow_\beta 0$ (recall Definition 7.15), more precisely:

$$\|g_n\|_\infty \rightarrow 0, \quad \|\delta g_n\|_\alpha \leq 2; \quad \|f_n\|_\infty \rightarrow 0, \quad \|\delta f_n\|_\beta \leq 2, \quad (8.10)$$

(in particular, $g_n \rightarrow 0$ in $\mathcal{C}^{\alpha'}$ and $f_n \rightarrow 0$ in $\mathcal{C}^{\beta'}$ for any $\alpha' < \alpha$ and $\beta' < \beta$).

However, if we fix $\alpha + \beta \leq 1$, we have $I(f_n, g_n) \not\rightarrow 0$, because

$$\forall t \in [0, 1]: \quad \lim_{n \rightarrow \infty} I(f_n, g_n)_t = \begin{cases} +\infty & \text{if } \alpha + \beta < 1 \\ \frac{1}{2}t & \text{if } \alpha + \beta = 1 \\ 0 & \text{if } \alpha + \beta > 1 \end{cases}.$$

Proof. Note that $\|g_n\|_\infty = n^{-\alpha}$ and $\|g'_n\|_\infty = n^{1-\alpha}$, hence

$$|g_n(t) - g_n(s)| \leq \min \{ \|g'_n\|_\infty |t - s|, 2 \|g_n\|_\infty \} \leq \min \{ n^{1-\alpha} |t - s|, 2 n^{-\alpha} \}.$$

Since $\min \{x, y\} \leq x^\gamma y^{1-\gamma}$, for any $\gamma \in [0, 1]$, choosing $\gamma = \alpha$ we obtain

$$|g_n(t) - g_n(s)| \leq 2^{1-\alpha} |t - s|^\alpha,$$

hence $\|\delta g_n\|_\alpha \leq 2^{1-\alpha} \leq 2$. Similar arguments apply to f_n , proving (8.10).

Next we observe that $\frac{1}{2\pi} \int_0^{2\pi} \cos^2(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \sin^2(x) dx = \frac{1}{2}$. Then, for fixed $t > 0$, as $n \rightarrow \infty$

$$\int_0^{nt} \cos^2(x) dx = \int_0^{2\pi \lfloor \frac{nt}{2\pi} \rfloor} \cos^2(x) dx + O(1) = \frac{1}{2} 2\pi \left\lfloor \frac{nt}{2\pi} \right\rfloor + O(1) = \frac{t}{2} n + O(1).$$

It follows that

$$I(f_n, g_n)_t = \frac{n}{n^{\alpha+\beta}} \int_0^t \cos^2(ns) ds = \frac{1}{n^{\alpha+\beta}} \int_0^{nt} \cos^2(x) dx \sim \frac{t}{2} n^{1-(\alpha+\beta)}. \quad \square$$

8.3. A CHOICE

We have seen in (7.11) above that, given $(X, Y) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta > 1$, we have an *explicit formula* for the remainder $R_{st} = I_t - I_s - Y_s(X_t - X_s)$, given by

$$R_{st} = \int_s^t (Y_u - Y_s) dX_u, \quad 0 \leq s \leq t \leq T, \quad (8.11)$$

where $I_t = \int_0^t Y_u dX_u$ is the *unique* function given by the Young integral of Theorem 7.1. Moreover $R_{st} = \int_s^t (Y_u - Y_s) dX_u$ is the unique function in C_2 which satisfies

$$R \in C_2^{\alpha+\beta}, \quad \delta R_{sut} = \delta Y_{su} \delta X_{ut}, \quad 0 \leq s \leq u \leq t \leq T. \quad (8.12)$$

In the regime $\alpha + \beta < 1$, the Young integral is not available anymore. However by Proposition 8.5 we know that we can find an integral $I \in \mathcal{C}^\alpha$ in the sense of Definition 8.1 by setting

$$\delta I_{st} := Y_s(X_t - X_s) - J_{\prec}(X, Y)_{st},$$

where $J_{\prec}$ is the paraintegral of Proposition 8.5, see also Remark 8.6. This shows that, in this setting, the remainder $R_{st} = I_t - I_s - Y_s(X_t - X_s)$ is not given by an explicit formula like (8.11) (which is now ill-defined), rather we have

$$R = -J_{\prec}(X, Y).$$

However formula (8.11) suggests that we can *define*

$$\int_s^t (Y_u - Y_s) dX_u := R_{st} = -J_{\prec}(X, Y)_{st}, \quad 0 \leq s \leq t \leq T. \quad (8.13)$$

In other words, the left hand side of (8.13) is *chosen* to be equal to the remainder R associated with the integral I as in (8.4). We recall that $R = -J_{\prec}(X, Y)$ satisfies

$$R \in C_2^{\alpha+\beta}, \quad \delta R_{sut} = \delta Y_{su} \delta X_{ut}, \quad 0 \leq s \leq u \leq t \leq T. \quad (8.14)$$

The difference between formula (8.14) and formula (8.12), is that in the former $\alpha + \beta < 1$ while in the latter $\alpha + \beta > 1$. Accordingly, in (8.14) the function R is *not* uniquely determined, while in (8.12) it is.

The comparison between formula (8.14) and formula (8.12), and the explicit expression (8.11) in the case $\alpha + \beta > 1$ show that (8.13) is a reasonable *definition* of the function $(s, t) \mapsto \int_s^t (Y_u - Y_s) dX_u$ in the setting $\alpha + \beta \leq 1$.

We also stress that R in (8.14) *can not be uniquely determined*. Indeed, by Remark 8.3, we have infinitely many possible choices given by

$$R' = R + \delta h, \quad h \in \mathcal{C}^{\alpha+\beta}, h_0 = 0. \quad (8.15)$$

Remark 8.9. In the special case $X = Y$ and $\alpha = \beta \leq \frac{1}{2}$, (8.4) becomes

$$I_0 = 0, \quad \delta I_{st} = X_s \delta X_{st} + R_{st}, \quad |R_{st}| \lesssim |t - s|^{2\alpha}. \quad (8.16)$$

Now the germ is $A_{st} = X_s(X_t - X_s)$ and we have a simple canonical solution which does not rely on the paraintegral and is given by

$$I_t := \frac{1}{2}(X_t^2 - X_0^2), \quad R_{st} := \frac{1}{2}(X_t - X_s)^2,$$

since

$$\underbrace{\frac{1}{2}(X_t^2 - X_s^2)}_{I_t - I_s} = \underbrace{X_s(X_t - X_s)}_{A_{st}} + \underbrace{\frac{1}{2}(X_t - X_s)^2}_{R_{st}}.$$

As we have seen in (7.15)-(7.16), if $\alpha > \frac{1}{2}$ then (I, R) is the *only* solution of (8.16) and moreover

$$R_{st} = \int_s^t (X_r - X_s) dX_r$$

where the integral is in the Young sense. If $\alpha \leq \frac{1}{2}$, then we have infinitely many possible solutions (I', R') .

8.4. ONE-DIMENSIONAL ROUGH PATHS

We have seen at the beginning of this chapter that for every $i, j \in \{1, \dots, d\}$, the function $(\mathbb{X}_{st}^2)^{ij}$ plays the role of the remainder R^{ij} associated with a generalised integral I^{ij} of (X^i, X^j) in the sense of Definition 8.1 with $\alpha = \beta < \frac{1}{2}$: in other words the choice of $\mathbb{X}^2$ is *equivalent* to the choice of integrals (in the sense of Definition 8.1) $I^{ij} \in \mathcal{C}^\alpha$ for all $i, j \in \{1, \dots, d\}$, such that

$$I_0^{ij} = 0, \quad \delta I_{st}^{ij} = X_s^i \delta X_{st}^j + (\mathbb{X}_{st}^2)^{ij}, \quad |(\mathbb{X}_{st}^2)^{ij}| \lesssim |t - s|^{2\alpha},$$

or, in more compact notations,

$$I_0 = 0, \quad \delta I_{st} = X_s \otimes \mathbb{X}_{st}^1 + \mathbb{X}_{st}^2, \quad |\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}. \quad (8.17)$$

Existence of $\mathbb{X}^2$ satisfying (8.17) with $\alpha < \frac{1}{2}$ is therefore granted by Lemma 8.2, e.g. via the paraintegral of Theorem 8.5. We also know that in the regime $\alpha < \frac{1}{2}$ we have infinitely many possible choices for $(I, \mathbb{X}^2)$, all of the form (8.15) above.

Suppose first that we are in the setting $d = 1$. Then Definition 3.2 becomes

DEFINITION 8.10. *Let $\alpha \in [\frac{1}{3}, \frac{1}{2}]$ and $X: [0, T] \rightarrow \mathbb{R}$ of class $\mathcal{C}^\alpha$. A α -Rough Path over X is a pair $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2) \in C_2^\alpha \times C_2^{2\alpha}$ such that*

$$\mathbb{X}_{st}^1 = X_t - X_s, \quad \delta \mathbb{X}_{sut}^2 = \mathbb{X}_{su}^1 \mathbb{X}_{ut}^1. \quad (8.18)$$

We recall that the conditions $X \in \mathcal{C}^\alpha$ and $\mathbb{X}^1 = \delta X \in C_2^\alpha$ are equivalent, and that $(\mathbb{X}^1, \mathbb{X}^2) \in C_2^\alpha \times C_2^{2\alpha}$ is equivalent to

$$|\mathbb{X}_{st}^1| \lesssim |t - s|^\alpha, \quad |\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}.$$

We have seen in Chapter 3 that it is possible to build an integration theory for every choice of the α -rough path $\mathbb{X}$ over X . In this theory we can recover existence *and* uniqueness of the integral function $\int_0^\cdot Y \, dX$ for a large class of choices of Y . For this we have to give very different roles to the integrator X and to the integrand Y , whereas in the case of the Young integral the two functions play a symmetric role: X will be a component of a rough path and Y a component of a *controlled path*, see Chapter 9.

We note that the algebraic condition $\delta \mathbb{X}_{sut}^2 = \mathbb{X}_{su}^1 \mathbb{X}_{ut}^1$ is *non-linear*, which implies that α -rough paths do not form a vector subspace of $C_2^\alpha \times C_2^{2\alpha}$.

For all $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, given any *real-valued* path $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R})$, there is always a rough path lying above X . Indeed, $I_t := \frac{1}{2} X_t^2$ is a generalised integral of X in dX integral in the sense of Definition 8.1, because

$$\delta I_{st} = \frac{1}{2} (X_t^2 - X_s^2) = X_s \delta X_{st} + \frac{1}{2} (\delta X_{st})^2 = X_s \delta X_{st} + O(|t - s|^{2\alpha}).$$

Then, by Remark 8.9, we can define a rough path $\mathbb{X}$ by setting

$$\mathbb{X}_{st}^2 = \frac{1}{2} (\delta X_{st})^2. \quad (8.19)$$

More directly, note that (8.19) satisfies the Chen relation (8.21), and clearly $\mathbb{X}^2 \in C_2^{2\alpha}$.

8.5. THE VECTOR CASE

Let us consider now a *vector valued* path $X: [0, T] \rightarrow \mathbb{R}^d$, with $X_t = (X_t^1, \dots, X_t^d)$. We suppose that X is of class $\mathcal{C}^\alpha$, namely that $X^i \in \mathcal{C}^\alpha$ for all $i = 1, \dots, d$, with $\alpha > \frac{1}{3}$.

We can now see that Definition 3.2 generalises Definition 8.10 to the vector case. The multi-dimensional case $d \geq 2$ is sensibly richer, because off-diagonal terms $I_t^{ij} = \int_0^t X^i dX^j$ with $i \neq j$ as in (8.17) do not have explicit candidates as in (8.19). We rephrase then Definition 3.2 as follows:

DEFINITION 8.11. *Let $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, $d \geq 1$ and $X: [0, T] \rightarrow \mathbb{R}^d$ of class $\mathcal{C}^\alpha$. An α -Rough Path on $\mathbb{R}^d$ over X is a pair $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$, with*

- $\mathbb{X}^1 = (\delta X^i)_{i=1, \dots, d} \in C_2^\alpha([0, T]; \mathbb{R}^d)$
- $\mathbb{X}^2 = (R^{ij})_{i, j=1, \dots, d} \in C_2^{2\alpha}([0, T]_{\leq}^2; \mathbb{R}^d \otimes \mathbb{R}^d)$

such that

$$(\delta \mathbb{X}_{sut}^2)^{ij} = (\mathbb{X}_{su}^1)^i (\mathbb{X}_{ut}^1)^j, \quad (8.20)$$

or equivalently

$$\mathbb{X}_{st}^2 - \mathbb{X}_{su}^2 - \mathbb{X}_{ut}^2 = \mathbb{X}_{su}^1 \otimes \mathbb{X}_{ut}^1. \quad (8.21)$$

We denote by $\mathcal{R}_{\alpha, d}$ the space of α -rough paths on $\mathbb{R}^d$ and by $\mathcal{R}_{\alpha, d}(X)$ the set of α -rough paths over X .

The condition (8.20)-(8.21) is the celebrated *Chen relation* (3.13). Via the equality (8.17), a choice of $\mathbb{X} \in \mathcal{R}_{\alpha, d}(X)$ is equivalent to the choice of $I: [0, T] \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ such that $I_0 = 0$ and

$$|I_t - I_s - X_s \otimes (X_t - X_s)| \lesssim |t - s|^{2\alpha}, \quad 0 \leq s \leq t \leq T,$$

see Exercise 8.1 below. We say that $I = \int_0^\cdot X_s \otimes dX_s$ and we know that in the regime $\alpha < \frac{1}{2}$ we have infinitely many possible choices for $(I, \mathbb{X}^2)$, all of the form (8.15) above.

In other words, for every fixed $x_0 \in \mathbb{R}^d$, we have a natural bijection between the space $\mathcal{R}_{\alpha, d}$ of α -rough paths on $\mathbb{R}^d$ and the pairs (X, I) where $X: [0, T] \rightarrow \mathbb{R}^d$ is of class $\mathcal{C}^\alpha$ with $X_0 = x_0$, and $I: [0, T] \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ satisfying (8.17).

We are going to see in Chapter 9 that it is possible to build an integration theory for every choice of an α -rough path $\mathbb{X}$. Again, we note that the condition (8.20)-(8.21) is *non-linear*, which implies that α -rough paths do not form a vector space.

Exercise 8.1. Given a α -rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ over X in $\mathbb{R}^d$, a process $I \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d \otimes \mathbb{R}^d)$ satisfying (8.17) is a generalised integral of X in dX in the sense of Definition 8.1.

Viceversa, given $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ and an integral $I \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d \otimes \mathbb{R}^d)$ of X in dX , in the sense of Definition 8.1, defining $\mathbb{X}^2$ by (8.17) we obtain a α -rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ over X in $\mathbb{R}^d$.

In the multi-dimensional case $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ with $d \geq 2$, building a rough path over X is non-trivial, because one has to define off-diagonal integrals $\int X^i dX^j$ for $i \neq j$. However, by the results we have proved on the existence of the paraintegral in Proposition 8.5, we can easily deduce the following.

PROPOSITION 8.12. *For any $d \in \mathbb{N}$, $\alpha \in (\frac{1}{3}, \frac{1}{2})$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$, there is a α -rough path $\mathbb{X}$ which lies above X (hence, by Lemma 8.17, there are infinitely many of them).*

Proof. For any fixed $i, j \in \{1, \dots, d\}$, let I^{ij} be a generalised integral of X^i in dX^j in the sense of Definition 8.1, whose existence is guaranteed by the paraintegral of Proposition 8.5. Then, by Exercise 8.1, defining $\mathbb{X}^2$ by (8.17) we obtain a rough path $\mathbb{X}$ which lies above X . $\square$

We conclude with an elementary observation, that will be useful later. By Exercise 8.1, any α -rough path $\mathbb{X}$ over $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ determines an integral I of (X, X) , given by (8.17). Applying the latter relation in a telescopic fashion, we can write

$$I_t = \sum_{[t_i, t_{i+1}] \in \mathcal{P}} (X_{t_i} \delta X_{t_i t_{i+1}} + \mathbb{X}_{t_i t_{i+1}}^2), \quad (8.22)$$

where $\mathcal{P} = \{0 = t_0 < t_1 < \dots < t_k = t\}$ is an arbitrary partition of $[0, t]$. We will see in Chapter 9 below that a generalization of (8.22), when we also take the limit of vanishing mesh $|\mathcal{P}| \rightarrow 0$, is the correct recipe for building “Riemann-sums”, in order to define a generalised integral of h in dX in the sense of Definition 8.1 for a wide class of functions h .

Remark 8.13. (ITÔ AND STRATONOVICH) Given a d -dimensional Brownian motion $(B_t)_{t \in [0, T]}$, with $B = (B^1, \dots, B^d)$ a vector of d independent standard BMs, we have seen in Theorem 4.1 and in Remark 5.1 that the Itô rough paths $\mathbb{B} = (\mathbb{B}^1, \mathbb{B}^2)$ defined in (4.2) and the Stratonovich rough path $\bar{\mathbb{B}} = (\bar{\mathbb{B}}^1, \bar{\mathbb{B}}^2)$ defined in (5.1) satisfy Definition 8.11 for all $\alpha \in (\frac{1}{3}, \frac{1}{2})$. We recall that this is based on the a.s. estimate on the Itô integral

$$\left| \int_s^t (B_r^i - B_s^i) dB_r^j \right| \lesssim |t - s|^{2\alpha}, \quad 0 \leq s \leq t \leq T,$$

which follows from (4.8) in Theorem 4.3. This is a special case of Remark 8.7.

8.6. DISTANCE ON ROUGH PATHS

We denote by $\mathcal{R}_{\alpha, d}$ the set of all α -rough paths in $\mathbb{R}^d$. For $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha, d}$ we set

$$\|\mathbb{X}\|_{\mathcal{R}_{\alpha, d}} := \|\mathbb{X}^1\|_\alpha + \|\mathbb{X}^2\|_{2\alpha} = \sup_{0 \leq s < t \leq T} \frac{|\mathbb{X}_{st}^1|}{|t - s|^\alpha} + \sup_{0 \leq s < t \leq T} \frac{|\mathbb{X}_{st}^2|}{|t - s|^{2\alpha}}. \quad (8.23)$$

We stress that $\mathcal{R}_{\alpha, d}$ is not a vector space, because the Chen relation (8.21) is not linear. However, it is meaningful to define for $\mathbb{X}, \bar{\mathbb{X}} \in \mathcal{R}_{\alpha, d}$

$$d_{\mathcal{R}_{\alpha, d}}(\mathbb{X}, \bar{\mathbb{X}}) := \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha + \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}. \quad (8.24)$$

Exercise 8.2. $d_{\mathcal{R}_{\alpha, d}}$ is a distance on $\mathcal{R}_{\alpha, d}$.

When we talk of convergence in $\mathcal{R}_{\alpha, d}$, we mean with respect to the distance $d_{\mathcal{R}_{\alpha, d}}$. Note that $d_{\mathcal{R}_{\alpha, d}}$ is equal on $\mathcal{R}_{\alpha, d}$ to the distance induced by the natural norm $\|F\|_\alpha + \|G\|_{2\alpha}$ for $(F, G) \in C_2^\alpha \times C_2^{2\alpha}$. In particular $\mathbb{X}_n = (\mathbb{X}_n^1, \mathbb{X}_n^2) \rightarrow \mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ in $\mathcal{R}_{\alpha, d}$ if and only if $\mathbb{X}_n^1 \rightarrow \mathbb{X}^1$ in C_2^α and $\mathbb{X}_n^2 \rightarrow \mathbb{X}^2$ in $C_2^{2\alpha}$.

LEMMA 8.14. *The metric space $(\mathcal{R}_{\alpha,d}, d_{\mathcal{R}_{\alpha,d}})$ is complete.*

Proof. Let $(\mathbb{X}_n)_{n \in \mathbb{N}} \subset \mathcal{R}_{\alpha,d}$ be a Cauchy sequence. Then, by definition of $d_{\mathcal{R}_{\alpha,d}}$, for every $\epsilon > 0$ there is $\bar{n}_\epsilon < \infty$ such that for all $n, m \geq \bar{n}_\epsilon$ and $0 \leq s < t \leq T$

$$|\mathbb{X}_n^1(s, t) - \mathbb{X}_m^1(s, t)| \leq \epsilon |t - s|^\alpha, \quad |\mathbb{X}_n^2(s, t) - \mathbb{X}_m^2(s, t)| \leq \epsilon |t - s|^{2\alpha}. \quad (8.25)$$

Note that

$$d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}}) \geq \frac{\|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\infty}{T^\alpha} + \frac{\|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_\infty}{T^{2\alpha}}.$$

It follows that the sequences of continuous functions $(\mathbb{X}_n^1)_{n \in \mathbb{N}}$ and $(\mathbb{X}_n^2)_{n \in \mathbb{N}}$ are Cauchy in the sup-norm, hence there are continuous functions $\mathbb{X}^1$ and $\mathbb{X}^2$ such that $\|\mathbb{X}_n^1 - \mathbb{X}^1\|_\infty \rightarrow 0$ and $\|\mathbb{X}_n^2 - \mathbb{X}^2\|_\infty \rightarrow 0$. In particular, we have pointwise convergence $\mathbb{X}_m^1(s, t) \rightarrow \mathbb{X}^1(s, t)$ and $\mathbb{X}_m^2(s, t) \rightarrow \mathbb{X}^2(s, t)$ as $m \rightarrow \infty$. Taking this limit in (8.25) shows that $d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}_n, \mathbb{X}) \leq 2\epsilon$ for all $n \geq \bar{n}_\epsilon$. $\square$

This allows to rephrase the continuity result of section 3.7. We fix

$$D \geq \|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty + \|\nabla^3 \sigma\|_\infty + \|\nabla \sigma_2\|_\infty + \|\nabla^2 \sigma_2\|_\infty.$$

We obtain from Proposition 3.11

PROPOSITION 8.15. *We suppose that $\alpha \in (\frac{1}{3}, \frac{1}{2}]$ and $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is of class C^3 , with $\|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty + \|\nabla^3 \sigma\|_\infty + \|\nabla \sigma_2\|_\infty + \|\nabla^2 \sigma_2\|_\infty < +\infty$ (without boundedness assumptions on σ and σ_2). For $\mathbb{X} \in \mathcal{R}_{\alpha,d}$ and $Z_0 \in \mathbb{R}^k$ we denote by $Z: [0, T] \rightarrow \mathbb{R}^k$ the unique solution to equation (3.19)*

$$Z_{st}^{[3]} = o(t - s), \quad Z_{st}^{[3]} = \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma_2(Z_s) \mathbb{X}_{st}^2,$$

Then the map $\mathbb{R}^k \times \mathcal{R}_{\alpha,d} \ni (Z_0, \mathbb{X}) \mapsto Z \in \mathcal{C}^\alpha$ is locally Lipschitz continuous.

Let us fix $x_0 \in \mathbb{R}^d$. Via the bijection constructed in the discussion after Definition 8.11, we have that $\mathbb{X}$ and $\bar{\mathbb{X}}$ determine $(X, I), (\bar{X}, \bar{I}): [0, T] \rightarrow \mathbb{R}^d \times (\mathbb{R}^d \otimes \mathbb{R}^d)$ such that $X_0 = \bar{X}_0 = x_0$, $I_0 = \bar{I}_0 = 0$, $\delta X := \mathbb{X}^1$, $\delta \bar{X} := \bar{\mathbb{X}}^1$, and

$$|R_{st}| + |\bar{R}_{st}| \lesssim |t - s|^{2\alpha}, \quad 0 \leq s \leq t \leq T,$$

where $R_{st} := I_t - I_s - X_s \otimes (X_t - X_s)$ and $\bar{R}_{st} := \bar{I}_t - \bar{I}_s - \bar{X}_s \otimes (\bar{X}_t - \bar{X}_s)$. Now, the distance between $\mathbb{X}$ and $\bar{\mathbb{X}}$ in $\mathcal{R}_{\alpha,d}$ is equal to

$$d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}}) = \|\delta X - \delta \bar{X}\|_\alpha + \|R - \bar{R}\|_{2\alpha}.$$

8.7. CANONICAL ROUGH PATHS FOR $\alpha > \frac{1}{2}$

Let $\frac{1}{3} < \alpha' \leq \frac{1}{2} < \alpha < 1$. Then it is well known that $\mathcal{C}^\alpha \subset \mathcal{C}^{\alpha'}$. Therefore, if $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ we have in particular $X \in \mathcal{C}^{\alpha'}([0, T]; \mathbb{R}^d)$ and therefore there is a α' -rough path $\mathbb{X}$ over X . However, is there a α -rough path over X ? Note that we have restricted Definition 8.11 to the range $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, while here we are discussing the existence of $\mathbb{X}^2: [0, T] \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ satisfying the Chen relation (8.21) and

$$|\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}$$

where now $\alpha > \frac{1}{2}$.

LEMMA 8.16. *Let $\alpha \in (\frac{1}{2}, 1]$. For every $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$, there is a unique $\mathbb{X}^2: [0, T]_{\leq}^2 \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ satisfying the Chen relation (8.21) and such that $\mathbb{X}^2 \in C_2^{2\alpha}$. We have the explicit formula*

$$\mathbb{X}_{st}^2 = \int_s^t \mathbb{X}_{su}^1 \otimes dX_u, \quad \mathbb{X}_{st}^1 = \delta X_{st}, \quad 0 \leq s \leq t \leq T, \quad (8.26)$$

where the integral is in the Young sense. Moreover the map $\mathcal{C}^\alpha \ni X \mapsto \mathbb{X}^2 \in C_2^{2\alpha}$ is continuous (in particular, locally Lipschitz-continuous).

Proof. It is easy to check that $\mathbb{X}^2$ in (8.26) satisfies the Chen relation (8.18), thanks to the bi-linearity of the Young integral. Indeed, we can rewrite (8.26) as

$$\mathbb{X}_{st}^2 = \int_s^t X_u \otimes dX_u - X_s \otimes (X_t - X_s), \quad (8.27)$$

hence for $s \leq u \leq t$ we have that

$$\begin{aligned} (\delta \mathbb{X}^2)_{sut} &= -X_s \otimes (X_t - X_s) + X_s \otimes (X_u - X_s) + X_u \otimes (X_t - X_u) \\ &= -X_s \otimes (X_t - X_u) + X_u \otimes (X_t - X_u) \\ &= \delta X_{su} \otimes \delta X_{ut}. \end{aligned}$$

We show now that $\mathbb{X}^2 \in C_2^{2\alpha}$. We recall that the Young integral satisfies the following key estimate, for $f \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$ with $\alpha + \beta > 1$:

$$\left| \int_s^t f dg - f_s (g_t - g_s) \right| \leq c_{\alpha+\beta} |t - s|^{\alpha+\beta}.$$

Choosing $f = X^i$ and $g = X^j$ shows that $\mathbb{X}^2$, given by (8.27), is $O(|t - s|^{2\alpha})$. Finally, we prove the continuity of $\mathcal{C}^\alpha \ni X \mapsto \mathbb{X}^2 \in C_2^{2\alpha}$. Given $X, \bar{X} \in \mathcal{C}^\alpha$ and the respective $\mathbb{X}^2, \bar{\mathbb{X}}^2 \in C_2^{2\alpha}$, we have

$$\mathbb{X}_{st}^2 - \bar{\mathbb{X}}_{st}^2 = \int_s^t (\mathbb{X}_{su}^1 - \bar{\mathbb{X}}_{su}^1) \otimes dX_u + \int_s^t \bar{\mathbb{X}}_{su}^1 \otimes d(X - \bar{X})_u,$$

with all integrals in the Young sense. Then by the Sewing Lemma

$$\|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} \leq K_{2\alpha} (\|\delta X\|_\alpha + \|\delta \bar{X}\|_\alpha) \|\delta X - \delta \bar{X}\|_\alpha.$$

The proof is complete. $\square$

Therefore, we could extend Definition 8.11 to α -rough paths for $\alpha \in (\frac{1}{3}, 1]$. For $\alpha \in (\frac{1}{2}, 1]$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ there is a unique α -rough path over X , which we call the *canonical rough path* over X .

While for $\alpha > \frac{1}{2}$ there is a unique rough path lying above a given path $X \in \mathcal{C}^\alpha$, for $\alpha \leq \frac{1}{2}$ there are infinitely many of them, that can be characterized explicitly.

LEMMA 8.17. *Let $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ be a α -rough path in $\mathbb{R}^d$, with $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. Then $\bar{\mathbb{X}} = (\bar{\mathbb{X}}^1, \bar{\mathbb{X}}^2)$ is a α -rough path if and only if for some $f \in \mathcal{C}^{2\alpha}([0, T]; \mathbb{R}^d \otimes \mathbb{R}^d)$ one has $\bar{\mathbb{X}}^2 = \mathbb{X}^2 + \delta f$, that is*

$$\bar{\mathbb{X}}_{st}^2 = \mathbb{X}_{st}^2 + f_t - f_s, \quad 0 \leq s \leq t \leq T.$$

Proof. By assumption $\mathbb{X}^2$ and $\bar{\mathbb{X}}^2$ satisfy the Chen relation (8.21). If $\bar{\mathbb{X}}^2 = \mathbb{X}^2 + \delta f$ then $\mathbb{X}^2 \in C_2^{2\alpha}$ if and only if $\delta \mathbb{X}^2 = \delta \bar{\mathbb{X}}^2$ and $\bar{\mathbb{X}}^2 \in C_2^{2\alpha}$. Therefore, if $\mathbb{X}$ is a α -rough path then so is $\bar{\mathbb{X}}$.

Viceversa, if $\bar{\mathbb{X}}$ is a α -rough path, then $\delta \mathbb{X}^2 = \delta \bar{\mathbb{X}}^2$ because both $\mathbb{X}$ and $\bar{\mathbb{X}}$ satisfy the Chen relation (8.21) with the same $\mathbb{X}^1$, hence $\bar{\mathbb{X}}^2 = \mathbb{X}^2 + \delta f$ for some f . Since both $\mathbb{X}^2, \bar{\mathbb{X}}^2$ belong to $C_2^{2\alpha}$, then also $\delta f \in C_2^{2\alpha}$, which is the same as $f \in C^{2\alpha}$. $\square$

Remark 8.18. We mainly work with α -Hölder rough paths for $\alpha \in (\frac{1}{3}, \frac{1}{2})$, excluding the boundary case $\alpha = \frac{1}{2}$ for technical reasons. Let us stress that, by doing so, *we are not throwing away any rough paths, but only giving up a tiny amount of regularity*, because any $\frac{1}{2}$ -rough path is a α -rough path, for any $\alpha < \frac{1}{2}$.

To summarize, the situation is the following:

1. For $\alpha \in (\frac{1}{2}, 1]$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ there is a unique α -rough path over X
2. For $\alpha \in (\frac{1}{3}, \frac{1}{2})$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$, there are infinitely many α -rough paths over X
3. For $\alpha = \frac{1}{2}$, either there is no α -rough path over X , or there are infinitely many of them.

In the range $\alpha \in (\frac{1}{2}, 1]$, the unique α -rough path $\mathbb{X}$ above X can be called the *canonical rough path* over X . We let $\mathcal{R}_{1,d}$ be the set of all canonical rough paths over paths $X \in C^1$ (see Lemma 8.16).

8.8. LACK OF CONTINUITY

We have seen in Lemma 8.16 that, for $\alpha > \frac{1}{2}$, the map $\mathcal{C}^\alpha \ni X \mapsto \mathbb{X}^2 \in C_2^{2\alpha}$ is continuous if $\mathbb{X}^2$ is defined by the Young integral (8.26). It is a crucial fact that this continuity property can *not* be extended to $\alpha \leq \frac{1}{2}$, as shown by the next example.

For $n \in \mathbb{N}$ consider the smooth paths $X_n^1, X_n^2: [0, 1] \rightarrow \mathbb{R}$

$$X_n^1(t) := \frac{1}{\sqrt{n}} \cos(nt), \quad X_n^2(t) := \frac{1}{\sqrt{n}} \sin(nt).$$

We have already shown in Lemma 8.8 that $X_n^1 \rightarrow 0$ and $X_n^2 \rightarrow 0$ in $\mathcal{C}^\alpha$, for all $\alpha \in (0, \frac{1}{2})$. More precisely, we have shown that $X_n^1 \rightsquigarrow_{\frac{1}{2}} 0$ and $X_n^2 \rightsquigarrow_{\frac{1}{2}} 0$, by showing that $\|\delta X_n^1\|_{\frac{1}{2}} \leq 2$, $\|\delta X_n^2\|_{\frac{1}{2}} \leq 2$ for all $n \in \mathbb{N}$ and, obviously, $\|X_n^1\|_\infty \rightarrow 0$, $\|X_n^2\|_\infty \rightarrow 0$. Next we set

$$I_n^{ij}(t) := \int_0^t X_n^i(u) dX_n^j(u), \quad \text{for } i, j \in \{1, 2\},$$

and correspondingly $(\mathbb{X}_n^1)_{st} := (X_n^1(t) - X_n^1(s), X_n^2(t) - X_n^2(s))$,

$$\begin{aligned} (\mathbb{X}_n^2)_{st}^{ij} &= \\ &= \int_s^t (X_n^i(u) - X_n^i(s)) dX_n^j(u) = I_n^{ij}(t) - I_n^{ij}(s) - X_n^i(s)(X_n^j(t) - X_n^j(s)). \end{aligned} \tag{8.28}$$

Then $\mathbb{X}_n = (\mathbb{X}_n^1, \mathbb{X}_n^2)$ is the canonical rough path associated with the smooth path (X_n^1, X_n^2) .

We also set $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$, with $\mathbb{X}^1 \equiv 0$ and $\mathbb{X}_{st}^2$ given by the 2×2 matrix

$$(\mathbb{X}^2)_{st}^{ij} = \begin{pmatrix} 0 & \frac{t-s}{2} \\ -\frac{t-s}{2} & 0 \end{pmatrix} = \begin{cases} \frac{t-s}{2} & \text{if } i=1, j=2 \\ -\frac{t-s}{2} & \text{if } i=2, j=1 \\ 0 & \text{if } i=j \end{cases} . \quad (8.29)$$

It is not difficult to show that $(\mathbb{X}_n^2)^{ij} \rightarrow (\mathbb{X}^2)^{ij}$ in C_2^θ , for any $\theta \in (0, 1)$. As a consequence, for any $\alpha \in (\frac{1}{3}, \frac{1}{2})$, we have $\mathbb{X}_n^1 \rightarrow 0$ in C^α and $\mathbb{X}_n^2 \rightarrow \mathbb{X}^2$ in $C_2^{2\alpha}$, that is *the canonical rough path $\mathbb{X}_n = (\mathbb{X}_n^1, \mathbb{X}_n^2)$ converge in $\mathcal{R}_{\alpha,d}$ to the rough path $\mathbb{X} = (0, \mathbb{X}^2)$* .

Let us prove that $(\mathbb{X}_n^2)^{ij} \rightarrow (\mathbb{X}^2)^{ij}$ in C_2^θ , for any $\theta \in (0, 1)$. We have already shown the pointwise (actually uniform) convergence $I_n^{12}(t) \rightarrow \frac{1}{2}t$. With similar arguments, one shows the uniform convergence $I_n^{ij} \rightarrow I^{ij}$ defined by

$$I^{ij}(t) = \begin{pmatrix} 0 & \frac{t}{2} \\ -\frac{t}{2} & 0 \end{pmatrix} = \begin{cases} \frac{t}{2} & \text{if } i=1, j=2 \\ -\frac{t}{2} & \text{if } i=2, j=1 \\ 0 & \text{if } i=j \end{cases} .$$

It follows by (8.28) that we have the uniform convergence $(\mathbb{X}_n^2)_{st}^{ij} \rightarrow I^{ij}(t) - I^{ij}(s) = (\mathbb{X}^2)_{st}^{ij}$. To prove convergence in C_2^θ , it suffices to show a uniform ‘‘Lipschitz-like’’ bound $|(\mathbb{X}_n^2)_{st}^{ij}| \leq 2|t-s|$, which is easy:

$$\begin{aligned} |(\mathbb{X}_n^2)_{st}^{ij}| &\leq \int_s^t |X_n^i(u) - X_n^i(s)| |(X_n^j)'(u)| du \\ &\leq 2 \|X_n^i\|_\infty \|(X_n^j)'\|_\infty |t-s| \\ &= 2 \frac{1}{\sqrt{n}} \frac{n}{\sqrt{n}} |t-s| \\ &= 2|t-s|. \end{aligned}$$

8.9. THE PARA INTEGRAL

In this section we prove Proposition 8.5. Given continuous functions $X, Y: [0, T] \rightarrow \mathbb{R}$, let us define $R^1, R^2 \in C_2$

$$R^1(X, Y)_{st} := -Y_s \delta X_{st}, \quad R^2(X, Y)_{st} := X_t \delta Y_{st}, \quad 0 \leq s \leq t \leq T, \quad (8.30)$$

and note that

$$R_{st}^2 = R_{st}^1 + \delta(XY)_{st}.$$

Recalling Remark 8.6, it is easy to check that R^1 and R^2 satisfy

$$\delta R^1(X, Y)_{sut} = \delta R^2(X, Y)_{sut} = \delta Y_{su} \delta X_{ut}. \quad (8.31)$$

However, neither R^1 nor R^2 are in $C_2^{\alpha+\beta}$ in general, because we can only estimate

$$\|R^1\|_\alpha \leq \|Y\|_\infty \|\delta X\|_\alpha, \quad \|R^2\|_\beta \leq \|X\|_\infty \|\delta Y\|_\beta. \quad (8.32)$$

We are going to show that, by combining R^1 and R^2 in a suitable way, one can build R which satisfies both (8.8) and (8.9). This yields the existence of an integral.

We start with a technical approximation lemma.

LEMMA 8.19. *Given $f \in \mathcal{C}^\alpha$, there is a sequence $(\tilde{f}_n)_n \subset C^\infty$ such that*

$$f(x) = f(0) + \sum_{n \geq 0} \tilde{f}_n(x), \quad \forall x \in [0, T]. \quad (8.33)$$

One can choose $\tilde{f}_n$ so that for every $n \geq 0$

$$\|\tilde{f}_n\|_\infty \leq C \|\delta f\|_\alpha 2^{-n\alpha}, \quad \|\tilde{f}'_n\|_\infty \leq C \|\delta f\|_\alpha 2^{n(1-\alpha)}, \quad (8.34)$$

where $C \in (0, \infty)$ depends only on T (e.g. one can take $C = 2(T^\alpha + 1)$).

Proof. We may assume without loss of generality that $f(x) = 0$ (it suffices to redefine $f(x)$ as $f(x) - f(0)$, which does not change $\|\delta f\|_\alpha$.)

We extend $f: \mathbb{R} \rightarrow \mathbb{R}$ (e.g. with $f(x) := f(0)$ for $x \leq 0$ and $f(x) := f(T)$ for $x \geq T$) so that $\|f\|_\alpha$ is not changed. Then we fix a probability density $\phi: [-1, 1] \rightarrow [0, \infty)$ with $\phi \in C^1$ and for $n \geq 0$ we define the rescaled density

$$\phi_n(x) := 2^n \phi(2^n x).$$

Next, for $n \geq 0$, we set $f_n(x) := (f * \phi_n)(x)$, that is

$$\begin{aligned} f_n(x) &:= \int_{\mathbb{R}} f(z) \phi_n(x - z) dz = \int_{\mathbb{R}} f(x - z) \phi_n(z) dz \\ &= \int_{\mathbb{R}} f(x - \frac{z}{2^n}) \phi(z) dz. \end{aligned} \quad (8.35)$$

It is easy to check that $\|f_n - f\|_\infty \rightarrow 0$. Next we define

$$\tilde{f}_0(x) := f_0(x), \quad \text{for } k \geq 1: \quad \tilde{f}_k(x) := f_k(x) - f_{k-1}(x).$$

Note that $\sum_{k=0}^n \tilde{f}_k = f_n$, hence relation (8.33) is proved (we recall that $f(0) = 0$).

We now prove the first relation in (8.34). Since $f(0) = 0$, for all $x \in [0, T]$ we can write

$$\begin{aligned} |\tilde{f}_0(x)| &= |f_0(x)| \leq \int_{\mathbb{R}} |f(x - z)| \phi(z) dz = \int_{\mathbb{R}} |f(x - z) - f(0)| \phi(z) dz \\ &\leq \|\delta f\|_\alpha \int_{\mathbb{R}} |x - z|^\alpha \phi(z) dz \leq (T^\alpha + 1) \|\delta f\|_\alpha, \end{aligned}$$

where for the last inequality we have used $(x + y)^\alpha \leq x^\alpha + y^\alpha$ (for $\alpha < 1$ and $x, y \geq 0$), $x \leq T$ and $\int_{\mathbb{R}} |z|^\alpha \phi(z) dz \leq \int_{[-1, 1]} \phi(z) dz = 1$, because ϕ is a density supported on $[-1, 1]$. For $k \geq 1$ we estimate

$$\begin{aligned} |\tilde{f}_k(x)| &= |f_k(x) - f_{k-1}(x)| \\ &\leq \int_{\mathbb{R}} |f(x - \frac{z}{2^k}) - f(x - \frac{z}{2^{k-1}})| \phi(z) dz \\ &\leq 2^{-k\alpha} \|\delta f\|_\alpha \end{aligned}$$

again because $\int_{\mathbb{R}} |z|^\alpha \phi(z) dz \leq 1$. We have proved the first relation in (8.34).

We finally prove the second relation in (8.34). Note that

$$f'_n(x) = \int_{\mathbb{R}} f(z) \phi'_n(x-z) dz = 2^n \int_{\mathbb{R}} f(x - \frac{z}{2^n}) \phi'(z) dz,$$

which has the same form as $f_n(x)$, see the last integral in (8.35), just with an extra multiplicative factor 2^n and with ϕ replaced by ϕ' . Arguing as before, we obtain

$$\begin{aligned} |\tilde{f}'_0(x)| &= |f'_0(x)| \leq (T^\alpha + 1) \left(\int_{[-1,1]} |\phi'(z)| dz \right) \|\delta f\|_\alpha, \\ |\tilde{f}'_k(x)| &= |f'_k(x) - f'_{k-1}(x)| \leq 2^{k(1-\alpha)} \left(\int_{[-1,1]} |\phi'(z)| dz \right) \|\delta f\|_\alpha, \end{aligned}$$

for $k \geq 1$. We can choose ϕ to be symmetric, decreasing on $[0, 1]$, with $\phi(0) = 1$ and $\phi(1) = 0$, so that

$$\int_{[-1,1]} |\phi'(z)| dz = 2 \int_0^1 (-\phi'(z)) dz = 2(\phi(0) - \phi(1)) = 2,$$

and this completes the proof. $\square$

Proof of Proposition 8.5. The existence of an integral is an immediate consequence of Remark 8.6, because if we define $R_{st} := J_{\prec}(X, Y)_{st}$, then both relations (8.8) and (8.9) are satisfied.

It remains to build $J_{\prec}$. Let us write, applying Lemma 8.19,

$$X(x) = X(0) + \sum_{m \geq 0} \tilde{X}_m(x), \quad Y(x) = Y(0) + \sum_{n \geq 0} \tilde{Y}_n(x).$$

Recalling (8.30), we define

$$J_{\prec}(X, Y) := \sum_{0 \leq m \leq n} R^1(\tilde{X}_n, \tilde{Y}_m) + \sum_{0 \leq n < m} R^2(\tilde{X}_n, \tilde{Y}_m). \quad (8.36)$$

We show below that the series converge uniformly. Note that $\sum_{n \geq 0} \tilde{X}_n(x) = X(x) - X(0)$, hence $\sum_{n \geq 0} \delta \tilde{X}_n = \delta(X - X(0)) = \delta X$, and similarly for Y . Applying (8.31), we get

$$\begin{aligned} \delta J_{\prec}(X, Y)_{sut} &= \sum_{0 \leq m \leq n} (\delta \tilde{Y}_n)_{su} (\delta \tilde{X}_m)_{ut} + \sum_{0 \leq n < m} (\delta \tilde{Y}_n)_{su} (\delta \tilde{X}_m)_{ut} \\ &= \left(\sum_{n \geq 0} (\delta \tilde{Y}_n)_{su} \right) \left(\sum_{m \geq 0} (\delta \tilde{X}_m)_{ut} \right) = \delta Y_{su} \delta X_{ut}, \end{aligned}$$

which proves (8.7). We now prove (8.6). Note that, by (8.34),

$$|(\delta \tilde{X}_n)_{st}| \leq \|\tilde{X}'_n\|_\infty |t - s| \leq C \|\delta X\|_\alpha 2^{-\alpha n} (2^n |t - s|),$$

but at the same time, always by (8.34),

$$|(\delta \tilde{X}_n)_{st}| \leq |\tilde{X}_n(s)| + |\tilde{X}_n(t)| \leq 2 \|\tilde{X}_n\|_\infty \leq 2C \|\delta X\|_\alpha 2^{-\alpha n}.$$

Altogether, using the notation $x \wedge y := \min\{x, y\}$,

$$|(\delta \tilde{X}_n)_{st}| \leq 2C \|\delta X\|_\alpha 2^{-\alpha n} (2^n |t - s| \wedge 1).$$

Similarly

$$|(\delta \tilde{Y}_m)_{st}| \leq 2C \|\delta Y\|_\beta 2^{-\beta m} (2^m |t-s| \wedge 1).$$

Recalling (8.30) and applying again (8.34), we get

$$\begin{aligned} |R^1(\tilde{X}_n, \tilde{Y}_m)_{st}| &\leq \|\tilde{Y}_m\|_\infty |(\delta \tilde{X}_n)_{st}| \\ &\leq 2C^2 \|\delta X\|_\alpha \|\delta Y\|_\beta 2^{-\alpha n} 2^{-\beta m} (2^n |t-s| \wedge 1). \end{aligned}$$

and similarly

$$\begin{aligned} |R^2(\tilde{X}_n, \tilde{Y}_m)_{st}| &\leq \|\tilde{X}_n\|_\infty |(\delta \tilde{Y}_m)_{st}| \\ &\leq 2C^2 \|\delta X\|_\alpha \|\delta Y\|_\beta 2^{-\alpha n} 2^{-\beta m} (2^m |t-s| \wedge 1) \end{aligned}$$

These relations show that the series in (8.36) converge indeed uniformly. We now plug these estimates into (8.36), getting

$$\begin{aligned} |J_{\prec}(X, Y)_{st}| &\leq 2C^2 \|\delta X\|_\alpha \|\delta Y\|_\beta \left(\sum_{0 \leq m \leq n} 2^{-\alpha n} 2^{-\beta m} (2^m |t-s| \wedge 1) \right. \\ &\quad \left. + \sum_{0 \leq n < m} 2^{-\alpha n} 2^{-\beta m} (2^n |t-s| \wedge 1) \right). \end{aligned} \quad (8.37)$$

Let us set for convenience

$$\bar{k} = \bar{k}_{st} := \log_2 \frac{1}{|t-s|},$$

so that $2^m |t-s| \leq 2$ if and only if $m \leq \bar{k}$. Since $\sum_{n=m}^{\infty} 2^{-\alpha n} \leq \frac{1}{1-2^{-\alpha}} 2^{-\alpha m}$, the first sum in (8.37) can be bounded as follows (neglecting the prefactor $(1-2^{-\alpha})^{-1}$):

$$\begin{aligned} \sum_{m \geq 0} 2^{-(\alpha+\beta)m} (2^m |t-s| \wedge 1) &\leq |t-s| \sum_{0 \leq m < \bar{k}} 2^{(1-\alpha-\beta)m} + \sum_{m \geq \bar{k}} 2^{-(\alpha+\beta)m} \\ &\leq |t-s| \frac{2^{(1-\alpha-\beta)\bar{k}}}{2^{1-\alpha-\beta} - 1} + \frac{2^{-(\alpha+\beta)\bar{k}}}{1 - 2^{-(\alpha+\beta)}} \\ &\leq \left\{ \frac{1}{2^{1-\alpha-\beta} - 1} + \frac{1}{1 - 2^{-(\alpha+\beta)}} \right\} |t-s|^{\alpha+\beta}. \end{aligned}$$

The same estimates apply to the second sum in (8.37), hence (8.6) is proved. $\square$

Remark 8.20. In the previous proof, if $\alpha + \beta = 1$, then we have

$$\sum_{0 \leq m < \bar{k}} \underbrace{2^{(1-\alpha-\beta)m}}_{=1} = \bar{k} = \log_2 \frac{1}{|t-s|}$$

and therefore we obtain, instead of (8.6), that

$$|J_{\prec}(f, g)_{st}| \lesssim |t-s| \left| \log \frac{1}{|t-s|} \right|, \quad 0 \leq s < t \leq T.$$

CHAPTER 9

ROUGH INTEGRATION

9.1. CONTROLLED PATHS

We fix $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$. We recall that fixing a α -rough path $\mathbb{X}$ over X as in Definition 8.10 is equivalent to choosing a solution $(I, \mathbb{X}^2)$ to (8.17), with I and $\mathbb{X}^2$ representing our choices of the integrals, respectively,

$$I_t =: \int_0^t X_r \otimes dX_r, \quad \mathbb{X}_{st}^2 =: \int_s^t (X_r - X_s) \otimes dX_r = I_t - I_s - X_s \otimes (X_t - X_s).$$

The key point is that, having fixed a choice of $\mathbb{X}^2$, it is now possible to give a *canonical definition of the integral* $\int_0^\cdot Y dX$ (depending on $\mathbb{X}^2$) for a wide class of $Y \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^k \otimes (\mathbb{R}^d)^*)$, namely those *paths Y which are controlled by $\mathbb{X}$* (see below). In order to introduce this notion, let us recall that, given $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ and $Y \in \mathcal{C}^\beta([0, T]; \mathbb{R}^k \otimes (\mathbb{R}^d)^*)$, we look now for $J: [0, T] \rightarrow \mathbb{R}^k$ and $R^J: [0, T]_{\leq}^2 \rightarrow \mathbb{R}^k$ such that, in analogy with (8.4),

$$\delta J_{st} = Y_s \delta X_{st} + R_{st}^J, \quad |R_{st}^J| \lesssim |t - s|^{\alpha+\beta}.$$

We say that J is *controlled by $\mathbb{X}$* . It is actually a very fruitful idea to require that *each component of Y is also controlled by $\mathbb{X}$* . This is the content of the next

DEFINITION 9.1. Let $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, $\eta \in]0, 1]$ and $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ an α -rough path on $\mathbb{R}^d$. A pair $\mathbf{Z} = (Z, Z^1): [0, T] \rightarrow \mathbb{R}^k \times (\mathbb{R}^k \otimes (\mathbb{R}^d)^*)$ with Z of class $\mathcal{C}^\alpha$ and Z^1 of class $\mathcal{C}^\eta$ is a path $(\alpha + \eta)$ -controlled by $\mathbb{X}$ if

$$\delta Z_{st} = Z_s^1 \mathbb{X}_{st}^1 + Z_{st}^{[2]}, \quad |Z_{st}^{[2]}| \lesssim |t - s|^{\alpha+\eta}, \quad (s, t) \in [0, T]_{\leq}^2. \quad (9.1)$$

The function Z^1 is called a *derivative of Z with respect to $\mathbb{X}$* and $Z^{[2]}$ is the *remainder of the couple (Z, Z^1)* .

For a fixed α -rough path $\mathbb{X}$ on $\mathbb{R}^d$, we denote by $\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}(\mathbb{R}^k)$ the space of paths $(\alpha + \eta)$ -controlled by $\mathbb{X}$ with values in $\mathbb{R}^k$.

Remark 9.2. Note that, if $\alpha + \eta \leq 1$, in general Z^1 is *not* determined by $(Z, \mathbb{X}^1)$, so that we say that Z^1 is a *derivative* rather than *the derivative* of Z . Viceversa, Z is *not* determined by $(Z^1, \mathbb{X}^1)$: if (Z, Z^1) is $(\alpha + \eta)$ -controlled by $\mathbb{X}$ and $f \in \mathcal{C}^{\alpha+\eta}([0, T]; \mathbb{R}^k)$ then $(Z + f, Z^1)$ is also $(\alpha + \eta)$ -controlled by $\mathbb{X}$.

It is now clear from the definitions that, unlike rough paths, $(\alpha + \eta)$ -controlled paths have a natural linear structure, in particular as a linear subspace of $\mathcal{C}^\alpha \times \mathcal{C}^\eta$.

Exercise 9.1. Show that for each $i, j = 1, \dots, d$, setting $[0, T] \ni t \mapsto (\mathbb{X}_{0t}^1, \text{Id}) \in \mathbb{R}^d \times (\mathbb{R}^d \otimes (\mathbb{R}^d)^*)$ and $[0, T] \ni t \mapsto (\mathbb{X}_{0t}^2, \mathbb{X}_{0t}^1 \otimes \text{Id}) \in \mathbb{R}^d \otimes \mathbb{R}^d \times (\mathbb{R}^d \otimes \mathbb{R}^d \otimes (\mathbb{R}^d)^*)$ are paths 2α -controlled by $\mathbb{X}$.

9.2. THE ROUGH INTEGRAL

Now we can finally show how to modify the germ $Y_s(X_t - X_s)$ in order to obtain a well-defined integration theory.

PROPOSITION 9.3. *Let $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, $\eta \in]0, 1]$ and $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ a α -rough path on $\mathbb{R}^d$. If $\mathbf{Z} = (Z, Z^1)$ is $(\alpha + \eta)$ -controlled by $\mathbb{X}$ as in Definition 9.1, with $Z: [0, T] \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ and $Z^1: [0, T] \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^d)^*$, then the germ*

$$A_{st} = Z_s \mathbb{X}_{st}^1 + Z_s^1 \mathbb{X}_{st}^2$$

satisfies $\delta A \in C_3^{2\alpha+\eta}$.

Therefore if $2\alpha + \eta > 1$ we can canonically define $J_t = \int_0^t Z \, d\mathbb{X}$ as the unique function $J: [0, T] \rightarrow \mathbb{R}^k$ such that $J_0 = 0$ and $\delta J - A \in C_2^{2\alpha+\eta}$, namely

$$|J_t - J_s - Z_s \mathbb{X}_{st}^1 - Z_s^1 \mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha+\eta},$$

and we have

$$J_t = \lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} (Z_{t_i} \mathbb{X}_{t_i t_{i+1}}^1 + Z_{t_i}^1 \mathbb{X}_{t_i t_{i+1}}^2)$$

along arbitrary partitions $\mathcal{P}$ of $[0, t]$ with vanishing mesh $|\mathcal{P}| \rightarrow 0$.

Proof. We compute by (8.20)

$$\begin{aligned} \delta A_{sut} &= -\delta Z_{su} \mathbb{X}_{ut}^1 + Z_s^1 \delta \mathbb{X}_{sut}^2 - \delta Z_{su}^1 \mathbb{X}_{ut}^2 \\ &= -(\delta Z_{su} - Z_s^1 \mathbb{X}_{su}^1) \mathbb{X}_{ut}^1 - \delta Z_{su}^1 \mathbb{X}_{ut}^2 \\ &= -Z_{su}^{[2]} \mathbb{X}_{ut}^1 - \delta Z_{su}^1 \mathbb{X}_{ut}^2. \end{aligned} \tag{9.2}$$

Then by (2.8)

$$\begin{aligned} |\delta A_{sut}| &\leq \|Z^{[2]}\|_{\alpha+\eta} |u - s|^{\alpha+\eta} \|\mathbb{X}^1\|_{\alpha} |t - u|^{\alpha} + \|\delta Z^1\|_{\eta} |u - s|^{\eta} \|\mathbb{X}^2\|_{2\alpha} |t - u|^{2\alpha} \\ &\leq (\|Z^{[2]}\|_{\alpha+\eta} \|\mathbb{X}^1\|_{\alpha} + \|\delta Z^1\|_{\eta} \|\mathbb{X}^2\|_{2\alpha}) |t - s|^{2\alpha+\eta}. \end{aligned} \tag{9.3}$$

Since $\delta A \in C_3^{2\alpha+\eta}$, if $2\alpha + \eta > 1$ we can apply the Sewing Lemma and define $J^{[3]} := -\Lambda(\delta A)$ and $J: [0, T] \rightarrow \mathbb{R}^k$ such that $J_0 = 0$ and $\delta J = A + J^{[3]}$ where Λ is the Sewing Map of Theorem 6.11, namely

$$J_0 = 0, \quad \delta J_{st} = Z_s \mathbb{X}_{st}^1 + Z_s^1 \mathbb{X}_{st}^2 + J_{st}^{[3]}, \quad |J_{st}^{[3]}| \lesssim |t - s|^{2\alpha+\eta}. \tag{9.4}$$

The last assertion on the convergence of the generalised Riemann sums follows from (6.13). $\square$

We have in particular proved by (6.14) and (9.3) that

$$\|J^{[3]}\|_{2\alpha+\eta} \leq K_{2\alpha+\eta} (\|Z^{[2]}\|_{\alpha+\eta} \|\mathbb{X}^1\|_{\alpha} + \|\delta Z^1\|_{\eta} \|\mathbb{X}^2\|_{2\alpha}). \tag{9.5}$$

We stress that the function J depends on $(\mathbf{Z}, \mathbb{X})$, in particular on Z^1 as well. We use the following notations

$$\mathbf{J} := (J, Z), \quad \int_0^t \mathbf{Z} \, d\mathbb{X} := (J_t, Z_t) = \mathbf{J}_t. \quad (9.6)$$

We shall see in Proposition 9.5 below that $\mathbf{J}: [0, T] \rightarrow \mathbb{R}^k \times (\mathbb{R}^k \otimes (\mathbb{R}^d)^*)$ is 2α -controlled by $\mathbb{X}$, i.e. Z is a derivative of J with respect to $\mathbb{X}$ as in Definition 9.1.

We define a norm $\|\cdot\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$ and a seminorm $[\cdot]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$ on the space $\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$ of paths $(\alpha + \eta)$ -controlled by $\mathbb{X}$, defined as follows:

$$\begin{aligned} \|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}} &:= |Z_0| + |Z_0^1| + [\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}, & \mathbf{Z} &= (Z, Z^1) \\ [\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}} &:= \|\delta Z^1\|_{\eta} + \|Z^{[2]}\|_{\alpha+\eta}, & Z_{st}^{[2]} &= \delta Z_{st} - Z_s^1 \mathbb{X}_{st}^1, \end{aligned} \quad (9.7)$$

as in (9.1). Recall that we defined the standard norm $\|f\|_{\mathcal{C}^\alpha} = \|f\|_{\infty} + \|\delta f\|_{\alpha}$ in (1.13).

LEMMA 9.4. *We have the equivalence of norms for all $\mathbf{Z} = (Z, Z^1) \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$*

$$\|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}} \leq \|Z\|_{\mathcal{C}^\alpha} + \|Z^1\|_{\mathcal{C}^\eta} + \|Z^{[2]}\|_{\alpha+\eta} \leq C \|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}, \quad (9.8)$$

where $C > 0$ is an explicit constant which depends only on $(\mathbb{X}, T, \alpha, \eta)$. In particular, $(\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}, \|\cdot\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}})$ is a Banach space.

Proof. The first inequality in (9.8) is obvious by the definition of the norm $\|\cdot\|_{\mathcal{C}^\alpha}$. In order to prove the second one, first we note that by (1.15)

$$\|f\|_{\mathcal{C}^\eta} = \|f\|_{\infty} + \|\delta f\|_{\eta} \leq (1 + T^\eta)(|f_0| + \|\delta f\|_{\eta}).$$

This shows that $\|Z^1\|_{\mathcal{C}^\eta} \lesssim \|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$ for $(Z, Z^1) \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$. Now, since $\delta Z_{st} = Z_s^1 \mathbb{X}_{st}^1 + Z_{st}^{[2]}$ by (9.1),

$$\begin{aligned} \|\delta Z\|_{\alpha} &\leq \|Z^1\|_{\infty} \|\mathbb{X}^1\|_{\alpha} + \|Z^{[2]}\|_{\alpha} \\ &\leq C_{T,\eta}(|Z_0^1| + \|\delta Z^1\|_{\eta}) \|\mathbb{X}^1\|_{\alpha} + T^\eta \|Z^{[2]}\|_{\alpha+\eta}, \end{aligned}$$

namely $\|Z\|_{\mathcal{C}^\alpha} \lesssim \|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$. Finally $\|Z^{[2]}\|_{\alpha+\eta} \leq \|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$. The proof is complete. $\square$

9.3. CONTINUITY PROPERTIES OF THE ROUGH INTEGRAL

We wrote before Definition 9.1 that the notion of controlled path aimed at making the rough integral map $(Z, Z^1) \mapsto (J, Z)$ iterable, where we use the notation of Proposition 9.3. In order to make this precise, we need the following important

PROPOSITION 9.5. *Let $\mathbb{X}$ be a α -rough path on $\mathbb{R}^d$ with $\alpha \in]\frac{1}{3}, \frac{1}{2}]$, $\eta \in]1 - 2\alpha, 1]$ and $\mathbf{Z} \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$ a path $(\alpha + \eta)$ -controlled by $\mathbb{X}$. Then, in the notation of (9.6),*

- $\mathbf{J} = \int_0^\cdot \mathbf{Z} \, d\mathbb{X}$ is 2α -controlled by $\mathbb{X}$

- the map $\mathcal{D}_{\mathbb{X}}^{\alpha+\eta} \ni \mathbf{Z} \mapsto \mathbf{J} \in \mathcal{D}_{\mathbb{X}}^{2\alpha}$ is linear and for all $\mathbf{Z} \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$

$$[\mathbf{J}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}} \leq 2(1 + \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}})[|Z_0^1| + T^\eta(1 + K_{2\alpha+\eta})[\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}]. \quad (9.9)$$

Proof. Recall first (9.5), so that in particular $\|J^{[3]}\|_{2\alpha+\eta} < +\infty$. Now $J_{st}^{[2]} = Z_s^1 \mathbb{X}_{st}^2 + J_{st}^{[3]}$ satisfies

$$\|J^{[2]}\|_{2\alpha} \leq \|Z^1\|_\infty \|\mathbb{X}^2\|_{2\alpha} + \|J^{[3]}\|_{2\alpha} \leq \|Z^1\|_\infty \|\mathbb{X}^2\|_{2\alpha} + T^\eta \|J^{[3]}\|_{2\alpha+\eta}. \quad (9.10)$$

Finally $\delta J_{st} = Z_s \mathbb{X}_{st}^1 + J_{st}^{[2]}$ and therefore

$$\|\delta J\|_\alpha \leq \|Z\|_\infty \|\mathbb{X}^1\|_\alpha + \|Z^1\|_\infty \|\mathbb{X}^2\|_{2\alpha} + T^{\alpha+\eta} \|J^{[3]}\|_{2\alpha+\eta}.$$

Therefore $(J, Z, J^{[2]}) \in \mathcal{C}^\alpha \times \mathcal{C}^\alpha \times C_2^{2\alpha}$ and we obtain that (J, Z) is 2α -controlled by $\mathbb{X}$.

We prove now the second assertion. Since $\delta Z_{st} = Z_s^1 \mathbb{X}_{st}^1 + Z_{st}^{[2]}$, by (1.40)

$$\begin{aligned} \|\delta Z\|_\alpha &\leq \|Z^1\|_\infty \|\mathbb{X}^1\|_\alpha + T^\eta \|Z^{[2]}\|_{\alpha+\eta} \\ &\leq (\|\mathbb{X}^1\|_\alpha + 1)(|Z_0^1| + T^\eta [\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}). \end{aligned}$$

Now, analogously to (9.10), again by (1.40)

$$\begin{aligned} \|J^{[2]}\|_{2\alpha} &\leq \|Z^1\|_\infty \|\mathbb{X}^2\|_{2\alpha} + \|J^{[3]}\|_{2\alpha} \\ &\leq T^\eta \|J^{[3]}\|_{2\alpha+\eta} + \|\mathbb{X}^2\|_{2\alpha}(|Z_0^1| + T^\eta \|\delta Z^1\|_\eta). \end{aligned}$$

Therefore, since $\|\mathbb{X}^1\|_\alpha + \|\mathbb{X}^2\|_{2\alpha} = \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}}$, recall (8.23),

$$\|\delta Z\|_\alpha + \|J^{[2]}\|_{2\alpha} \leq T^\eta \|J^{[3]}\|_{2\alpha+\eta} + (1 + \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}})[|Z_0^1| + T^\eta [\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}].$$

By (9.5) we obtain

$$\begin{aligned} [\mathbf{J}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}} &= \|\delta Z\|_\alpha + \|J^{[2]}\|_{2\alpha} \leq \\ &\leq 2(1 + \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}})[|Z_0^1| + (1 + K_{2\alpha+\eta})T^\eta [\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}] \end{aligned}$$

The proof is complete. $\square$

We note that the estimate of the seminorm $[\mathbf{J}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}}$ in terms of $[\mathbf{Z}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$ rather than of the norm $\|\mathbf{J}\|_{\mathcal{D}_{\mathbb{X}}^{2\alpha}}$ in terms of $\|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$ plays an important role in Chapter 10 (with $\eta = \alpha$), see in particular (10.9). In any case, from (9.9) it is easy to obtain an estimate of $\|\mathbf{J}\|_{\mathcal{D}_{\mathbb{X}}^{2\alpha}}$: since $J_0 = 0$ and $J_0^1 = Z_0$, we obtain

$$\begin{aligned} \|\mathbf{J}\|_{\mathcal{D}_{\mathbb{X}}^{2\alpha}} &= |Z_0| + [\mathbf{J}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}} \leq \\ &\leq 2(1 + K_{2\alpha+\eta})(1 + \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}})(1 + T^\eta) \|\mathbf{Z}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}. \end{aligned}$$

Therefore the linear operator $\mathcal{D}_{\mathbb{X}}^{\alpha+\eta} \ni \mathbf{Z} \mapsto \int_0^\cdot \mathbf{Z} d\mathbb{X} \in \mathcal{D}_{\mathbb{X}}^{2\alpha}$ is continuous. In fact a stronger property holds: we have continuity of the map $(\mathbb{X}, \mathbf{Z}) \mapsto \int_0^\cdot \mathbf{Z} d\mathbb{X}$. In order to prove this, we need to introduce the following space

$$\mathcal{S}_{\alpha,\eta} := \{(\mathbb{X}, \mathbf{Z}): \mathbb{X} \text{ is a } \alpha\text{-rough path, } \mathbf{Z} \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}\},$$

and the following quantity for $\mathbf{Z} \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$ and $\bar{\mathbf{Z}} \in \mathcal{D}_{\bar{\mathbb{X}}}^{\alpha+\eta}$

$$[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta} := \|\delta Z^1 - \delta \bar{Z}^1\|_{\eta} + \|Z^{[2]} - \bar{Z}^{[2]}\|_{\alpha+\eta},$$

where $Z^{[2]} = \delta Z - Z^1 \mathbb{X}^1$ and $\bar{Z}^{[2]} = \delta \bar{Z} - \bar{Z}^1 \bar{\mathbb{X}}^1$, recall (9.7). We endow $\mathcal{S}_{\alpha, \eta}$ with the distance (see (8.24) for the definition of $d_{\mathcal{R}_{\alpha, d}}$)

$$d_{\alpha, \eta}((\mathbb{X}, \mathbf{Z}), (\bar{\mathbb{X}}, \bar{\mathbf{Z}})) = d_{\mathcal{R}_{\alpha, d}}(\mathbb{X}, \bar{\mathbb{X}}) + |Z_0 - \bar{Z}_0| + |Z_0^1 - \bar{Z}_0^1| + [\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta}.$$

Let us note that in the case $\mathbb{X} = \bar{\mathbb{X}}$, we have

$$[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta} = [\mathbf{Z} - \bar{\mathbf{Z}}]_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}, \quad d_{\alpha, \eta}((\mathbb{X}, \mathbf{Z}), (\mathbb{X}, \bar{\mathbf{Z}})) = \|\mathbf{Z} - \bar{\mathbf{Z}}\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}},$$

see the definition (9.7) of the norm $\|\cdot\|_{\mathcal{D}_{\mathbb{X}}^{\alpha+\eta}}$. Note that $[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta}$ is *not* a function of $\mathbf{Z} - \bar{\mathbf{Z}}$ when $\mathbb{X} \neq \bar{\mathbb{X}}$.

PROPOSITION 9.6. (LOCAL LIPSCHITZ ESTIMATE) *Let $\alpha \in]\frac{1}{3}, \frac{1}{2}]$ and $\eta \in]1 - 2\alpha, 1]$. The function $\mathcal{S}_{\alpha, \eta} \ni (\mathbb{X}, \mathbf{Z}) \mapsto (\mathbb{X}, \int_0 \cdot \mathbf{Z} d\mathbb{X}) \in \mathcal{S}_{\alpha, \alpha}$ is continuous with respect to the distances $d_{\alpha, \eta}$ and $d_{\alpha, \alpha}$.*

More precisely, for every $M \geq 0$ there is $K_{M, \alpha, \eta} \geq 0$ such that for all $(\mathbb{X}, \mathbf{Z}), (\bar{\mathbb{X}}, \bar{\mathbf{Z}}) \in \mathcal{S}_{\alpha, \eta}$ satisfying

$$1 + T^{\eta} + \|\mathbb{X}\|_{\mathcal{R}_{\alpha, d}} + \|\bar{\mathbf{Z}}\|_{\mathcal{D}_{\bar{\mathbb{X}}}^{\alpha+\eta}} \leq M,$$

setting $\mathbf{J} := \int_0 \cdot \mathbf{Z} d\mathbb{X}$ and $\bar{\mathbf{J}} := \int_0 \cdot \bar{\mathbf{Z}} d\bar{\mathbb{X}}$ we have

$$\begin{aligned} d_{\alpha, \alpha}((\mathbb{X}, \mathbf{J}), (\bar{\mathbb{X}}, \bar{\mathbf{J}})) &\leq \\ &\leq 2M^2(1 + K_{2\alpha+\eta})[d_{\mathcal{R}_{\alpha, d}}(\mathbb{X}, \bar{\mathbb{X}}) + |Z_0 - \bar{Z}_0| + |Z_0^1 - \bar{Z}_0^1| + T^{\eta}[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta}] \\ &\leq 2M^3(1 + K_{2\alpha+\eta}) d_{\alpha, \eta}((\mathbb{X}, \mathbf{Z}), (\bar{\mathbb{X}}, \bar{\mathbf{Z}})). \end{aligned}$$

Proof. Let $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ and $\bar{\mathbb{X}} = (\bar{\mathbb{X}}^1, \bar{\mathbb{X}}^2)$ be α -rough paths and $\mathbf{Z} \in \mathcal{D}_{\mathbb{X}}^{\alpha+\eta}$, $\bar{\mathbf{Z}} \in \mathcal{D}_{\bar{\mathbb{X}}}^{\alpha+\eta}$. We argue as in the proof of (9.9), using furthermore a number of times the simple estimate

$$|ab - \bar{a}\bar{b}| \leq |a - \bar{a}| |b| + |\bar{a}| |b - \bar{b}|. \quad (9.11)$$

We set for notational convenience $\varepsilon := T^{\eta}$. Then, since $\delta Z_{st} = Z_s^1 \mathbb{X}_{st}^1 + Z_{st}^{[2]}$, by (1.40)

$$\begin{aligned} \|\delta Z - \delta \bar{Z}\|_{\alpha} &\leq \|Z^1 - \bar{Z}^1\|_{\infty} \|\mathbb{X}^1\|_{\alpha} + \|\bar{Z}^1\|_{\infty} \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} + \varepsilon \|Z^{[2]} - \bar{Z}^{[2]}\|_{\alpha+\eta} \\ &\leq (\|\mathbb{X}^1\|_{\alpha} + 1)(|Z_0^1 - \bar{Z}_0^1| + \varepsilon [\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta}) + M^2 \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha}, \end{aligned}$$

since by assumption

$$\|\bar{Z}^1\|_{\infty} \leq |\bar{Z}_0^1| + \varepsilon \|\delta \bar{Z}^1\|_{\eta} \leq (1 + \varepsilon)(|\bar{Z}_0^1| + \|\delta \bar{Z}^1\|_{\eta}) \leq M^2.$$

Now $J_{st}^{[2]} = Z_s^1 \mathbb{X}_{st}^2 + J_{st}^{[3]}$, so that arguing similarly

$$\begin{aligned} \|J^{[2]} - \bar{J}^{[2]}\|_{2\alpha} &\leq \|J^{[3]} - \bar{J}^{[3]}\|_{2\alpha} + \|Z^1 \mathbb{X}^2 - \bar{Z}^1 \bar{\mathbb{X}}^2\|_{2\alpha} \leq \\ &\leq \varepsilon \|J^{[3]} - \bar{J}^{[3]}\|_{2\alpha+\eta} + \|\mathbb{X}^2\|_{2\alpha}(|Z_0^1 - \bar{Z}_0^1| + \varepsilon \|\delta Z^1 - \delta \bar{Z}^1\|_{\eta}) + M^2 \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha}. \end{aligned}$$

Therefore, since $1 + \|\mathbb{X}^1\|_\alpha + \|\mathbb{X}^2\|_{2\alpha} = 1 + \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}} \leq M$,

$$\begin{aligned} & \|\delta Z - \delta \bar{Z}\|_\alpha + \|J^{[2]} - \bar{J}^{[2]}\|_{2\alpha} \leq \\ & \leq \varepsilon \|J^{[3]} - \bar{J}^{[3]}\|_{2\alpha+\eta} + M^2(|Z_0^1 - \bar{Z}_0^1| + \varepsilon[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta} + d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}})). \end{aligned}$$

We can estimate in the same way

$$\begin{aligned} \|\delta A - \delta \bar{A}\|_{2\alpha+\eta} & \leq \|Z^{[2]} - \bar{Z}^{[2]}\|_{\alpha+\eta} \|\mathbb{X}^1\|_\alpha + \|\bar{Z}^{[2]}\|_{\alpha+\eta} \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha + \\ & \quad + \|\delta Z^1 - \delta \bar{Z}^1\|_\eta \|\mathbb{X}^2\|_{2\alpha} + \|\delta \bar{Z}^1\|_\eta \|\mathbb{X}^2 - \bar{\mathbb{X}}^2\|_{2\alpha} \\ & \leq [\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta} \|\mathbb{X}\|_{\mathcal{R}_{\alpha,d}} + [\bar{\mathbf{Z}}]_{\mathcal{D}_{\bar{\mathbb{X}}}^{\alpha+\eta}} d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}}) \\ & \leq M([\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta} + d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}})). \end{aligned}$$

By the Sewing bound (1.42)

$$\|J^{[3]} - \bar{J}^{[3]}\|_{2\alpha+\eta} \leq K_{2\alpha+\eta} M([\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta} + d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}})).$$

We obtain

$$\begin{aligned} [\mathbf{J}; \bar{\mathbf{J}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \alpha} & = \|\delta Z - \delta \bar{Z}\|_\alpha + \|J^{[2]} - \bar{J}^{[2]}\|_{2\alpha} \leq \\ & \leq M^2(1 + K_{2\alpha+\eta})[|Z_0^1 - \bar{Z}_0^1| + d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}}) + \varepsilon[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta}]. \end{aligned}$$

Since $J_0 - \bar{J}_0 = 0$, $J_0^1 - \bar{J}_0^1 = Z_0 - \bar{Z}_0$, we obtain

$$\begin{aligned} d_{\alpha,\alpha}((\mathbb{X}, \mathbf{J}), (\bar{\mathbb{X}}, \bar{\mathbf{J}})) & = d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}}) + |Z_0 - \bar{Z}_0| + [\mathbf{J}; \bar{\mathbf{J}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \alpha} \\ & \leq 2M^2(1 + K_{2\alpha+\eta})[|Z_0 - \bar{Z}_0| + |Z_0^1 - \bar{Z}_0^1| + d_{\mathcal{R}_{\alpha,d}}(\mathbb{X}, \bar{\mathbb{X}}) + \varepsilon[\mathbf{Z}; \bar{\mathbf{Z}}]_{\mathbb{X}, \bar{\mathbb{X}}, \alpha, \eta}]. \end{aligned}$$

The second estimate follows since we have assumed that $1 + \varepsilon \leq M$. $\square$

9.4. STOCHASTIC AND ROUGH INTEGRALS

In this section we explore the connections between Itô integrals and Young or rough integrals. We fix $\alpha \in]0, \frac{1}{2}[$ and a realisation of the Itô rough path $\mathbb{B}$ defined in (4.3) satisfying a.s. (4.4). We consider an adapted process $h: [0, T] \rightarrow (\mathbb{R}^d)^*$ with continuous paths and its Itô integral

$$I_t := \int_0^t h_s dB_s, \quad t \in [0, T]. \quad (9.12)$$

Let us suppose also that a.s. h is of class $\mathcal{C}^\beta$ with $\beta \in]0, 1[$. By (4.7) we have

$$|\delta I_{st} - h_s \mathbb{B}_{st}^1| = \left| \int_s^t \delta h_{sr} dB_r \right| \lesssim (t-s)^{\alpha+\beta}, \quad \forall 0 \leq s \leq t \leq T.$$

By Theorem 4.3, this means a.s. the Itô integral in (9.12) is a generalised integral of h in $d\mathbb{B}$ in the sense of Definition 8.1.

The situation is different according to the value of $\alpha + \beta$. If $\alpha + \beta > 1$, then we can apply Theorem 7.1 and we obtain that I_t is equal to the integral $\int_0^t h_s dB_s$ also in the Young sense. In this regime, we have uniqueness of generalised integrals in the sense of Definition 8.1. Moreover, by (6.13) we have a.s.

$$I_t = \lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} h_{t_i} (B_{t_{i+1}} - B_{t_i}),$$

where $\mathcal{P} = \{0 = t_0 < t_1 < \dots < t_k = t\}$ is a partition of $[0, t]$.

If $\alpha + \beta \leq 1$, then I is indeed only one of the generalised integrals as in Definition 8.1: for any $f: [0, T] \rightarrow \mathbb{R}$ of class $\mathcal{C}^{\alpha+\beta}$, then $I + f$ is also such a generalised integral. In this setting, in order to characterise uniquely the Itô integral among all generalised integrals, one can use (4.9): if we assume that, almost surely,

$$|\delta h_{sr} - h_s^1 \mathbb{B}_{sr}^1| \lesssim (r - s)^{\eta+\alpha},$$

for some adapted process $h^1 = (h_t^1)_{t \in [0, T]}$ of class $\mathcal{C}^\eta$ with $\eta \in]0, 1]$, then a.s.

$$|\delta I_{st} - h_s \mathbb{B}_{st}^1 - h_s^1 \mathbb{B}_{st}^2| = \left| \int_s^t (\delta h_{sr} - h_s^1 \mathbb{B}_{sr}^1) dB_r \right| \lesssim (t - s)^{2\alpha+\eta}.$$

By Proposition 9.3, if $2\alpha + \eta > 1$ then (I, h) is the rough integral of (h, h^1) with respect to $\mathbb{B}$, namely

$$(I_t, h_t) = \int_0^t (h, h^1) d\mathbb{B}, \quad t \geq 0,$$

as in (9.6). Moreover, by (6.13) we have a.s.

$$I_t = \lim_{|\mathcal{P}| \rightarrow 0} \sum_{i=0}^{\#\mathcal{P}-1} [h_{t_i} \mathbb{B}_{t_i t_{i+1}}^1 + h_{t_i}^1 \mathbb{B}_{t_i t_{i+1}}^2],$$

where $\mathcal{P} = \{0 = t_0 < t_1 < \dots < t_k = t\}$.

Let $(B_t)_{t \geq 0}$ be a Brownian motion in $\mathbb{R}^d$ and $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ be such that the SDE

$$Z_t = Z_0 + \underbrace{\int_0^t \sigma(Z_s) dB_s}_{\text{Itô integral}} \quad (9.13)$$

has a unique solution for all $Z_0 \in \mathbb{R}^k$.

THEOREM 9.7. (LYONS 98, DAVIE 07) *A.s. $(Z_t)_{t \geq 0}$ is the only continuous function $Y: [0, \infty) \rightarrow \mathbb{R}^k$ such that $Y_0 = Z_0$ and*

$$Y_t - Y_s - \sigma(Y_s) (B_t - B_s) - \underbrace{\sigma_2(Y_s) \int_s^t (B_r - B_s) \otimes dB_r}_{\text{Itô integral}} = o(t - s) \quad (9.14)$$

(where $\sigma_2(y) := \nabla \sigma(y) \sigma(y)$) uniformly over $0 \leq s \leq t \leq T$ for any $T \geq 0$.

Here the distribution on $\mathbb{R}$

$${}^{\circ}\sigma(Z_t)dB_t := \frac{d}{dt} \int_0^t \sigma(Z_s)dB_s$$

is analytically a singular product, since

$$Z \in \mathcal{C}^{\frac{1}{2}-\varepsilon}, \quad B \in \mathcal{C}^{\frac{1}{2}-\varepsilon}.$$

It is therefore not canonically defined if one only knows $\sigma(Z)$ and B .

Let us now consider the SDE

$$Z_t = Z_0 + \underbrace{\int_0^t \sigma(Z_s) \circ dB_s}_{\text{Stratonovich integral}} \quad (9.15)$$

and let us suppose that there is a unique solution for all $Z_0 \in \mathbb{R}^k$.

THEOREM 9.8. (LYONS 98, DAVIE 07) *A.s. $(Z_t)_{t \geq 0}$ is the only continuous function $Y: [0, \infty) \rightarrow \mathbb{R}^k$ such that $Y_0 = Z_0$ and*

$$Y_t - Y_s - \underbrace{\sigma(Y_s)(B_t - B_s) - \sigma_2(Y_s) \int_s^t (B_r - B_s) \otimes \circ dB_r}_{\text{Stratonovich integral}} = o(t - s) \quad (9.16)$$

uniformly over $0 \leq s \leq t \leq T$ for any $T \geq 0$.

Here the distribution on $\mathbb{R}$

$${}^{\circ}\sigma(Z_t) \circ dB_t := \frac{d}{dt} \int_0^t \sigma(Z_s) \circ dB_s \left[= \frac{d}{dt} \int_0^t \sigma(Z_s)dB_s + \underbrace{\frac{1}{2}\text{Tr}[\sigma_2(Z_t)]}_{\text{It\^o-Strato correction}} \right]$$

is analytically a singular product, since

$$Z \in \mathcal{C}^{\frac{1}{2}-\varepsilon}, \quad B \in \mathcal{C}^{\frac{1}{2}-\varepsilon}.$$

It is therefore not canonically defined if one only knows $\sigma(Z)$ and B .

The main message of Rough Paths is that:

- If we choose the product $B_t \dot{B}_t$, then many products like $\sigma(Z_t) \dot{B}_t$ are canonically defined.

This is a pathwise construction: for sample trajectories (fixed $\omega \in \Omega$)

- $(B_s(\omega), \int_0^t B_s dB_s(\omega))_{s \geq 0}$,
- $(h_s(\omega))_{s \geq 0}$,

we can compute canonically $(\int_0^t h_s dB_s(\omega))_{t \geq 0}$.

CHAPTER 10

ROUGH INTEGRAL EQUATIONS

In this chapter we go back to the finite difference equation (3.19) in the rough setting $\alpha \in (\frac{1}{2}, \frac{1}{3}]$, and we discuss its integral formulation that we already mentioned at the end of Section 7.3. Now that we have studied the rough integral in Chapter 9, we can indeed show that the equation

$$|Z_{st}^{[3]}| \lesssim |t-s|^{3\alpha}, \quad Z_{st}^{[3]} = \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma_2(Z_s) \mathbb{X}_{st}^2, \quad (10.1)$$

recall (3.18), can be interpreted in the context of controlled paths. Indeed, (10.1) suggests that, for any candidate solution Z , the pair $\mathbf{Z} = (Z, \sigma(Z))$ should be controlled by $\mathbb{X}$. At the same time, in order to apply Proposition 9.3 and interpret (10.1) as an integral equation, we are going to show that $(\sigma(Z), \sigma_2(Z))$ is controlled by $\mathbb{X}$. This is guaranteed by the following

LEMMA 10.1. *Let $\phi: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$ be of class C^2 and $\mathbf{f} = (f, f^1) \in \mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$. Set*

$$\phi(\mathbf{f}) := (\phi(f), \nabla \phi(f) f^1),$$

where $\phi(f): [0, T] \rightarrow \mathbb{R}^\ell$ is defined by $\phi(f)_t := \phi(f_t)$ and

$$\nabla \phi(f) f^1: [0, T] \rightarrow \mathbb{R}^\ell \otimes \mathbb{R}^d, \quad (\nabla \phi(f) f^1)_t^{ab} = \sum_{j=1}^k \partial_j \phi^a(f_t) \cdot (f_t^1)^{jb}.$$

Then $\phi(\mathbf{f}) \in \mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^\ell)$.

Proof. Analogously to (3.22) we have for $\mathbf{f} = (f, f^1) \in \mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$, setting $f_{st}^{[2]} := \delta f_{st} - f_s^1 \mathbb{X}_{st}^1$ as in (9.1),

$$\begin{aligned} \phi(\mathbf{f})_{st}^{[2]} &:= \phi(f_t) - \phi(f_s) - \nabla \phi(f_s) f_s^1 \mathbb{X}_{st}^1 \\ &= \nabla \phi(f_s) f_{st}^{[2]} + \int_0^1 [\nabla \phi(f_s + r \delta f_{st}) - \nabla \phi(f_s)] dr \delta f_{st} \\ &= \nabla \phi(f_s) f_{st}^{[2]} + \int_0^1 (1-u) \nabla^2 \phi(f_s + u \delta f_{st}) du \delta f_{st} \otimes \delta f_{st}. \end{aligned} \quad (10.2)$$

Then we can write using the estimate $|ab - \bar{a}\bar{b}| \leq |a - \bar{a}| |b| + |\bar{a}| |b - \bar{b}|$

$$\begin{aligned} |\nabla \phi(f_t) f_t^1 - \nabla \phi(f_s) f_s^1| &\leq c_{\phi, f}^{(1)} |f_t^1 - f_s^1| + c_{\phi, f}^{(2)} |f_t - f_s| \|f^1\|_\infty, \\ |\phi(\mathbf{f})_{st}^{[2]}| &\leq c_{\phi, f}^{(1)} |f_{st}^{[2]}| + c_{\phi, f}^{(2)} |\delta f_{st}|^2, \end{aligned} \quad (10.3)$$

where

$$c_{\phi, f}^{(1)} := \sup_{s \in [0, T]} |\nabla \phi(f_s)|, \quad c_{\phi, f}^{(2)} := \sup_{s, t \in [0, T], u \in [0, 1]} |\nabla^2 \phi(f_s + u \delta f_{st})|. \quad (10.4)$$

Therefore $(\phi(f), \nabla \phi(f) f^1)$ is controlled by $\mathbb{X}$. $\square$

This suggests that we can reinterpret the finite difference equation (10.1) as follows: we look for $Z: [0, T] \rightarrow \mathbb{R}^k$ such that $\mathbf{Z} = (Z, \sigma(Z))$ is controlled by $\mathbb{X}$ (namely it belongs to $\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$) and

$$\mathbf{Z}_t = (Z_0, 0) + \int_0^t \sigma(\mathbf{Z}) d\mathbb{X}, \quad \forall t \in [0, T]. \quad (10.5)$$

By Lemma 10.1, $\sigma(\mathbf{Z}) = (\sigma(Z), \nabla \sigma(Z) Z^1)$, but here $Z^1 = \sigma(Z)$, so that

$$\sigma(\mathbf{Z}) = (\sigma(Z), \nabla \sigma(Z) \sigma(Z)) = (\sigma(Z), \sigma_2(Z)),$$

is controlled by $\mathbb{X}$, where we use the notation $\sigma_2: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^* \otimes (\mathbb{R}^d)^*$

$$\sigma_2(y) := \nabla \sigma(y) \sigma(y), \quad [\sigma_2(y)]_{j\ell}^i := \sum_{a=1}^k \partial_a \sigma_j^i(y) \sigma_\ell^a(y).$$

By Proposition 9.3, the integral equation in (10.5) is equivalent to

$$|Z_{st}^{[3]}| \lesssim |t - s|^{3\alpha}, \quad Z_{st}^{[3]} = \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma_2(Z_s) \mathbb{X}_{st}^2. \quad (10.6)$$

Viceversa, if $Z \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^k)$ is such that $Z^{[3]} \in C_2^{3\alpha}$, then setting $Z^1 := \sigma(Z)$ the path $\mathbf{Z} = (Z, Z^1)$ is controlled by $\mathbb{X}$ and satisfies (10.5). Therefore, the integral equation (10.5) is equivalent to the finite difference equation (10.6).

10.1. LOCALIZATION ARGUMENT

PROPOSITION 10.2. *If we can prove local existence for the rough differential equation (10.6) under the assumption that σ is of class C^3 and $\sigma, \nabla \sigma, \nabla^2 \sigma, \nabla^3 \sigma$ are bounded, then we can prove local existence for (10.6) assuming only that σ is of class C^3 .*

Proof. Let σ be of class C^3 . Note that σ and its derivatives are bounded on the closed unit ball $B := \{z \in \mathbb{R}^k: |z - Z_0| \leq 1\}$, which is a compact subset of $\mathbb{R}^k$. Then we can find a function $\hat{\sigma}$ of class C^3 which is bounded with all its derivatives up to the third on the whole $\mathbb{R}^k$ and coincides with σ on B . By local existence for $\hat{\sigma}$, there is a solution $\hat{Z}: [0, T] \rightarrow \mathbb{R}^k$ of the RDE (10.6) with σ replaced by $\hat{\sigma}$. Since Z is continuous with $Z_0 \in B$, we can find $T' > 0$ such that $Z_t \in B$ for all $t \in [0, T']$. Then $\sigma(Z_t) = \hat{\sigma}(Z_t)$ and $\sigma_2(Z_t) = \hat{\sigma}_2(Z_t)$ for all $t \in [0, T']$, so that Z is a solution of the original RDE (10.6) on the shorter time interval $[0, T']$. We have proved *local existence* assuming only that σ is of class C^3 . $\square$

10.2. INVARIANCE

In this section we prepare the ground for a contraction argument to be proved in the next section. We start with an estimate of $[\sigma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}$ in terms of $[\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}$, under the assumption that σ is of class C^2 with bounded first and second derivative. We fix $D > 0$ such that

$$D \geq \max \{ \|\nabla \sigma\|_\infty, \|\nabla^2 \sigma\|_\infty \}.$$

LEMMA 10.3. *Let $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ be of class C^2 with $\|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty \leq D$, for some $D < +\infty$. Then for some $C > 0$ and any $\mathbf{f} = (f, f^1) \in \mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$*

$$[\sigma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes \mathbb{R}^d)} \leq D([\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} + \|f^1\|_\infty \|\delta f\|_\alpha + \|\delta f\|_\alpha^2). \quad (10.7)$$

Proof. By (10.3) we have

$$\|\delta(\nabla \sigma(f) f^1)\|_\alpha \leq D(\|\delta f^1\|_\alpha + \|f^1\|_\infty \|\delta f\|_\alpha),$$

$$\|\sigma(\mathbf{f})^{[2]}\|_{2\alpha} \leq D(\|f^{[2]}\|_{2\alpha} + \|\delta f\|_\alpha^2).$$

Therefore, recalling (9.7),

$$\begin{aligned} [\sigma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes \mathbb{R}^d)} &= \|\delta(\nabla \sigma(f) f^1)\|_\alpha + \|\sigma(\mathbf{f})^{[2]}\|_{2\alpha} \\ &\leq D([\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} + \|f^1\|_\infty \|\delta f\|_\alpha + \|\delta f\|_\alpha^2). \end{aligned}$$

where, in the last inequality, we apply (9.8). $\square$

We define $\Gamma: \mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k) \rightarrow \mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$

$$\Gamma(\mathbf{f}) := (Z_0, 0) + \int_0^\cdot \sigma(\mathbf{f}) d\mathbb{X},$$

(we know that indeed Γ maps $\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$ into $\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)$ by Lemma 10.1). In other words, $\Gamma(f, f^1)$ is equal to the only $(J, J^1) \in \mathcal{D}_{\mathbb{X}}^{2\alpha}$ such that

$$J_0 = Z_0, \quad J_s^1 = \sigma(f_s), \quad \delta J_{st} - \sigma(f_s) \mathbb{X}_{st}^1 - \nabla \sigma(f_s) f_s^1 \mathbb{X}_{st}^2 \in C_2^{3\alpha}. \quad (10.8)$$

We want to construct solutions to (10.6) by a fixed point argument for T small enough. Let $M > 0$ and $\mathbb{X}$ such that $\|\mathbb{X}^1\|_\alpha + \|\mathbb{X}^2\|_{2\alpha} \leq M$ and

$$\mathcal{B} := \{\mathbf{f} = (f, f^1) \in \mathcal{D}_{\mathbb{X}}^{2\alpha}: (f_0, f_0^1) = (Z_0, \sigma(Z_0)), [\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq 4C\}, \quad (10.9)$$

where

$$C := (1 + M)D\|\sigma\|_\infty. \quad (10.10)$$

LEMMA 10.4. *If $T^\alpha \leq \varepsilon_0$ given by*

$$\varepsilon_0 := \frac{1}{8(1 + K_{3\alpha})(1 + D)(1 + \|\sigma\|_\infty)(1 + M)^2}, \quad (10.11)$$

then $\Gamma(\mathcal{B}) \subseteq \mathcal{B}$. Moreover, setting

$$L := 2(1 + M)\|\sigma\|_\infty = \frac{2C}{D}, \quad (10.12)$$

for any $\mathbf{f} = (f, f^1) \in \mathcal{B}$ we have

$$\max\{\|\delta f\|_\alpha, \|f^1\|_\infty\} \leq L. \quad (10.13)$$

Proof. Let $\mathbf{f} \in \mathcal{B}$. Setting $\varepsilon := T^\alpha$, if $\varepsilon \leq \varepsilon_0$ then in particular

$$\varepsilon C \leq \frac{\|\sigma\|_\infty}{8(1 + K_{3\alpha})(1 + \|\sigma\|_\infty)(1 + M)} \leq \frac{\|\sigma\|_\infty}{8}.$$

We obtain

$$\|f^1\|_\infty \leq |\sigma(Z_0)| + \varepsilon \|\delta f^1\|_\alpha \leq \|\sigma\|_\infty + \varepsilon_0 [\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq 2\|\sigma\|_\infty \leq L,$$

since $\varepsilon_0 4C \leq \|\sigma\|_\infty$. Similarly

$$\begin{aligned} \|\delta f\|_\alpha &\leq \varepsilon \|f^{[2]}\|_{2\alpha} + \|f^1\|_\infty \|\mathbb{X}^1\|_\alpha \leq \varepsilon_0 4C + (\|\sigma\|_\infty + \varepsilon_0 4C)M \\ &= \varepsilon_0 4C(1+M) + \|\sigma\|_\infty M \leq 2(1+M)\|\sigma\|_\infty = L. \end{aligned}$$

Therefore (10.13) is proved.

We prove now that $\Gamma(\mathbf{f}) \in \mathcal{B}$. We recall that $\Gamma(\mathbf{f}) = (J, \sigma(f))$, where J is uniquely determined by (10.8). By (9.9)

$$[\Gamma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq 2(1+M)(|\nabla \sigma(Z_0) \sigma(Z_0)| + \varepsilon(1+K_{3\alpha})[\sigma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}).$$

By (10.7) and (10.13) we obtain

$$[\Gamma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq 2(1+M)(D\|\sigma\|_\infty + \varepsilon(1+K_{3\alpha})D([\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} + 2L^2)).$$

Now $(1+M)D\|\sigma\|_\infty = C$, and

$$D([\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} + 2L^2) \leq D\left(4C + 2\frac{4C^2}{D^2}\right) \leq 8C\left(D + \frac{C}{D}\right).$$

Note that

$$D + \frac{C}{D} = D + (1+M)\|\sigma\|_\infty \leq (1+M)(1+D)(1+\|\sigma\|_\infty), \quad (10.14)$$

so that by (10.11)

$$[\Gamma(\mathbf{f})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq 2C + 2C = 4C.$$

Therefore, $\Gamma(\mathbf{f}) \in \mathcal{B}$. □

10.3. LOCAL LIPSCHITZ CONTINUITY

We suppose that σ is of class C^3 , with $\|\sigma\|_\infty + \|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty + \|\nabla^3 \sigma\|_\infty < +\infty$ and we fix $D > 0$ such that

$$D \geq \|\nabla \sigma\|_\infty + \|\nabla^2 \sigma\|_\infty + \|\nabla^3 \sigma\|_\infty.$$

LEMMA 10.5. (LOCAL LIPSCHITZ ESTIMATE) *If $T^\alpha \in]0, \varepsilon_0]$ where ε_0 is as in (10.11), then for $\mathbf{f}, \bar{\mathbf{f}} \in \mathcal{B}$, with $\mathcal{B}$ defined in (10.9), we have the local Lipschitz estimate*

$$[\sigma(\mathbf{f}) - \sigma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes (\mathbb{R}^d)^*)} \leq (2 + D + \|\sigma\|_\infty) [\mathbf{f} - \bar{\mathbf{f}}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \quad (10.15)$$

Proof. By Lemma 10.4 we have for $\mathbf{f} = (f, f^1)$, $\bar{\mathbf{f}} = (\bar{f}, \bar{f}^1)$

$$\max \{ \|\delta f\|_\alpha, \|\delta \bar{f}\|_\alpha, \|\bar{f}^1\|_\infty \} \leq L,$$

with L as in (10.12). Now, we want to estimate

$$\begin{aligned} [\sigma(\mathbf{f}) - \sigma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes (\mathbb{R}^d)^*)} &= \underbrace{\|\delta(\nabla\sigma(f) f^1 - \nabla\sigma(\bar{f}) \bar{f}^1)\|_{\alpha}}_A \\ &\quad + \underbrace{\|\sigma(\mathbf{f})^{[2]} - \sigma(\bar{\mathbf{f}})^{[2]}\|_{2\alpha}}_B. \end{aligned}$$

We set $\Delta := f - \bar{f}$, $\Delta^1 := f^1 - \bar{f}^1$, $\Delta^{[2]} := f^{[2]} - \bar{f}^{[2]}$. We first estimate A :

$$\begin{aligned} |\delta(\nabla\sigma(f) f^1 - \nabla\sigma(\bar{f}) \bar{f}^1)_{st}| &= \\ &= |\delta(\nabla\sigma(f))_{st} f_t^1 + \nabla\sigma(f_s) \delta f_{st}^1 - \delta(\nabla\sigma(\bar{f}))_{st} \bar{f}_t^1 - \nabla\sigma(\bar{f}_s) \delta \bar{f}_{st}^1| \\ &\leq |\delta(\nabla\sigma(f) - \nabla\sigma(\bar{f}))_{st} f_t^1| + |\delta(\nabla\sigma(\bar{f}))_{st} (f_t^1 - \bar{f}_t^1)| + \\ &\quad + |(\nabla\sigma(f_s) - \nabla\sigma(\bar{f}_s)) \delta f_{st}^1| + |\nabla\sigma(\bar{f}_s) (\delta f - \delta \bar{f})_{st}|. \end{aligned}$$

By Lemma 2.8 and (1.40) we have for $\varepsilon = T^\alpha$

$$\begin{aligned} A &\leq D[\|f^1\|_{\infty}(\|\delta\Delta\|_{\alpha} + (\|\delta f\|_{\alpha} + \|\delta\bar{f}\|_{\alpha})\|\Delta\|_{\infty}) + \|\delta\bar{f}\|_{\alpha}\|\Delta^1\|_{\infty} + \\ &\quad + \|\Delta\|_{\infty}\|\delta f^1\|_{\alpha} + \|\delta\Delta^1\|_{\alpha}] \\ &\leq D[(\|\delta f\|_{\alpha} + \|\delta\bar{f}\|_{\alpha})\|f^1\|_{\infty} + \|\delta f^1\|_{\alpha}\|\Delta\|_{\infty} + \|f^1\|_{\infty}\|\delta\Delta\|_{\alpha} + \\ &\quad + (1 + \varepsilon\|\delta\bar{f}\|_{\alpha})\|\delta\Delta^1\|_{\alpha}] \\ &\leq D[(2L^2 + \|\delta f^1\|_{\alpha})\|\Delta\|_{\infty} + L\|\delta\Delta\|_{\alpha} + (1 + \varepsilon L)\|\delta\Delta^1\|_{\alpha}] \end{aligned}$$

We show now that

$$\begin{aligned} B &\leq D((\|f^{[2]}\|_{2\alpha} + 3\|\delta f\|_{\alpha}^2)\|\Delta\|_{\infty} + (\|\delta f\|_{\alpha} + \|\delta\bar{f}\|_{\alpha})\|\delta\Delta\|_{\alpha} + \|\Delta^{[2]}\|_{2\alpha}) \\ &\leq D((\|f^{[2]}\|_{2\alpha} + 3L^2)\|\Delta\|_{\infty} + 2L\|\delta\Delta\|_{\alpha} + \|\Delta^{[2]}\|_{2\alpha}). \end{aligned} \quad (10.16)$$

We have by (10.2)

$$\begin{aligned} B &\leq \|\nabla\sigma(f) f^{[2]} - \nabla\sigma(\bar{f}) \bar{f}^{[2]}\|_{2\alpha} + \\ &\quad + \int_0^1 \|\nabla^2\sigma(f + u\delta f) \delta f \otimes \delta f - \nabla^2\sigma(\bar{f} + u\delta\bar{f}) \delta\bar{f} \otimes \delta\bar{f}\|_{2\alpha} du. \end{aligned}$$

With the usual estimate $|ab - \bar{a}\bar{b}| \leq |a - \bar{a}||b| + |\bar{a}||b - \bar{b}|$ we can write

$$\begin{aligned} &\|\nabla\sigma(f) f^{[2]} - \nabla\sigma(\bar{f}) \bar{f}^{[2]}\|_{2\alpha} \leq \\ &\leq \|\nabla\sigma(f) - \nabla\sigma(\bar{f})\|_{\infty} \|f^{[2]}\|_{2\alpha} + \|\nabla\sigma(\bar{f})\|_{\infty} \|\Delta^{[2]}\|_{2\alpha} \\ &\leq \|\nabla^2\sigma\|_{\infty} \|\Delta\|_{\infty} \|f^{[2]}\|_{2\alpha} + \|\nabla\sigma\|_{\infty} \|\Delta^{[2]}\|_{2\alpha} \\ &\leq D(\|\Delta\|_{\infty} \|f^{[2]}\|_{2\alpha} + \|\Delta^{[2]}\|_{2\alpha}). \end{aligned}$$

For the other term

$$\begin{aligned} &\int_0^1 \|\nabla^2\sigma(f + u\delta f) \cdot \delta f \otimes \delta f - \nabla^2\sigma(\bar{f} + u\delta\bar{f}) \cdot \delta\bar{f} \otimes \delta\bar{f}\|_{2\alpha} du \leq \\ &\leq \|\nabla^3\sigma\|_{\infty} \|\delta f\|_{\alpha}^2 (\|\Delta\|_{\infty} + \|\delta\Delta\|_{\infty}) + \|\nabla^2\sigma\|_{\infty} (\|\delta f\|_{\alpha} + \|\delta\bar{f}\|_{\alpha}) \|\delta\Delta\|_{\alpha} \\ &\leq D(\|\delta f\|_{\alpha}^2 (\|\Delta\|_{\infty} + \|\delta\Delta\|_{\infty}) + (\|\delta f\|_{\alpha} + \|\delta\bar{f}\|_{\alpha}) \|\delta\Delta\|_{\alpha}). \end{aligned}$$

Recalling that $\|\delta\Delta\|_{\infty} \leq 2\|\Delta\|_{\infty}$, we have finished the proof of (10.16).

Since $\Delta_0 = f_0 - \bar{f}_0 = 0$, we have $\|\Delta\|_\infty \leq \varepsilon \|\delta\Delta\|_\alpha$. Summing up, we obtain

$$\begin{aligned} & [\sigma(\mathbf{f}) - \sigma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes (\mathbb{R}^d)^*)} = A + B \\ & \leq \{(3L + \varepsilon(5L^2 + [\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}))\|\delta\Delta\|_\alpha + (1 + \varepsilon L)[\mathbf{f} - \bar{\mathbf{f}}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}\}. \end{aligned}$$

On the other hand

$$\begin{aligned} \|\delta\Delta\|_\alpha & \leq \varepsilon \|\Delta^{[2]}\|_{2\alpha} + \|\Delta^1\|_\infty \|\mathbb{X}^1\|_\alpha \\ & \leq \varepsilon \|\Delta^{[2]}\|_{2\alpha} + \varepsilon M \|\delta\Delta^1\|_\alpha \\ & \leq \varepsilon(1 + M)[\mathbf{f} - \bar{\mathbf{f}}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}. \end{aligned}$$

Therefore

$$[\sigma(\mathbf{f}) - \sigma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes (\mathbb{R}^d)^*)} \leq (\varepsilon(1 + M)c_1 + c_2)[\mathbf{f} - \bar{\mathbf{f}}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)},$$

where we set

$$c_1 := D(3L + \varepsilon([\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} + 5L^2)), \quad c_2 := D(1 + \varepsilon L).$$

Since $[\mathbf{f}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq 4C$ we obtain, recalling that $DL = 2C$ by (10.12),

$$\begin{aligned} c_1 & \leq D(3L + \varepsilon(4C + 5L^2)) \leq 6C + 20\varepsilon C \left(D + \frac{C}{D} \right) \\ & \leq 6C + 20\varepsilon C(1 + D)(1 + \|\sigma\|_\infty)(1 + M) \\ & \leq 6C + 3C = 9C, \end{aligned}$$

where we have used first (10.14) and then (10.10)-(10.11). Similarly

$$\varepsilon(1 + M)c_1 \leq 9\varepsilon C(1 + M) = 9\varepsilon D\|\sigma\|_\infty(1 + M)^2 \leq 2,$$

and

$$c_2 = D + \varepsilon DL = D + 2\varepsilon C \leq D + \|\sigma\|_\infty.$$

Therefore

$$\varepsilon(1 + M)c_1 + c_2 \leq 2 + D + \|\sigma\|_\infty.$$

The proof is finished. $\square$

10.4. CONTRACTION

In this section we prove local existence by means of a fixed point argument, assuming σ to be of class C^3 and bounded with its first, second and third derivatives, namely $\|\sigma\|_\infty + \|\nabla\sigma\|_\infty + \|\nabla^2\sigma\|_\infty + \|\nabla^3\sigma\|_\infty < +\infty$. Therefore the assumptions are stronger than for the discrete approximation of Section 3.9. However this method has the advantage of not requiring compactness of the image of Γ and therefore this approach works also for rough equations with values in infinite-dimensional spaces.

Let us fix $D > 0$ such that

$$D \geq \max \{ \|\nabla\sigma\|_\infty, \|\nabla^2\sigma\|_\infty, \|\nabla^3\sigma\|_\infty \}.$$

Recalling that $\mathcal{B}$ was defined in (10.9), we can now show the following

LEMMA 10.6. *If $T^\alpha \in]0, \varepsilon_0]$ where ε_0 is as in (10.11), then $\Gamma: \mathcal{B} \rightarrow \mathcal{B}$ is a contraction for $\|\cdot\|_{\mathcal{D}_{\mathbb{X}}^{2\alpha}}$.*

Proof. Let $\mathbf{f} = (f, f^1)$ and $\bar{\mathbf{f}} = (\bar{f}, \bar{f}^1)$ be in $\mathcal{B}$. Since $f_0 = \bar{f}_0$ and $f_0^1 = \bar{f}_0^1$, by the definitions, see in particular (9.7),

$$\|\Gamma(\mathbf{f}) - \Gamma(\bar{\mathbf{f}})\|_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} = [\Gamma(\mathbf{f}) - \Gamma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}.$$

We set $\varepsilon := T^\alpha$. By (9.9)

$$[\Gamma(\mathbf{f}) - \Gamma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq \varepsilon 2(1+M)(1+K_{3\alpha}) [\sigma(\mathbf{f}) - \sigma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}.$$

Now by Lemma 10.5

$$[\sigma(\mathbf{f}) - \sigma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k \otimes (\mathbb{R}^d)^*)} \leq (2+D+\|\sigma\|_\infty) [\mathbf{f} - \bar{\mathbf{f}}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)}.$$

Now $2+D+\|\sigma\|_\infty \leq 2(1+D)(1+\|\sigma\|_\infty)$. Therefore

$$[\Gamma(\mathbf{f}) - \Gamma(\bar{\mathbf{f}})]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)} \leq c_4 [\mathbf{f} - \bar{\mathbf{f}}]_{\mathcal{D}_{\mathbb{X}}^{2\alpha}(\mathbb{R}^k)},$$

with

$$c_4 = \varepsilon 2(1+M)(1+K_{3\alpha}) 2(1+D)(1+\|\sigma\|_\infty) \leq \frac{1}{2}$$

by (10.11). This concludes the proof. $\square$

CHAPTER 11

GEOMETRIC ROUGH PATHS

11.1. GEOMETRIC ROUGH PATHS

We recall that the set of smooth paths C^1 is not dense in $\mathcal{C}^\alpha$, but its closure is quite large, because it contains $\mathcal{C}^{\alpha'}$ for all $\alpha' > \alpha$ (see Theorem 7.11). The situation is different for rough paths: the set $\mathcal{R}_{1,d}$ of canonical rough paths over smooth paths is again not dense in $\mathcal{R}_{\alpha,d}$, but its closure is a significantly smaller set, that we now describe.

DEFINITION 11.1. *The closure of $\mathcal{R}_{1,d}$ in $\mathcal{R}_{\alpha,d}$ for $\alpha \in]\frac{1}{3}, 1]$ is denoted by $\mathcal{R}_{\alpha,d}^g$ and its elements are called geometric rough paths.*

For smooth paths $f, g \in C^1$, the integration by parts formula holds:

$$\int_s^t f(u) \, dg(u) = f(t)g(t) - f(s)g(s) - \int_s^t g(u) \, df(u).$$

It follows that

$$\int_s^t (f(u) - f(s)) \, dg(u) + \int_s^t (g(u) - g(s)) \, df(u) = (f(t) - f(s))(g(t) - g(s)).$$

We have seen in Proposition 7.8 that the same formula holds if $(f, g) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ with $\alpha + \beta > 1$ and the integral is in the Young sense.

Given a smooth path $X \in C^1$, define $\mathbb{X}^2$ by (8.26) as an ordinary integral (i.e. $(\mathbb{X}^1, \mathbb{X}^2)$ is the canonical rough path over X). The previous relation for $f = X_i$ and $g = X_j$ shows that

$$(\mathbb{X}_{st}^2)^{ij} + (\mathbb{X}_{st}^2)^{ji} = (\mathbb{X}_{st}^1)^i (\mathbb{X}_{st}^1)^j. \quad (11.1)$$

This relation is called the *shuffle relation*: for $i = j$ it identifies $(\mathbb{X}^2)^{ii}$ in terms of $(\mathbb{X}^1)^i$:

$$(\mathbb{X}_{st}^2)^{ii} = \frac{1}{2} ((\mathbb{X}_{st}^1)^i)^2, \quad (11.2)$$

while for $i \neq j$ it expresses $(\mathbb{X}^2)^{ij}$ in terms of $(\mathbb{X}^1)^i, (\mathbb{X}^1)^j, (\mathbb{X}^2)^{ji}$. Denoting by $\text{Sym}(\mathbb{X}^2) := \frac{1}{2} (\mathbb{X}^2 + (\mathbb{X}^2)^T)$ the symmetric part of $\mathbb{X}^2$, we can rewrite the shuffle relation more compactly as follows:

$$\text{Sym}(\mathbb{X}^2) = \frac{1}{2} \mathbb{X}^1 \otimes \mathbb{X}^1. \quad (11.3)$$

DEFINITION 11.2. *Rough paths in $\mathcal{R}_{\alpha,d}$ that satisfy the shuffle relation (11.1)-(11.3) are called weakly geometric and denoted by $\mathcal{R}_{\alpha,d}^{\text{wg}}$.*

Exercise 11.1. For $\alpha > \frac{1}{2}$ we have $\mathcal{R}_{\alpha,d} = \mathcal{R}_{\alpha,d}^{\text{wg}}$ (every rough path is weakly geometric).

We can now show that the closure of $\mathcal{R}_{1,d}$ in $\mathcal{R}_{\alpha,d}$ is included in $\mathcal{R}_{\alpha,d}^{\text{wg}}$.

LEMMA 11.3. *Geometric rough paths are weakly geometric: $\mathcal{R}_{\alpha,d}^g \subset \mathcal{R}_{\alpha,d}^{\text{wg}}$ for any $\alpha \in (\frac{1}{3}, 1)$, with a strict inclusion.*

Proof. Canonical rough paths $(\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{1,d}$ over smooth paths satisfy the shuffle relation (11.1)-(11.3). Geometric rough paths are by definition limits in $\mathcal{R}_{\alpha,d}$ of smooth paths in $\mathcal{R}_{1,d}$. Since convergence in $\mathcal{R}_{\alpha,d}$ implies pointwise convergence, geometric rough paths satisfy the shuffle relation too. This shows that $\mathcal{R}_{\alpha,d}^g \subset \mathcal{R}_{\alpha,d}^{\text{wg}}$.

To prove that the inclusion $\mathcal{R}_{\alpha,d}^g \subset \mathcal{R}_{\alpha,d}^{\text{wg}}$ is strict, it suffices to consider a weakly geometric rough path $(\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha,d}^{\text{wg}}$ which lies above a path $X \in \mathcal{C}^\alpha$ which is not in the closure of $\mathcal{C}^1$. Such a path is not geometric (recall that $(\mathbb{X}_n^1, \mathbb{X}_n^2) \rightarrow (\mathbb{X}^1, \mathbb{X}^2)$ in $\mathcal{R}_{\alpha,d}$ implies $\mathbb{X}_n^1 \rightarrow \mathbb{X}^1$ in $\mathcal{C}_2^\alpha$).

To prove the existence of such a rough path, in the one-dimensional case $d = 1$ it is enough to consider the one provided by (8.19), which is by construction weakly geometric, since the shuffle relation reduces to $\mathbb{X}_{st}^2 := \frac{1}{2}(\mathbb{X}_{st}^1)^2$. $\square$

Although the inclusion $\mathcal{R}_{\alpha,d}^g \subset \mathcal{R}_{\alpha,d}^{\text{wg}}$ is strict, what is left out turns out to be not so large. More precisely, recalling that $\mathcal{R}_{\alpha,d}^g$ is the closure of $\mathcal{R}_{1,d}$ in $\mathcal{R}_{\alpha,d}$, we have a result which is similar to what happens for Hölder spaces, with the important difference that the whole space $\mathcal{R}_{\alpha,d}$ is replaced by $\mathcal{R}_{\alpha,d}^{\text{wg}}$. The proof is non-trivial and we omit it.

PROPOSITION 11.4. *For any $\frac{1}{3} < \alpha' < \alpha < 1$ one has $\mathcal{R}_{\alpha,d}^{\text{wg}} \subseteq \mathcal{R}_{\alpha',d}^g$. This means that for any $\mathbb{X} \in \mathcal{R}_{\alpha,d}^{\text{wg}}$ there is a sequence $\mathbb{X}_n \in \mathcal{R}_{1,d}$ such that $\mathbb{X}_n \rightarrow \mathbb{X}$ in $\mathcal{R}_{\alpha',d}$.*

We stress that the notion of “weakly geometric” rough path depends only on the function $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$, but the notion of “geometric” rough path depends also on the chosen space $\mathcal{R}_{\alpha,d}$. Given a weakly geometric rough path $\mathbb{X} \in \mathcal{R}_{\alpha,d}$, even though $\mathbb{X}$ may fail to be geometric in $\mathcal{R}_{\alpha,d}$, it is certainly geometric in $\mathcal{R}_{\alpha',d}$ for all $\alpha' < \alpha$. In this sense, *every weakly geometric rough path is a geometric rough path, of a possibly slightly lower regularity.*

Finally we note the following

PROPOSITION 11.5. *Let $\alpha \in (\frac{1}{2}, 1)$ and $X \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$. The canonical α -rough path constructed in Lemma 8.16 is geometric.*

Proof. We recall that by the Chen relation

$$\delta(\mathbb{X}^2)_{sut}^{ij} = \delta X_{su}^i \delta X_{ut}^j, \quad \delta(\mathbb{X}^2)_{sut}^{ji} = \delta X_{su}^j \delta X_{ut}^i,$$

so that

$$\delta[(\mathbb{X}^2)^{ij} + (\mathbb{X}^2)^{ji}]_{sut} = \delta X_{su}^i \delta X_{ut}^j + \delta X_{su}^j \delta X_{ut}^i.$$

On the other hand by a simple computation

$$\delta[\delta X^i \delta X^j]_{sut} = \delta X_{su}^i \delta X_{ut}^j + \delta X_{su}^j \delta X_{ut}^i.$$

Therefore $(\mathbb{X}^2)^{ij} + (\mathbb{X}^2)^{ji} - \delta X^i \delta X^j = \delta f$ for some $f \in C_1$ such that $\delta f \in C_2^{2\alpha}$. Since $2\alpha > 1$, we obtain that $\delta f \equiv 0$. $\square$

Note that Proposition 11.5 can be seen as a consequence of the integration by parts formula satisfied by the Young integral, see Proposition 7.8.

11.2. THE STRATONOVICH ROUGH PATH

Let $(B_t)_{t \geq 0}$ be a d -dimensional Brownian motion. We have seen in Theorem 4.2 that the $\mathbb{B} = (\mathbb{B}^1, \mathbb{B}^2)$, defined by

$$\mathbb{B}_{st}^1 := \delta B_{st}, \quad \mathbb{B}_{st}^2 := \int_s^t \mathbb{B}_{sr}^1 \otimes dB_r, \quad 0 \leq s \leq t \leq T,$$

with an Itô integral, yields a.s. an α -rough path for all $\alpha \in (\frac{1}{3}, \frac{1}{2})$, that we can call the *Itô rough path*. As in Section 5.1 we can modify this definition and set

$$\bar{\mathbb{B}}_{st}^1 := \delta B_{st}, \quad \bar{\mathbb{B}}_{st}^2 := \int_s^t \bar{\mathbb{B}}_{sr}^1 \otimes \circ dB_r, \quad 0 \leq s \leq t \leq T,$$

where $\circ$ denotes Stratonovich integration, namely

$$\bar{\mathbb{B}}_{st}^1 := \delta B_{st}, \quad \bar{\mathbb{B}}_{st}^2 := \int_s^t (B_r - B_s) \otimes dB_r + \frac{t-s}{2} I, \quad 0 \leq s \leq t \leq T,$$

with I the identity matrix in $\mathbb{R}^d \otimes \mathbb{R}^d$. By Lemma 8.17, $\bar{\mathbb{B}} = (\bar{\mathbb{B}}^1, \bar{\mathbb{B}}^2)$ defines a α -rough path for all $\alpha \in (\frac{1}{3}, \frac{1}{2})$, that we call the *Stratonovich rough path*. Now we show that $\bar{\mathbb{B}}$ is geometric. We recall that the integration by parts formula reads in this case

$$B_t^i B_t^j - B_s^i B_s^j = \int_s^t B_r^i \circ dB_r^j + \int_s^t B_r^j \circ dB_r^i, \quad 0 \leq s \leq t.$$

Moreover

$$\int_s^t B_s^i \circ dB_r^j = B_s^i (B_t^j - B_s^j).$$

Therefore

$$\begin{aligned} (\bar{\mathbb{B}}_{st}^2)^{ij} + (\bar{\mathbb{B}}_{st}^2)^{ji} &= B_t^i B_t^j - B_s^i B_s^j - B_s^i (B_t^j - B_s^j) - B_s^j (B_t^i - B_s^i) \\ &= (B_t^i - B_s^i)(B_t^j - B_s^j) = [\bar{\mathbb{B}}_{st}^1 \otimes \bar{\mathbb{B}}_{st}^1]^{ij}. \end{aligned}$$

As in the remark following Proposition 11.5, also in the case of the Stratonovich rough path an integration by parts formula is at the heart of the geometric property.

On the other hand, the Itô rough path is *not* geometric, since the integration by parts formula with Itô integrals reads for $i = j$

$$(B_t^i)^2 - (B_s^i)^2 = 2 \int_s^t B_r^i dB_r^i + (t-s), \quad 0 \leq s \leq t,$$

and moreover we have

$$\int_s^t B_s^i dB_r^i = B_s^i (B_t^i - B_s^i).$$

Therefore by the definition of $\mathbb{B}_{st}^2$

$$\begin{aligned} 2(\mathbb{B}_{st}^2)^{ii} &= (B_t^i)^2 - (B_s^i)^2 - 2B_s^i (B_t^i - B_s^i) - (t - s) \\ &= (B_t^i - B_s^i)^2 - (t - s) \\ &= [\mathbb{B}_{st}^1 \otimes \mathbb{B}_{st}^1]^{ii} - (t - s) \\ &\neq [\mathbb{B}_{st}^1 \otimes \mathbb{B}_{st}^1]^{ii}. \end{aligned}$$

Note that for $i \neq j$ we do obtain $(\mathbb{B}_{st}^2)^{ij} + (\mathbb{B}_{st}^2)^{ji} = [\mathbb{B}_{st}^1 \otimes \mathbb{B}_{st}^1]^{ij}$.

11.3. NON-GEOMETRIC ROUGH PATHS

We next consider generic rough paths. These cannot be approximated by canonical rough paths over smooth paths. However we have

LEMMA 11.6. *Given an arbitrary rough path $(\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha, d}$ lying above X , there is always a weakly geometric rough path $(\mathbb{X}^1, \tilde{\mathbb{X}}^2) \in \mathcal{R}_{\alpha, d}^{\text{wg}}$ lying above the same path X .*

Proof. It suffice to define $\tilde{\mathbb{X}}_{ij}^2 := \mathbb{X}_{ij}^2$ for all $i > j$ and use the shuffle relation to define the remaining entries of $\tilde{\mathbb{X}}^2$, i.e. $\tilde{\mathbb{X}}_{ii}^2 := \frac{1}{2}(\mathbb{X}_i^1)^2$ and $\tilde{\mathbb{X}}_{ij}^2 := \mathbb{X}_i^1 \mathbb{X}_j^1 - \mathbb{X}_{ji}^2$ for all $i < j$. In this way $(\mathbb{X}^1, \tilde{\mathbb{X}}^2)$ satisfies the shuffle relation by construction and it is easy to check that $\tilde{\mathbb{X}}^2 \in C_2^{2\alpha}$.

It remains to prove that the Chen relation (8.21) holds for $(\mathbb{X}^1, \tilde{\mathbb{X}}^2)$, that is

$$\delta \tilde{\mathbb{X}}_{ij}^2(s, u, t) = \mathbb{X}_i^1(s, u) \mathbb{X}_j^1(u, t).$$

If $i > j$ this holds because $\tilde{\mathbb{X}}_{ij}^2 = \mathbb{X}_{ij}^2$, so we only need to consider $i = j$ and $i < j$. Note that if we define $A_{st} := \delta f_{st} \delta g_{st}$, for arbitrary $f, g: [a, b] \rightarrow \mathbb{R}$, we have

$$\begin{aligned} \delta A_{sut} &= \delta f_{st} \delta g_{st} - \delta f_{su} \delta g_{su} - \delta f_{ut} \delta g_{ut} \\ &= (\delta f_{su} + \delta f_{ut}) \delta g_{st} - \delta f_{su} \delta g_{su} - \delta f_{ut} \delta g_{ut} \\ &= \delta f_{su} \delta g_{ut} + \delta g_{su} \delta f_{ut}. \end{aligned}$$

Applying this to $f = X^i$ and $g = X^j$ yields, for $i < j$,

$$\begin{aligned} \delta \tilde{\mathbb{X}}_{ij}^2(s, u, t) &= \delta(\mathbb{X}_i^1 \mathbb{X}_j^1 - \mathbb{X}_{ji}^2)(s, u, t) \\ &= \mathbb{X}_i^1(s, u) \mathbb{X}_j^1(u, t) + \mathbb{X}_j^1(s, u) \mathbb{X}_i^1(u, t) - \mathbb{X}_j^1(s, u) \mathbb{X}_i^1(u, t) \\ &= \mathbb{X}_i^1(s, u) \mathbb{X}_j^1(u, t). \end{aligned}$$

Similarly, choosing $f = g = X_i$ gives $\delta \tilde{\mathbb{X}}_{ii}^2(s, u, t) = \mathbb{X}_i^1(s, u) \mathbb{X}_i^1(u, t)$. $\square$

As a corollary, we obtain a useful approximation result.

PROPOSITION 11.7. *For any rough path $(\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha,d}$, there is a function $f \in \mathcal{C}^{2\alpha}([0, T]; \mathbb{R}^d \otimes \mathbb{R}^d)$ and a sequence of canonical rough paths over smooth paths $(\mathbb{X}_n^1, \mathbb{X}_n^2) \in \mathcal{R}_{1,d}$ such that*

$$(\mathbb{X}_n^1, \mathbb{X}_n^2 + \delta f) \rightarrow (\mathbb{X}^1, \mathbb{X}^2) \quad \text{in } \mathcal{R}_{\alpha',d}, \quad \forall \alpha' \in \left(\frac{1}{3}, \alpha\right).$$

Proof. By Lemma 11.6 there is a weakly geometric rough path $(\mathbb{X}^1, \tilde{\mathbb{X}}^2)$ lying above the same path X . Then $\mathbb{X}^2 - \tilde{\mathbb{X}}^2 = \delta f$ for some $f \in \mathcal{C}^{2\alpha}([0, T]; \mathbb{R}^d \otimes \mathbb{R}^d)$, by Lemma 8.17. By Proposition 11.4, there is a sequence $(\mathbb{X}_n^1, \mathbb{X}_n^2) \in \mathcal{R}_{1,d}$ such that $(\mathbb{X}_n^1, \mathbb{X}_n^2) \rightarrow (\mathbb{X}^1, \tilde{\mathbb{X}}^2)$ in $\mathcal{R}_{\alpha',d}$, for any $\alpha' < \alpha$. It follows that $(\mathbb{X}_n^1, \mathbb{X}_n^2 + \delta f) \rightarrow (\mathbb{X}^1, \tilde{\mathbb{X}}^2 + \delta f) = (\mathbb{X}^1, \mathbb{X}^2)$. $\square$

11.4. PURE AREA ROUGH PATHS

Given $X \in \mathcal{C}^\alpha$, we have defined in Definition 3.2 the subset $\mathcal{R}_{\alpha,d}(X)$ of rough paths $(\mathbb{X}^1, \mathbb{X}^2) \in \mathcal{R}_{\alpha,d}$ lying above X , i.e. such that $\mathbb{X}^1 = \delta X$. The case of $\mathbb{X}^1 \equiv 0$ is particularly interesting:

DEFINITION 11.8. *The elements of $\mathcal{R}_{\alpha,d}(0)$, i.e. those of the form $\mathbb{X} = (0, \mathbb{X}^2)$, are called pure area rough paths.*

Pure area rough paths are very explicit. Let us denote by $(\mathbb{R}^{d \times d})^a$ the subspace of $\mathbb{R}^{d \times d}$ given by antisymmetric matrices.

LEMMA 11.9. *$\mathbb{X} = (0, \mathbb{X}^2)$ is a pure area α -rough path if and only if $\mathbb{X}^2 = \delta f$, for some $f \in \mathcal{C}^{2\alpha}([0, T]; \mathbb{R}^{d \times d})$. Such rough path is weakly geometric if and only if $\mathbb{X}_{st}^2 \in (\mathbb{R}^{d \times d})^a$, i.e. $\mathbb{X}_{st}^2$ is an antisymmetric matrix, for all $s, t \in [0, T]_{\leq}^2$; equivalently, we can take $f \in \mathcal{C}^{2\alpha}([0, T]; (\mathbb{R}^{d \times d})^a)$.*

Proof. Since $(0, 0)$ is a rough path, it follows by Lemma 8.17 that for all (pure area) rough paths $(0, \mathbb{X}^2)$ we have $\mathbb{X}^2 = \delta f$ for some $f \in \mathcal{C}^{2\alpha}$. We may assume that $f(0) = 0$ (just redefine $f(t)$ as $f(t) - f(0)$). Since $\mathbb{X}^1 = 0$, the shuffle relation (11.3) becomes $\text{Sym}(\mathbb{X}^2) = 0$, i.e. $\mathbb{X}_{st}^2$ is an antisymmetric matrix. Then $f(t) = f(t) - f(0) = \mathbb{X}_{0t}^2$ is antisymmetric too. $\square$

Note that the set $\mathcal{R}_{\alpha,d}(0)$ of pure area rough paths is a *vector space*, because the Chen relation (8.21) reduces to the linear relation $\delta \mathbb{X}^2 = 0$. Here is the link with general rough paths.

PROPOSITION 11.10. *The set $\mathcal{R}_{\alpha,d}(X)$ of rough paths laying above a given path X is an affine space, with associated vector space $\mathcal{R}_{\alpha,d}(0)$, the space of pure area rough paths.*

Proof. Given rough paths $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ and $\bar{\mathbb{X}} = (\mathbb{X}^1, \bar{\mathbb{X}}^2)$ lying above the same path X , their difference $\mathbb{X} - \bar{\mathbb{X}} = (0, \mathbb{X}^2 - \bar{\mathbb{X}}^2)$ is a pure area rough path, because it satisfies the Chen relation $\delta(\mathbb{X}^2 - \bar{\mathbb{X}}^2) = 0$ (since $\delta \mathbb{X}^2 = \mathbb{X}^1 \otimes \mathbb{X}^1 = \delta \bar{\mathbb{X}}^2$).

Alternatively, Lemma 8.17 yields $\mathbb{X}^2 - \bar{\mathbb{X}}^2 = \delta f$ for some $f \in \mathcal{C}^{2\alpha}$, hence $(0, \mathbb{X}^2 - \bar{\mathbb{X}}^2)$ is a pure area rough path by Lemma 11.9. $\square$

We have seen in Section 8.8 how pure area rough paths can arise concretely as limits of canonical rough paths associated with smooth paths.

11.5. DOSS-SUSSMANN

In this section we suppose that $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ is such that for all $i \in \{1, \dots, k\}$ the $d \times d$ matrix $((\sigma_2)_{j\ell}^i)_{j\ell}$ is symmetric, namely by (3.5) we have the *Frobenius condition*

$$\sum_{a=1}^k \sigma_\ell^a(y) \frac{\partial \sigma_j^i(y)}{\partial y_a} = \sum_{a=1}^k \sigma_j^a(y) \frac{\partial \sigma_\ell^i(y)}{\partial y_a}, \quad \forall y \in \mathbb{R}^k, i \in \{1, \dots, k\}, j, \ell \in \{1, \dots, d\}. \quad (11.4)$$

If we introduce the vector fields $(X_j)_{j=1, \dots, d}$ on $\mathbb{R}^k$:

$$X_j f := \sum_{a=1}^k \sigma_j^a \frac{\partial f}{\partial y_a}, \quad f \in C^\infty(\mathbb{R}^k),$$

then the Frobenius condition (11.4) is equivalent to the commutation relation

$$X_j \circ X_\ell = X_\ell \circ X_j, \quad j, \ell \in \{1, \dots, d\}.$$

Indeed:

$$(X_j \circ X_\ell) f = \sum_{a=1}^k X_j \left(\sigma_\ell^a \frac{\partial f}{\partial y_a} \right) = \sum_{a,b=1}^k \left(\sigma_j^a \sigma_\ell^b \frac{\partial^2 f}{\partial y_a \partial y_b} + \sigma_j^b \frac{\partial \sigma_\ell^a}{\partial y_b} \frac{\partial f}{\partial y_a} \right).$$

The first term in the latter expression is symmetric in j, ℓ , while the second is symmetric for all $f \in C^\infty(\mathbb{R}^k)$ if and only if the Frobenius condition (11.4) holds.

Example 11.11. For $k = d = 2$ we $\sigma: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \otimes (\mathbb{R}^2)^*$

$$\sigma_j^i(y) = \mathbb{1}_{\{i=j\}} y_i,$$

namely

$$\sigma(y) = \begin{pmatrix} y_1 & 0 \\ 0 & y_2 \end{pmatrix}, \quad y = (y_1, y_2) \in \mathbb{R}^2.$$

Then

$$\frac{\partial \sigma_j^i(y)}{\partial y_a} = \mathbb{1}_{\{i=j=a\}},$$

and

$$(\sigma_2)_{j\ell}^i(y) = \sum_{a=1}^2 \frac{\partial \sigma_j^i(y)}{\partial y_a} \sigma_\ell^a(y) = \mathbb{1}_{\{i=j=\ell\}} y_i,$$

which is clearly symmetric in (j, ℓ) . In this case the Frobenius condition (11.4) is satisfied.

Example 11.12. For $k = d = 2$ we consider $\sigma: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \otimes (\mathbb{R}^2)^*$

$$\sigma(y) := \begin{pmatrix} y_2 & 0 \\ 0 & y_1 \end{pmatrix}, \quad y = (y_1, y_2) \in \mathbb{R}^2.$$

In coordinates,

$$\sigma_j^i(y) = \mathbb{1}_{\{i=j=1\}} y_2 + \mathbb{1}_{\{i=j=2\}} y_1.$$

If we compute the partial derivative

$$\frac{\partial \sigma_j^i(y)}{\partial y_a} = \mathbb{1}_{\{i=j \neq a\}}, \quad a \in \{1, 2\},$$

we obtain the expression for σ_2

$$(\sigma_2)_{j\ell}^i(y) = \sum_{a=1}^2 \frac{\partial \sigma_j^i(y)}{\partial y_a} \sigma_\ell^a(y) = \mathbb{1}_{\{i=j \neq \ell\}} y_\ell.$$

Note that σ_2 is not symmetric with respect to (j, ℓ) i.e. $(\sigma_2)_{j\ell}^i \neq (\sigma_2)_{\ell j}^i$, namely it does not satisfy the Frobenius condition (11.4).

If the Frobenius condition (11.4) holds and $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ is a weakly geometric α -rough path, we obtain

$$\begin{aligned} (\sigma_2(y) \mathbb{X}^2)^i &= \sum_{a,b=1}^d (\sigma_2)_{ab}^i(y) (\mathbb{X}^2)^{ba} \\ &= \sum_{a,b=1}^d \frac{1}{2} \{(\sigma_2)_{ab}^i + (\sigma_2)_{ba}^i\}(y) (\mathbb{X}^2)^{ba} \\ &= \frac{1}{2} \sum_{a,b=1}^d (\sigma_2)_{ab}^i(y) \{(\mathbb{X}^2)^{ab} + (\mathbb{X}^2)^{ba}\} \\ &= \frac{1}{2} \sum_{a,b=1}^d (\sigma_2)_{ab}^i(y) (\mathbb{X}^1)^a (\mathbb{X}^1)^b \\ &= \frac{1}{2} (\sigma_2(y) (\mathbb{X}^1 \otimes \mathbb{X}^1))^i. \end{aligned}$$

In this case it turns out that the solution Z to the associated finite difference equation is a function of $\mathbb{X}^1$ alone since (3.19) is equivalent to

$$|Z_{st}^{[3]}| \lesssim |t-s|^{3\alpha}, \quad Z_{st}^{[3]} = \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \frac{1}{2} \sigma_2(Z_s) (\mathbb{X}_{st}^1 \otimes \mathbb{X}_{st}^1). \quad (11.5)$$

Arguing as in the proof of the proof of Theorem 3.11, it can be seen that the map $(Z_0, \mathbb{X}^1) \mapsto Z$ is continuous.

PROPOSITION 11.13. *Let $M > 0$ and let us suppose that $\mathbb{X}$ is a weakly geometric rough path and $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ satisfies the Frobenius condition (11.4). If*

$$\max \{|\sigma(Z_0)| + |\sigma(\bar{Z}_0)| + |\sigma_2(\bar{Z}_0)|, \|\mathbb{X}^1\|_\alpha, \|\bar{\mathbb{X}}^1\|_\alpha\} \leq M,$$

then for every $T > 0$ there are $\hat{\tau}_{M,D,T}, C_{M,D,T} > 0$ such that for $\tau \in]0, \hat{\tau}_{M,D,T}]$

$$\begin{aligned} \|Z - \bar{Z}\|_{\infty, \tau} + \|\delta Z - \delta \bar{Z}\|_{\alpha, \tau} + \|Z^{[2]} - \bar{Z}^{[2]}\|_{2\alpha, \tau} &\leq \\ &\leq C_{M,D,T} (|Z_0 - \bar{Z}_0| + \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_\alpha). \end{aligned}$$

In particular, the solution to (3.19) depends only on $\mathbb{X}^1$ if $\mathbb{X}$ is a geometric rough path.

Proof. The proof follows from the same arguments as in the proof of Theorem 3.11, if one uses the algebraic relations for $Y^{[3]} := Z^{[3]} - \bar{Z}^{[3]}$ and $\delta Y^{[3]} := \delta Z^{[3]} - \delta \bar{Z}^{[3]}$ obtained by replacing $\mathbb{X}^2$ with $\frac{1}{2}\mathbb{X}^1 \otimes \mathbb{X}^1$, as in

$$\begin{aligned} Z_{st}^{[3]} &= \delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \frac{1}{2} \sigma_2(Z_s) (\mathbb{X}_{st}^1 \otimes \mathbb{X}_{st}^1), \\ \delta Z_{sut}^{[3]} &= (\sigma(Z_u) - \sigma(Z_s) - \sigma_2(Z_s) \mathbb{X}_{su}^1) \mathbb{X}_{ut}^1 + \frac{1}{2} \delta \sigma_2(Z)_{su} (\mathbb{X}_{ut}^1 \otimes \mathbb{X}_{ut}^1), \end{aligned}$$

and analogously for $\bar{Z}^{[3]}, \delta \bar{Z}^{[3]}$. One can also note the simple estimate

$$\|\mathbb{X}^1 \otimes \mathbb{X}^1 - \bar{\mathbb{X}}^1 \otimes \bar{\mathbb{X}}^1\|_{2\alpha} \leq \|\mathbb{X}^1 - \bar{\mathbb{X}}^1\|_{\alpha} (\|\mathbb{X}^1\|_{\alpha} + \|\bar{\mathbb{X}}^1\|_{\alpha}).$$

The rest of the proof is identical to that of Theorem 3.11. $\square$

Remark 11.14. Doss and Sussmann prove a continuity result in the sup-norm.

11.6. LACK OF CONTINUITY (AGAIN)

In section 11.5 we have seen that, under appropriate conditions on σ , the map $(Z_0, \mathbb{X}^1) \mapsto Z$ is continuous if $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ varies in the class of weakly geometric rough paths. In this section we show that this is not a general fact, and the continuity result of Proposition 3.11 can not be improved in general.

More precisely, we consider the sequence $\mathbb{X}_n = (\mathbb{X}_n^1, \mathbb{X}_n^2)$ such that $\mathbb{X}_n^1 \rightarrow 0$, $\mathbb{X}_n^2 \rightarrow \mathbb{X}^2 \neq 0$ constructed in Section 8.8 and we provide an explicit $\sigma: \mathbb{R}^k \rightarrow \mathbb{R}^k \otimes (\mathbb{R}^d)^*$ one such that the solution Z^n to the controlled integral equation

$$Z_t^n = Z_0 + \int_0^t \sigma(Z_s^n) \dot{X}_n(s) ds, \quad t \geq 0,$$

is *not* a continuous function of $(Z_0, \mathbb{X}^1)$ (but only of $(Z_0, \mathbb{X}^1, \mathbb{X}^2)$).

We consider $\sigma: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \otimes (\mathbb{R}^2)^*$ as in the Example 11.12:

$$\sigma(y) := \begin{pmatrix} y_2 & 0 \\ 0 & y_1 \end{pmatrix}, \quad y = (y_1, y_2) \in \mathbb{R}^2.$$

Moreover we consider the rough paths $\mathbb{X}_n$ and $\mathbb{X}$ considered in Section 8.8 and given by $\mathbb{X}_{st} = (0, (t-s)J)$, where J is the 2×2 matrix

$$J := \begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix},$$

while $\mathbb{X}_n$ is the canonical rough path associated with the smooth function path $X_n(t) = n^{-\frac{1}{2}}(\cos(nt), \sin(nt))$, $t \geq 0$. Since $X_n \rightarrow 0$ in the sup-norm as $n \rightarrow \infty$, if the map $X \mapsto Z$ were continuous then we would expect Z_n to converge to a constant function $Z \equiv Z_0$, solution of $\dot{Z} = \sigma(Z) \cdot 0$.

Since $\mathbb{X}_n$ converges to $\mathbb{X}$ as a α -rough path for $\alpha \in (\frac{1}{3}, \frac{1}{2})$, the solution Z^n to the difference equation above converges in $\mathcal{C}^a$ to the solution Z to the equation

$$\delta Z_{st} - \sigma(Z_s) \mathbb{X}_{st}^1 - \sigma_2(Z_s) \mathbb{X}_{st}^2 = o(t-s).$$

Since $\mathbb{X}^1 \equiv 0$ and $\mathbb{X}_{st}^2 = (t-s)J$, we obtain that

$$\delta Z_{st} - (t-s)\sigma_2(Z_s)J = o(t-s).$$

Then, recalling that $(\sigma_2)_{j\ell}^i(y) = \mathbb{1}_{\{i=j \neq \ell\}} y_\ell$, we can compute

$$(\sigma_2(y)J)^i = \sum_{a,b=1}^2 (\sigma_2)_{ab}^i(y) J^{ba} = \frac{1}{2}(\mathbb{1}_{\{i=2\}} y_1 - \mathbb{1}_{\{i=1\}} y_2) = (Jy)^i.$$

Therefore

$$\delta Z_{st} - (t-s)JZ_s = o(t-s),$$

which means that

$$\dot{Z}_t = JZ_t \implies Z_t = \exp(tJ)Z_0, \quad t \geq 0.$$

Since we have already shown that $\mathbb{X}^1 = 0$, we obtain that $Z: [0, T] \rightarrow \mathbb{R}^2$, limit of Z^n as $n \rightarrow \infty$, satisfies

$$Z_t = Z_0 + \int_0^t \frac{1}{2} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} Z_s ds.$$

Then if $Z_0 \neq 0$ the solution Z is not constant.

11.7. THE ROUGH INTEGRAL IN THE GEOMETRIC CASE

We have seen in Proposition 7.8 that the Young integral satisfies the classical integration by parts formula. We consider now a weakly geometric rough path $\mathbb{X}$ and two paths $\mathbf{f} = (f, f^1)$, $\mathbf{g} = (g, g^1)$ which are 2α -controlled by $\mathbb{X}$. We set

$$F_t := F_0 + \int_0^t f_s d\mathbb{X}_s, \quad G_t := G_0 + \int_0^t g_s d\mathbb{X}_s, \quad t \geq 0.$$

We want to show that, under the assumption that $\mathbb{X}$ is weakly geometric, an analogous integration by parts formula holds, namely:

$$F_t G_t = F_0 G_0 + \underbrace{\int_0^t F_s g_s d\mathbb{X}_s + \int_0^t G_s f_s d\mathbb{X}_s}_{I_t}.$$

We start by showing that $(F_s g_s, F_s g_s^1 + f_s g_s)_{s \in [0, T]}$ is 2α -controlled by $\mathbb{X}$:

$$\begin{aligned} F_t g_t - F_s g_s &= F_t \delta g_{st} + g_s \delta F_{st} \\ &= F_s \delta g_{st} + g_s \delta F_{st} + \delta F_{st} \delta g_{st} \\ &= (F_s g_s^1 + f_s g_s) \mathbb{X}_{st}^1 + O(|t-s|^{2\alpha}). \end{aligned}$$

The same holds of course for $(f_s G_s, G_s f_s^1 + f_s g_s)_{s \in [0, T]}$. Now we know that I_t is the integral uniquely associated with the germ

$$A_{st} = (F_s g_s + G_s f_s) \mathbb{X}_{st}^1 + (F_s g_s^1 + G_s f_s^1 + 2f_s g_s) \mathbb{X}_{st}^2.$$

By the weakly geometric condition, we have $2\mathbb{X}_{st}^2 = (\mathbb{X}_{st}^1)^2$ and therefore we obtain

$$A_{st} = (F_s g_s + G_s f_s) \mathbb{X}_{st}^1 + (F_s g_s^1 + G_s f_s^1) \mathbb{X}_{st}^2 + f_s g_s (\mathbb{X}_{st}^1)^2.$$

Now we write

$$\begin{aligned} \delta(FG)_{st} &= \delta F_{st} G_t + F_s \delta G_{st} \\ &= G_s \delta F_{st} + F_s \delta G_{st} + \delta F_{st} \delta G_{st} \\ &= (F_s g_s + G_s f_s) \mathbb{X}_{st}^1 + (F_s g_s^1 + G_s f_s^1) \mathbb{X}_{st}^2 + \delta F_{st} \delta G_{st} + O(|t-s|^{3\alpha}). \end{aligned}$$

Now since $\mathbb{X}^2 \in C_2^{2\alpha}$

$$\begin{aligned} \delta F_{st} \delta G_{st} &= (f_s \mathbb{X}_{st}^1 + f_s^1 \mathbb{X}_{st}^2)(g_s \mathbb{X}_{st}^1 + g_s^1 \mathbb{X}_{st}^2) + O(|t-s|^{3\alpha}) \\ &= f_s g_s (\mathbb{X}_{st}^1)^2 + O(|t-s|^{3\alpha}). \end{aligned}$$

Then we obtain that

$$\delta(FG)_{st} = A_{st} + O(|t-s|^{3\alpha}).$$

Since $3\alpha > 1$, it follows that $F_t G_t - F_0 G_0 = I_t$ for all $t \geq 0$.

Example 11.15. It is well known that the Stratonovich stochastic integral satisfies the above integration by parts formula. This section extends this result to all (weakly) geometric rough paths.

CHAPTER 12

ALGEBRA

Let us recall that a d -dimensional α -rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ with $\alpha > \frac{1}{3}$ is such that $\mathbb{X}_{st}$ takes values in $G := \mathbb{R}^d \times (\mathbb{R}^d \otimes \mathbb{R}^d)$ for all $0 \leq s \leq t \leq T$. We want to show that the Chen relation (8.21) has a very natural algebraic interpretation if we endow G with a suitable group structure.

12.1. A NON-COMMUTATIVE GROUP

We denote in the following generic elements $x \in G = \mathbb{R}^d \times (\mathbb{R}^d \otimes \mathbb{R}^d)$ by $x = (x_1, x_2)$ with $x_1 \in \mathbb{R}^d$ and $x_2 \in \mathbb{R}^d \otimes \mathbb{R}^d$. We define an operation $\star: G \times G \rightarrow G$ as follows: for $x, y \in G$ with $x = (x_1, x_2)$ and $y = (y_1, y_2)$ we set

$$x \star y := z = (z_1, z_2), \quad z_1 := x_1 + y_1, \quad z_2 := x_2 + y_2 + x_1 \otimes y_1.$$

It is simple to see that $(G, \star, \mathbf{1})$, is a group, where $\mathbf{1} := (0, 0)$. First associativity of the product:

$$\begin{aligned} (x \star y) \star z &= (x_1 + y_1 + z_1, x_2 + y_2 + z_2 + x_1 \otimes y_1 + (x_1 + y_1) \otimes z_1) \\ &= (x_1 + y_1 + z_1, x_2 + y_2 + z_2 + x_1 \otimes (y_1 + z_1) + y_1 \otimes z_1) \\ &= x \star (y \star z). \end{aligned}$$

Now the fact that $\mathbf{1}$ is the neutral element is obvious. Finally the inverse is given explicitly by

$$x^{\star(-1)} = (-x_1, -x_2 + x_1 \otimes x_1). \quad (12.1)$$

Let us note that $(G, \star, \mathbf{1})$ is non-commutative for $d \geq 2$, since in general $x_1 \otimes y_1 \neq y_1 \otimes x_1$.

Now we want to interpret the Chen relation (8.21) in this setting. Given a α -rough path $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$, we write

$$\mathbb{X}: [0, T]_{\leq}^2 \rightarrow G, \quad \mathbb{X}_{st} := (\mathbb{X}_{st}^1, \mathbb{X}_{st}^2).$$

Then the Chen formula (8.21) yields

$$\mathbb{X}_{st} = \mathbb{X}_{su} \star \mathbb{X}_{ut}, \quad 0 \leq s \leq t \leq T.$$

Indeed it is enough to note that for $0 \leq s \leq u \leq t \leq T$

$$\mathbb{X}_{st}^1 = \mathbb{X}_{su}^1 + \mathbb{X}_{ut}^1, \quad \mathbb{X}_{st}^2 = \mathbb{X}_{su}^2 + \mathbb{X}_{ut}^2 + \mathbb{X}_{su}^1 \otimes \mathbb{X}_{ut}^1.$$

Note that we also have, by the analytical estimates $|\mathbb{X}_{st}^i| \lesssim |t-s|^{i\alpha}$ that $\mathbb{X}_{tt} = \mathbf{1}$.

12.2. SHUFFLE GROUP

We can consider the subset $H \subset G$ given by

$$H := \{x = (x_1, x_2) \in G : x_2 + x_2^T = x_1 \otimes x_1\}, \quad (12.2)$$

where $(a \otimes b)^T := b \otimes a$ for $a, b \in \mathbb{R}^d$.

We can see that H is a subgroup of G : if $x, y \in H$ then $z := x \star y$ satisfies

$$\begin{aligned} z_2 + z_2^T &= x_2 + y_2 + x_1 \otimes y_1 + x_2^T + y_2^T + y_1 \otimes x_1 \\ &= x_1 \otimes x_1 + y_1 \otimes y_1 + x_1 \otimes y_1 + y_1 \otimes x_1 \\ &= (x_1 + y_1) \otimes (x_1 + y_1) = z_1 \otimes z_1. \end{aligned}$$

Moreover if $x \in H$ then its inverse $y = x^{\star(-1)} \in G$ satisfies

$$\begin{aligned} y_2 + y_2^T &= -x_2 + x_1 \otimes x_1 - x_2^T + x_1 \otimes x_1 \\ &= -x_1 \otimes x_1 + 2x_1 \otimes x_1 \\ &= (-x_1) \otimes (-x_1) = y_1 \otimes y_1 \end{aligned}$$

so that $x^{\star(-1)} \in H$. Finally $\mathbf{1} \in H$. Therefore H is indeed a (proper) subgroup of G . Moreover by (12.1) and the relation defining elements of H we have the simpler expression for the inverse

$$x^{\star(-1)} = (-x_1, x_2^T), \quad x \in H. \quad (12.3)$$

Indeed for $x \in H$ we obtain

$$\begin{aligned} (x_1, x_2) \star (-x_1, x_2^T) &= (-x_1, x_2^T) \star (x_1, x_2) \\ &= (x_1 - x_1, x_2 + x_2^T - x_1 \otimes x_1) \\ &= (0, 0). \end{aligned}$$

Therefore by (11.1) we have the following

LEMMA 12.1. *A rough path $\mathbb{X}$ is weakly geometric if and only if the associated map $\mathbb{X}: [0, T]_{\leq}^2 \rightarrow G$ takes values in H .*

12.3. UNORDERED TIMES

As we explained at the beginning of Chapter 8, given $\mathbb{X}^1 = \delta X \in C_2^\alpha$, a choice of $\mathbb{X}^2$ is equivalent to a choice of a generalised integral $I_t = \int_0^t X_s \otimes dX_s$, $t \in [0, T]$, as in Definition 8.1, namely $I: [0, T] \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ such that $I_0 = 0$ and

$$\delta I_{st} - X_s \otimes \delta X_{st} = \mathbb{X}_{st}^2, \quad \mathbb{X}^2 \in C_2^{2\alpha}. \quad (12.4)$$

An interesting feature of the equality in (12.4) is that the left hand side makes sense also for $s > t$, which suggests how to define $\mathbb{X}_{st}^2$ for $s > t$ in a way which is consistent with the properties of $(\mathbb{X}_{st}^2)_{s \leq t}$, in particular the Chen relation.

We define therefore, given $\mathbb{X} = (\mathbb{X}^1, \mathbb{X}^2)$ and I as above, for $s > t$:

$$\begin{aligned}\mathbb{X}_{st}^1 &:= \delta X_{st} = -\mathbb{X}_{ts}^1, \\ \mathbb{X}_{st}^2 &:= \delta I_{st} - X_s \otimes \delta X_{st} = -\mathbb{X}_{ts}^2 - \mathbb{X}_{ts}^1 \otimes \mathbb{X}_{ts}^1,\end{aligned}$$

where the latter equality follows from

$$\delta I_{st} - X_s \otimes \delta X_{st} = -(\delta I_{ts} - X_t \otimes \delta X_{ts}) - \delta X_{ts} \otimes \delta X_{ts}.$$

We can now compute for $s, u, t \in [0, T]$:

$$\begin{aligned}\mathbb{X}_{su} \star \mathbb{X}_{ut} &= (\delta X_{su}, \delta I_{su} - X_s \otimes \delta X_{su}) \star (\delta X_{ut}, \delta I_{ut} - X_u \otimes \delta X_{ut}) \\ &= (\delta X_{st}, \delta I_{st} - X_s \otimes \delta X_{su} - X_u \otimes \delta X_{ut} + \delta X_{su} \otimes \delta X_{ut}) \\ &= (\delta X_{st}, \delta I_{st} - X_s \otimes \delta X_{st}) = \mathbb{X}_{st}.\end{aligned}$$

Therefore with the above definition we obtain the *Chen relation for unordered times*

$$\mathbb{X}_{st} = \mathbb{X}_{su} \star \mathbb{X}_{ut}, \quad \forall s, u, t \in [0, T]. \quad (12.5)$$

Moreover, if $|\mathbb{X}_{st}^1| \lesssim |t - s|^\alpha$ and $|\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}$ for $0 \leq s \leq t \leq T$, we have

$$|\mathbb{X}_{st}^1| \lesssim |t - s|^\alpha, \quad |\mathbb{X}_{st}^2| \lesssim |t - s|^{2\alpha}, \quad \forall s, t \in [0, T].$$

By (12.1) we see also that with this definition we have $\mathbb{X}_{ts} = (\mathbb{X}_{st})^{\star(-1)}$ for all s, t . In particular we obtain that

$$\mathbb{X}_{st} = (\mathbb{X}_{0s})^{\star(-1)} \star \mathbb{X}_{0t},$$

namely we realise that the two-parameter map $\mathbb{X}: [0, T]^2 \rightarrow G$ can be recovered as a *purely algebraic function* of $(\mathbb{X}_{0t})_{0 \leq t \leq T}$. Actually, for any $u \in [0, T]$ we have likewise

$$\mathbb{X}_{st} = (\mathbb{X}_{us})^{\star(-1)} \star \mathbb{X}_{ut}.$$

For this reason, one can look for a map $\mathbb{X}: [0, T] \rightarrow G$ such that

$$\mathbb{X}_{st} := \mathbb{X}_s^{\star(-1)} \star \mathbb{X}_t, \quad s, t \in [0, T]. \quad (12.6)$$

From the above, we see that any map $[0, T] \ni t \mapsto \mathbb{X}_{ut} \in G$, for a fixed $u \in [0, T]$, has this property. On other hand, (12.6) is also equivalent to

$$\mathbb{X}_t = \mathbb{X}_0 \star \mathbb{X}_{0t},$$

and for any $g \in G$ setting $\mathbb{X}_t^g := g \star \mathbb{X}_{0t}$ we have a $\mathbb{X}^g: [0, T] \rightarrow G$ that satisfies (12.6) and $\mathbb{X}_0^g = g$, since $\mathbb{X}_{00} = (0, 0) = \mathbf{1}$, the neutral element in G .

By the definitions, we have $\mathbb{X}_{0t} = (\delta X_{0t}, I_t - X_0 \otimes \delta X_{0t})$, $g = (g_1, g_2)$ and

$$\mathbb{X}_t^g = g \star \mathbb{X}_{0t} = (g_1 + \delta X_{0t}, g_2 + I_t + (g_1 - X_0) \otimes \delta X_{0t}). \quad (12.7)$$

For example a natural choice is for $g = (X_0, 0)$ and in this case $\mathbb{X}_t^g = (X_t, I_t)$. Another natural choice is $g = \mathbf{1} = (0, 0)$ and then $\mathbb{X}_t^g = (\delta X_{0t}, I_t - X_0 \otimes \delta X_{0t})$.

If now $\mathbb{X}$ is geometric, namely if $\mathbb{X}_{st} \in H$ for all s, t as in (12.2), then it is natural to require that $\mathbb{X}_t \in H$ as well for all t . For $\mathbb{X}_t = g \star \mathbb{X}_{0t}$, this is equivalent to $g \in H$. Note that if $X_0 \neq 0$ then $(X_0, 0) \notin H$ and therefore in this case $(X_t, I_t) \notin H$, while $(\delta X_{0t}, I_t - X_0 \otimes \delta X_{0t}) \in H$, which corresponds to the case $g = \mathbf{1}$.

The reason why we discuss this in detail is the following: instead of writing the theory of rough paths in terms of the two-parameter function $(\mathbb{X}_{st})_{0 \leq s \leq t \leq T}$, we may have written everything in terms of a choice of $(\mathbb{X}_t)_{0 \leq t \leq T}$ in G as above. However we must still how the analytical constraints on $(\mathbb{X}_{st})_{0 \leq s \leq t \leq T}$ are expressed in terms of $(\mathbb{X}_t)_{0 \leq t \leq T}$: these can be translated by (12.6) and (12.7) back to

$$|\delta X_{st}| \lesssim |t - s|^\alpha, \quad |\delta I_{st} - X_s \otimes \delta X_{st}| \lesssim |t - s|^{2\alpha}, \quad 0 \leq s \leq t \leq T,$$

namely the property of I being a generalised integral of $X \otimes dX$.

12.4. AN EXAMPLE: THE BROWNIAN CASE

Let consider the Itô Brownian rough paths in $\mathbb{R}^d$

$$\mathbb{B}_{st}^1 = B_t - B_s, \quad \mathbb{B}_{st}^2 = \int_s^t (B_r - B_s) \otimes dB_r, \quad 0 \leq s \leq t \leq T.$$

Note that if $s > t$ we can not use naively this definition for $\mathbb{B}_{st}^2$ since the stochastic integral $\int_t^s (B_s - B_r) \otimes dB_r$ is anticipative, namely $(B_s - B_r)_{t \leq r \leq s}$ is not adapted to the filtration of $(B_r)_{t \leq r \leq s}$, and therefore some care is required. Let us rather apply the algebraic definition $\mathbb{B}_{st} := \mathbb{B}_{ts}^{*(-1)}$ of Section 12.3, namely for $0 \leq t < s \leq T$ we set

$$\begin{aligned} \mathbb{B}_{st}^1 &:= B_t - B_s, \\ \mathbb{B}_{st}^2 &:= - \int_t^s (B_r - B_t) \otimes dB_r + (B_s - B_t) \otimes (B_s - B_t) \\ &= \int_t^s dB_r \otimes (B_r - B_t) + (s - t)I, \end{aligned}$$

where I is the identity matrix of $\mathbb{R}^d$. In other words

$$(\mathbb{B}_{st}^2)^{ij} = \int_t^s dB_r^i (B_r - B_t)^j + (s - t)\mathbb{1}_{(i=j)}.$$

Here a one-parameter function $\mathbb{B}: [0, T] \rightarrow G$ such that $\mathbb{B}_{st} = \mathbb{B}_s^{*(-1)} \star \mathbb{B}_t$ is given by

$$\mathbb{B}_t = \left(B_t, \int_0^t B_r \otimes dB_r \right), \quad t \geq 0.$$

Let us consider now the Stratonovich case:

$$\bar{\mathbb{B}}_{st}^1 = B_t - B_s, \quad \bar{\mathbb{B}}_{st}^2 = \int_s^t (B_r - B_s) \otimes \circ dB_r, \quad 0 \leq s \leq t \leq T.$$

Then we obtain from the definitions of the previous section for $0 \leq t < s \leq T$

$$\bar{\mathbb{B}}_{st}^1 = B_t - B_s,$$

and if one applies (12.3) then we have for $0 \leq t < s \leq T$

$$\bar{\mathbb{B}}_{st}^2 = (\bar{\mathbb{B}}_{ts}^2)^T = \int_t^s \circ dB_r \otimes (B_r - B_t),$$

namely

$$(\bar{\mathbb{B}}_{st}^2)^{ij} = \int_t^s (B_r - B_t)^j \circ dB_r^i.$$

Here a one-parameter function $\bar{\mathbb{B}}: [0, T] \rightarrow H$ such that $\bar{\mathbb{B}}_{st} = \bar{\mathbb{B}}_s^{*(-1)} \star \bar{\mathbb{B}}_t$ is given by

$$\bar{\mathbb{B}}_t = \left(B_t, \int_0^t B_r \otimes \circ dB_r \right), \quad t \geq 0.$$

As discussed at the end of Section 12.3, with this definition $\bar{\mathbb{B}}_t \in H$ for all $t \in [0, T]$ as long as $B_0 = 0$.

12.5. SEWING FOR UNORDERED TIMES

Now, suppose that we have a general germ $A: [0, T]^2 \rightarrow \mathbb{R}$. We suppose that it satisfies for some $\eta > 1$

$$|A_{st} - A_{su} - A_{ut}| \leq C_A(|u - s| \vee |t - u|)^\eta, \quad s, u, t \in [0, T].$$

In particular, the restriction $A: [0, T]_{\leq}^2 \rightarrow \mathbb{R}$ is such that $\delta A: [0, T]_{\leq}^3 \rightarrow \mathbb{R}$ belongs to C_3^η . By the Sewing Lemma, we have a unique choice for (I, R) such that

$$I_0 = 0, \quad \delta I_{st} = A_{st} + R_{st}, \quad |R_{st}| \lesssim |t - s|^\eta, \quad 0 \leq s \leq t \leq T.$$

We want to extend R to a function on $[0, T]^2$ in such a way that the previous formula holds over $[0, T]^2$. We set

$$R_{st} = -A_{st} - A_{ts} - R_{ts}, \quad 0 \leq t \leq s \leq T. \quad (12.8)$$

Since $\delta I_{ts} = -\delta I_{st}$, we have for $t \leq s$

$$R_{st} = -A_{st} - (\delta I_{ts} - R_{ts}) - R_{ts} = -A_{st} - \delta I_{ts} = \delta I_{st} - A_{st},$$

so that $\delta I = A + R$ on $[0, T]^2$. Moreover, since $A_{ss} = 0$ by Remark 6.6,

$$|R_{st}| \leq |(\delta A)_{sts}| + |R_{ts}| \leq (C_\eta + 1) C_A |t - s|^\eta, \quad 0 \leq t \leq s \leq T. \quad (12.9)$$

12.6. CONTROLLED PATHS

We define

$$S := \{\mathbb{1}\} \sqcup \{1, \dots, d\} \sqcup (\{1, \dots, d\} \times \{1, \dots, d\})$$

and $\mathcal{T}$ as the linear span of S . Given $g \in G = \mathbb{R}^d \oplus (\mathbb{R}^d \otimes \mathbb{R}^d)$ we define $\Gamma_x: \mathcal{T} \rightarrow \mathcal{T}$:

$$\Gamma_g \mathbb{1} := \mathbb{1}, \quad \Gamma_g i := i + (g^1)^i \mathbb{1}$$

$$\Gamma_g(i, j) := (i, j) + (g^1)^i j + (g^2)^{ij} \mathbb{1}.$$

Then we have the formula for $x, y \in G$

$$\begin{aligned}\Gamma_g \circ \Gamma_h i &= i + (g^1)^i \mathbb{1} + (h^1)^i \mathbb{1} = i + (g^1 + h^1)^i \mathbb{1} = \Gamma_{h \star g} i \\ \Gamma_g \circ \Gamma_h (i, j) &= (i, j) + (g^1)^i j + (g^2)^{ij} \mathbb{1} + (h^1)^i (j + (g^1)^j \mathbb{1}) + (h^2)^{ij} \mathbb{1} \\ &= (i, j) + (g^1 + h^1)^i j + (g^2 + h^2 + h^1 \otimes g^1)^{ij} \mathbb{1} = \Gamma_{h \star g} (i, j),\end{aligned}$$

namely

$$\Gamma_g \circ \Gamma_h = \Gamma_{h \star g}, \quad \forall g, h \in G,$$

and therefore the map $G \ni g \mapsto \Gamma_g \in \text{End}(\mathcal{T})$ is an anti-morphism of semigroups from $(G, \star)$ to $(\text{End}(\mathcal{T}), \circ)$. Since $\Gamma_{(0,0)} = \mathbb{1}$, we obtain

$$(\Gamma_g)^{-1} = \Gamma_{g^{\star(-1)}}.$$

Given a α -rough path $(\mathbb{X}_{st})_{s,t \in [0,T]}$ on $\mathbb{R}^d$, we set $\Gamma_{st}: \mathcal{T} \rightarrow \mathcal{T}$ as $\Gamma_{\mathbb{X}_{ts}}$, namely

$$\begin{aligned}\Gamma_{st} \mathbb{1} &:= \mathbb{1}, \quad \Gamma_{st} i := i + (\mathbb{X}_{ts}^1)^i \mathbb{1}, \\ \Gamma_{st} (i, j) &:= (i, j) + (\mathbb{X}_{ts}^1)^i j + (\mathbb{X}_{ts}^2)^{ij} \mathbb{1}.\end{aligned}$$

Note the somewhat strange choice of $\mathbb{X}_{ts}$ instead of $\mathbb{X}_{st}$ everywhere, which is due to the anti-morphism property. Then by the Chen relation (12.5)

$$\Gamma_{su} \circ \Gamma_{ut} = \Gamma_{\mathbb{X}_{us}} \circ \Gamma_{\mathbb{X}_{tu}} = \Gamma_{\mathbb{X}_{tu} \star \mathbb{X}_{us}} = \Gamma_{\mathbb{X}_{ts}} = \Gamma_{st}$$

for all $s, u, t \in [0, T]$. In particular, by the above formula for the inverse of Γ_g ,

$$(\Gamma_{st})^{-1} = \Gamma_{ts},$$

which shows the importance of defining $\mathbb{X}_{st}$ for all $s, t \in [0, T]$ and not only for $s \leq t$ (see also below).

Let us consider a controlled path $(Z, Z^1) \in C^\alpha([0, T]) \times C^\alpha([0, T]; (\mathbb{R}^d)^\star)$, namely we suppose that

$$|\delta Z_{st} - Z_s^1 \mathbb{X}_{st}^1| \lesssim |t - s|^{2\alpha},$$

where according to the rule on contraction of tensors we have

$$Z_s^1 \mathbb{X}_{st}^1 = \sum_{i=1}^d (Z_s^1)_i (\mathbb{X}_{st}^1)^i.$$

We set $F: [0, T] \mapsto \mathcal{T}$

$$F_t := Z_t \mathbb{1} + \sum_{i=1}^d (Z_t^1)_i i.$$

Now we can let the Γ_{st} operators act on F :

$$\Gamma_{ts} F_s = Z_s \mathbb{1} + \sum_{i=1}^d (Z_s^1)_i (i + (\mathbb{X}_{st}^1)^i \mathbb{1})$$

so that

$$F_t - \Gamma_{ts} F_s = (Z_t - Z_s - Z_s^1 \mathbb{X}_{st}^1) \mathbb{1} + \sum_{i=1}^d (Z_t^1 - Z_s^1)_i i.$$

Now the coefficients of this expression satisfy

$$|Z_t - Z_s - Z_s^1 \mathbb{X}_{st}^1| \lesssim |t - s|^{2\alpha}, \quad |Z_t^1 - Z_s^1| \lesssim |t - s|^\alpha.$$

Here we understand the reason for the "strange" definition of Γ_{ts} in terms of $\mathbb{X}_{st}$ instead of $\mathbb{X}_{ts}$: this is the correct definition to obtain the correct expression in the first bound.

Viceversa, given

$$F: [0, T] \rightarrow \mathcal{T}, \quad F_t = f_t^1 \mathbb{1} + \sum_{i=1}^d (f_t^1)_i i,$$

and

$$F_t - \Gamma_{ts} F_s = f_{ts}^1 \mathbb{1} + \sum_{i=1}^d (f_t^1 - f_s^1)_i i$$

the condition

$$|f_{ts}^1| \lesssim |t - s|^{2\alpha}, \quad |f_t^1 - f_s^1| \lesssim |t - s|^\alpha$$

is equivalent to (f, f^1) being a controlled path.

We can note that in this context it would be enough to define $(\Gamma_{st})_{s,t \in [0,T]}$ on the linear span of $\{\mathbb{1}, i: i = 1, \dots, d\}$, which is actually invariant under the action of $(\Gamma_{st})_{s,t \in [0,T]}$.

Let us now consider a controlled path (Z, Z^1) and its integral (I, Z) with respect to $\mathbb{X}$ as in Proposition 9.3. We can now define $U: [0, T] \mapsto \mathcal{T}$

$$U_t := I_t \mathbb{1} + \sum_{i=1}^d (Z_t)^i i + \sum_{i,j=1}^d (Z_t^1)^{ij} (i, j)$$

and compute as for $F: \mathcal{T} \mapsto \mathcal{T}$ above

$$\begin{aligned} U_t - \Gamma_{ts} U_s &= (U_t - U_s - Z_s \mathbb{X}_{st}^1 - Z_s^1 \mathbb{X}_{st}^2) \mathbb{1} \\ &\quad + \sum_{i=1}^d (Z_t - Z_s - Z_s^1 \mathbb{X}_{st}^1)_i i + \sum_{i,j=1}^d (Z_t^1 - Z_s^1)^{ij} (i, j). \end{aligned}$$

CHAPTER 13

NEW TOPICS

- Regularization by noise
- Stochastic sewing lemma
- Machine learning (Rama Cont, rough neural)
- Unbounded rough drivers
- Equazioni riflesse
- Rough Gronwall
- BM in magnetic field
- Parte algebrica, alberi, prodotto tensore, rough equations

BIBLIOGRAPHY

- [1] D. Feyel and A. de La Pradelle, “Curvilinear integrals along enriched paths,” *Electron. J. Probab.* **11** (2006) no. 34, 860–892. <https://doi.org/10.1214/EJP.v11-356>.
- [2] M. Gubinelli, “Controlling Rough Paths,” *J. Funct. Anal.* **216** no. 1, (2004) 86–140, [arXiv:math/0306433](https://arxiv.org/abs/math/0306433) [math.PR].